

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/31, C07K 14/22, A61K 39/095, G01N 33/53, C12Q 1/68, C07K 16/12		A2	(11) International Publication Number: WO 99/36544
			(43) International Publication Date: 22 July 1999 (22.07.99)
(21) International Application Number: PCT/IB99/00103		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 14 January 1999 (14.01.99)			
(30) Priority Data: 9800760.2 14 January 1998 (14.01.98) GB 9819015.0 1 September 1998 (01.09.98) GB 9822143.5 9 October 1998 (09.10.98) GB			
(71) Applicant (for all designated States except US): CHIRON S.P.A. [IT/IT]; Via Fiorentina, 1, I-53100 Siena (IT).			
(72) Inventors; and		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(75) Inventors/Applicants (for US only): MASIGNANI, Vega [IT/IT]; Via Pantaneto, 105, I-53100 Siena (IT). RAP-PUOLI, Rino [IT/IT]; Via delle Rocche, 1, Vagliagli, I-53019 Castelnuovo Berardenga (IT). PIZZA, Mariagrazia [IT/IT]; Strada di Montalbuccio, 160, I-53100 Siena (IT). SCARLATO, Vincenzo [IT/IT]; Via Firenze, 3/37, I-53134 Colle Val d'Elsa (IT). GRANDI, Guido [IT/IT]; 9° Strada, 4, I-20090 Segrate (IT).			
(74) Agent: HALLYBONE, Huw, George; Carpmaels & Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB).			
(54) Title: <i>NEISSERIA MENINGITIDIS</i> ANTIGENS			
(57) Abstract The invention provides proteins from <i>Neisseria meningitidis</i> (strains A and B), including amino acid sequences, the corresponding nucleotide sequences, expression data, and serological data. The proteins are useful antigens for vaccines, immunogenic compositions, and/or diagnostics.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

NEISSERIA MENINGITIDIS ANTIGENS

This invention relates to antigens from the bacterium *Neisseria meningitidis*.

BACKGROUND

Neisseria meningitidis is a non-motile, gram negative diplococcus human pathogen. It colonises the pharynx, causing meningitis and, occasionally, septicaemia in the absence of meningitis. It is closely related to *N.gonorrhoeae*, although one feature that clearly differentiates meningococcus from gonococcus is the presence of a polysaccharide capsule that is present in all pathogenic meningococci.

N.meningitidis causes both endemic and epidemic disease. In the United States the attack rate is 0.6-1 per 100,000 persons per year, and it can be much greater during outbreaks (see Lieberman *et al.* (1996) Safety and Immunogenicity of a Serogroups A/C *Neisseria meningitidis* Oligosaccharide-Protein Conjugate Vaccine in Young Children. *JAMA* 275(19):1499-1503; Schuchat *et al* (1997) Bacterial Meningitis in the United States in 1995. *N Engl J Med* 337(14):970-976). In developing countries, endemic disease rates are much higher and during epidemics incidence rates can reach 500 cases per 100,000 persons per year. Mortality is extremely high, at 10-20% in the United States, and much higher in developing countries. Following the introduction of the conjugate vaccine against *Haemophilus influenzae*, *N. meningitidis* is the major cause of bacterial meningitis at all ages in the United States (Schuchat *et al* (1997) *supra*).

Based on the organism's capsular polysaccharide, 12 serogroups of *N.meningitidis* have been identified. Group A is the pathogen most often implicated in epidemic disease in sub-Saharan Africa. Serogroups B and C are responsible for the vast majority of cases in the United States and in most developed countries. Serogroups W135 and Y are responsible for the rest of the cases in the United States and developed countries. The meningococcal vaccine currently in use is a tetravalent polysaccharide vaccine composed of serogroups A, C, Y and W135. Although efficacious in adolescents and adults, it induces a poor immune response and short duration of protection, and cannot be used in infants [*eg.* Morbidity and Mortality weekly report, Vol.46, No. RR-5 (1997)]. This is because polysaccharides are T-cell independent antigens that induce a weak immune response that cannot be boosted by repeated immunization. Following the success of the

vaccination against *H. influenzae*, conjugate vaccines against serogroups A and C have been developed and are at the final stage of clinical testing (Zollinger WD "New and Improved Vaccines Against Meningococcal Disease" in: *New Generation Vaccines*, *supra*, pp. 469-488; Lieberman *et al* (1996) *supra*; Costantino *et al* (1992) Development and phase I clinical testing of a conjugate vaccine against meningococcus A and C. *Vaccine* 10:691-698).

Meningococcus B remains a problem, however. This serotype currently is responsible for approximately 50% of total meningitis in the United States, Europe, and South America. The polysaccharide approach cannot be used because the menB capsular polysaccharide is a polymer of $\alpha(2-8)$ -linked *N*-acetyl neuraminic acid that is also present in mammalian tissue. This results in tolerance to the antigen; indeed, if an immune response were elicited, it would be anti-self, and therefore undesirable. In order to avoid induction of autoimmunity and to induce a protective immune response, the capsular polysaccharide has, for instance, been chemically modified substituting the *N*-acetyl groups with *N*-propionyl groups, leaving the specific antigenicity unaltered (Romero & Outschoorn (1994) Current status of Meningococcal group B vaccine candidates: capsular or non-capsular? *Clin Microbiol Rev* 7(4):559-575).

Alternative approaches to menB vaccines have used complex mixtures of outer membrane proteins (OMPs), containing either the OMPs alone, or OMPs enriched in porins, or deleted of the class 4 OMPs that are believed to induce antibodies that block bactericidal activity. This approach produces vaccines that are not well characterized. They are able to protect against the homologous strain, but are not effective at large where there are many antigenic variants of the outer membrane proteins. To overcome the antigenic variability, multivalent vaccines containing up to nine different porins have been constructed (*eg.* Poolman JT (1992) Development of a meningococcal vaccine. *Infect. Agents Dis.* 4:13-28). Additional proteins to be used in outer membrane vaccines have been the opa and opc proteins, but none of these approaches have been able to overcome the antigenic variability (*eg.* Ala' Aldeen & Borriello (1996) The meningococcal transferrin-binding proteins 1 and 2 are both surface exposed and generate bactericidal antibodies capable of killing homologous and heterologous strains. *Vaccine* 14(1):49-53).

A certain amount of sequence data is available for meningococcal and gonococcal genes and proteins (*eg.* EP-A-0467714, WO96/29412), but this is by no means complete. The provision of further sequences could provide an opportunity to identify secreted or surface-exposed proteins that

are presumed targets for the immune system and which are not antigenically variable. For instance, some of the identified proteins could be components of efficacious vaccines against meningococcus B, some could be components of vaccines against all meningococcal serotypes, and others could be components of vaccines against all pathogenic *Neisseriae*.

5 THE INVENTION

The invention provides proteins comprising the *N.meningitidis* amino acid sequences disclosed in the examples.

It also provides proteins comprising sequences homologous (*ie.* having sequence identity) to the *N.meningitidis* amino acid sequences disclosed in the examples. Depending on the particular
10 sequence, the degree of sequence identity is preferably greater than 50% (*eg.* 60%, 70%, 80%, 90%, 95%, 99% or more). These homologous proteins include mutants and allelic variants of the sequences disclosed in the examples. Typically, 50% identity or more between two proteins is considered to be an indication of functional equivalence. Identity between the proteins is preferably determined by the Smith-Waterman homology search algorithm as implemented in the MPSRCH
15 program (Oxford Molecular), using an affine gap search with parameters *gap open penalty=12* and *gap extension penalty=1*.

The invention further provides proteins comprising fragments of the *N.meningitidis* amino acid sequences disclosed in the examples. The fragments should comprise at least *n* consecutive amino acids from the sequences and, depending on the particular sequence, *n* is 7 or more (*eg.* 8, 10, 12,
20 14, 16, 18, 20 or more). Preferably the fragments comprise an epitope from the sequence.

The proteins of the invention can, of course, be prepared by various means (*eg.* recombinant expression, purification from cell culture, chemical synthesis *etc.*) and in various forms (*eg.* native, fusions *etc.*). They are preferably prepared in substantially pure form (*ie.* substantially free from other *N.meningitidis* or host cell proteins)

25 According to a further aspect, the invention provides antibodies which bind to these proteins. These may be polyclonal or monoclonal and may be produced by any suitable means.

According to a further aspect, the invention provides nucleic acid comprising the *N.meningitidis* nucleotide sequences disclosed in the examples. In addition, the invention provides nucleic acid comprising sequences homologous (*ie.* having sequence identity) to the *N.meningitidis* nucleotide sequences disclosed in the examples.

- 5 Furthermore, the invention provides nucleic acid which can hybridise to the *N.meningitidis* nucleic acid disclosed in the examples, preferably under "high stringency" conditions (*eg.* 65°C in a 0.1xSSC, 0.5% SDS solution).

- Nucleic acid comprising fragments of these sequences are also provided. These should comprise at least *n* consecutive nucleotides from the *N.meningitidis* sequences and, depending on the particular sequence, *n* is 10 or more (*eg.* 12, 14, 15, 18, 20, 25, 30, 35, 40 or more).
- 10

According to a further aspect, the invention provides nucleic acid encoding the proteins and protein fragments of the invention.

It should also be appreciated that the invention provides nucleic acid comprising sequences complementary to those described above (*eg.* for antisense or probing purposes).

- 15 Nucleic acid according to the invention can, of course, be prepared in many ways (*eg.* by chemical synthesis, from genomic or cDNA libraries, from the organism itself *etc.*) and can take various forms (*eg.* single stranded, double stranded, vectors, probes *etc.*).

In addition, the term "nucleic acid" includes DNA and RNA, and also their analogues, such as those containing modified backbones, and also peptide nucleic acids (PNA) *etc.*

- 20 According to a further aspect, the invention provides vectors comprising nucleotide sequences of the invention (*eg.* expression vectors) and host cells transformed with such vectors.

According to a further aspect, the invention provides compositions comprising protein, antibody, and/or nucleic acid according to the invention. These compositions may be suitable as vaccines, for instance, or as diagnostic reagents, or as immunogenic compositions.

The invention also provides nucleic acid, protein, or antibody according to the invention for use as medicaments (eg. as vaccines) or as diagnostic reagents. It also provides the use of nucleic acid, protein, or antibody according to the invention in the manufacture of: (i) a medicament for treating or preventing infection due to Neisserial bacteria; (ii) a diagnostic reagent for detecting the presence of Neisserial bacteria or of antibodies raised against Neisserial bacteria; and/or (iii) a reagent which can raise antibodies against Neisserial bacteria. Said Neisserial bacteria may be any species or strain (such as *N.gonorrhoeae*) but are preferably *N.meningitidis*, especially strain A, strain B or strain C.

The invention also provides a method of treating a patient, comprising administering to the patient a therapeutically effective amount of nucleic acid, protein, and/or antibody according to the invention.

According to further aspects, the invention provides various processes.

A process for producing proteins of the invention is provided, comprising the step of culturing a host cell according to the invention under conditions which induce protein expression.

A process for producing protein or nucleic acid of the invention is provided, wherein the protein or nucleic acid is synthesised in part or in whole using chemical means.

A process for detecting polynucleotides of the invention is provided, comprising the steps of: (a) contacting a nucleic probe according to the invention with a biological sample under hybridizing conditions to form duplexes; and (b) detecting said duplexes.

A process for detecting proteins of the invention is provided, comprising the steps of: (a) contacting an antibody according to the invention with a biological sample under conditions suitable for the formation of an antibody-antigen complexes; and (b) detecting said complexes.

Unlike the sequences disclosed in PCT/IB98/01665, the sequences disclosed in the present application are believed not to have any significant homologs in *N.gonorrhoeae*. Accordingly, the sequences of the present invention also find use in the preparation of reagents for distinguishing between *N.meningitidis* and *N.gonorrhoeae*

A summary of standard techniques and procedures which may be employed in order to perform the invention (eg. to utilise the disclosed sequences for vaccination or diagnostic purposes) follows. This summary is not a limitation on the invention but, rather, gives examples that may be used, but are not required.

5 General

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature eg. Sambrook *Molecular Cloning; A Laboratory Manual, Second Edition* (1989); *DNA Cloning, Volumes I and*
10 *ii* (D.N. Glover ed. 1985); *Oligonucleotide Synthesis* (M.J. Gait ed, 1984); *Nucleic Acid Hybridization* (B.D. Hames & S.J. Higgins eds. 1984); *Transcription and Translation* (B.D. Hames & S.J. Higgins eds. 1984); *Animal Cell Culture* (R.I. Freshney ed. 1986); *Immobilized Cells and Enzymes* (IRL Press, 1986); B. Perbal, *A Practical Guide to Molecular Cloning* (1984); the *Methods in Enzymology* series (Academic Press, Inc.), especially volumes 154 & 155; *Gene*
15 *Transfer Vectors for Mammalian Cells* (J.H. Miller and M.P. Calos eds. 1987, Cold Spring Harbor Laboratory); Mayer and Walker, eds. (1987), *Immunochemical Methods in Cell and Molecular Biology* (Academic Press, London); Scopes, (1987) *Protein Purification: Principles and Practice*, Second Edition (Springer-Verlag, N.Y.), and *Handbook of Experimental Immunology, Volumes I-IV* (D.M. Weir and C. C. Blackwell eds 1986).

20 Standard abbreviations for nucleotides and amino acids are used in this specification.

All publications, patents, and patent applications cited herein are incorporated in full by reference. In particular, the contents of UK patent applications 9800760.2, 9819015.0 and 9822143.5 are incorporated herein.

Definitions

25 A composition containing X is "substantially free of" Y when at least 85% by weight of the total X+Y in the composition is X. Preferably, X comprises at least about 90% by weight of the total of X+Y in the composition, more preferably at least about 95% or even 99% by weight.

The term "comprising" means "including" as well as "consisting" *eg.* a composition "comprising" X may consist exclusively of X or may include something additional to X, such as X+Y.

The term "heterologous" refers to two biological components that are not found together in nature. The components may be host cells, genes, or regulatory regions, such as promoters. Although the
5 heterologous components are not found together in nature, they can function together, as when a promoter heterologous to a gene is operably linked to the gene. Another example is where a Neisserial sequence is heterologous to a mouse host cell. A further examples would be two epitopes from the same or different proteins which have been assembled in a single protein in an arrangement not found in nature.

10 An "origin of replication" is a polynucleotide sequence that initiates and regulates replication of polynucleotides, such as an expression vector. The origin of replication behaves as an autonomous unit of polynucleotide replication within a cell, capable of replication under its own control. An origin of replication may be needed for a vector to replicate in a particular host cell. With certain
15 origins of replication, an expression vector can be reproduced at a high copy number in the presence of the appropriate proteins within the cell. Examples of origins are the autonomously replicating sequences, which are effective in yeast; and the viral T-antigen, effective in COS-7 cells.

A "mutant" sequence is defined as DNA, RNA or amino acid sequence differing from but having sequence identity with the native or disclosed sequence. Depending on the particular sequence, the
20 degree of sequence identity between the native or disclosed sequence and the mutant sequence is preferably greater than 50% (*eg.* 60%, 70%, 80%, 90%, 95%, 99% or more, calculated using the Smith-Waterman algorithm as described above). As used herein, an "allelic variant" of a nucleic acid molecule, or region, for which nucleic acid sequence is provided herein is a nucleic acid molecule, or region, that occurs essentially at the same locus in the genome of another or second
25 isolate, and that, due to natural variation caused by, for example, mutation or recombination, has a similar but not identical nucleic acid sequence. A coding region allelic variant typically encodes a protein having similar activity to that of the protein encoded by the gene to which it is being compared. An allelic variant can also comprise an alteration in the 5' or 3' untranslated regions of the gene, such as in regulatory control regions (*eg.* see US patent 5,753,235).

Expression systems

The Neisserial nucleotide sequences can be expressed in a variety of different expression systems; for example those used with mammalian cells, baculoviruses, plants, bacteria, and yeast.

i. Mammalian Systems

5 Mammalian expression systems are known in the art. A mammalian promoter is any DNA sequence capable of binding mammalian RNA polymerase and initiating the downstream (3') transcription of a coding sequence (eg. structural gene) into mRNA. A promoter will have a transcription initiating region, which is usually placed proximal to the 5' end of the coding sequence, and a TATA box, usually located 25-30 base pairs (bp) upstream of the transcription
10 initiation site. The TATA box is thought to direct RNA polymerase II to begin RNA synthesis at the correct site. A mammalian promoter will also contain an upstream promoter element, usually located within 100 to 200 bp upstream of the TATA box. An upstream promoter element determines the rate at which transcription is initiated and can act in either orientation [Sambrook et al. (1989) "Expression of Cloned Genes in Mammalian Cells." In *Molecular Cloning: A*
15 *Laboratory Manual*, 2nd ed.].

Mammalian viral genes are often highly expressed and have a broad host range; therefore sequences encoding mammalian viral genes provide particularly useful promoter sequences. Examples include the SV40 early promoter, mouse mammary tumor virus LTR promoter, adenovirus major late promoter (Ad MLP), and herpes simplex virus promoter. In addition, sequences derived from non-
20 viral genes, such as the murine metallothionein gene, also provide useful promoter sequences. Expression may be either constitutive or regulated (inducible), depending on the promoter can be induced with glucocorticoid in hormone-responsive cells.

The presence of an enhancer element (enhancer), combined with the promoter elements described above, will usually increase expression levels. An enhancer is a regulatory DNA sequence that can
25 stimulate transcription up to 1000-fold when linked to homologous or heterologous promoters, with synthesis beginning at the normal RNA start site. Enhancers are also active when they are placed upstream or downstream from the transcription initiation site, in either normal or flipped orientation, or at a distance of more than 1000 nucleotides from the promoter [Maniatis et al. (1987) *Science* 236:1237; Alberts et al. (1989) *Molecular Biology of the Cell*, 2nd ed.]. Enhancer elements
30 derived from viruses may be particularly useful, because they usually have a broader host range.

Examples include the SV40 early gene enhancer [Dijkema et al (1985) *EMBO J.* 4:761] and the enhancer/promoters derived from the long terminal repeat (LTR) of the Rous Sarcoma Virus [Gorman et al. (1982b) *Proc. Natl. Acad. Sci.* 79:6777] and from human cytomegalovirus [Boshart et al. (1985) *Cell* 41:521]. Additionally, some enhancers are regulatable and become active only
5 in the presence of an inducer, such as a hormone or metal ion [Sassone-Corsi and Borelli (1986) *Trends Genet.* 2:215; Maniatis et al. (1987) *Science* 236:1237].

A DNA molecule may be expressed intracellularly in mammalian cells. A promoter sequence may be directly linked with the DNA molecule, in which case the first amino acid at the N-terminus of the recombinant protein will always be a methionine, which is encoded by the ATG start codon. If desired,
10 the N-terminus may be cleaved from the protein by *in vitro* incubation with cyanogen bromide.

Alternatively, foreign proteins can also be secreted from the cell into the growth media by creating chimeric DNA molecules that encode a fusion protein comprised of a leader sequence fragment that provides for secretion of the foreign protein in mammalian cells. Preferably, there are processing sites encoded between the leader fragment and the foreign gene that can be cleaved either *in vivo*
15 or *in vitro*. The leader sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the secretion of the protein from the cell. The adenovirus tripartite leader is an example of a leader sequence that provides for secretion of a foreign protein in mammalian cells.

Usually, transcription termination and polyadenylation sequences recognized by mammalian cells
20 are regulatory regions located 3' to the translation stop codon and thus, together with the promoter elements, flank the coding sequence. The 3' terminus of the mature mRNA is formed by site-specific post-transcriptional cleavage and polyadenylation [Birnstiel et al. (1985) *Cell* 41:349; Proudfoot and Whitelaw (1988) "Termination and 3' end processing of eukaryotic RNA. In *Transcription and splicing* (ed. B.D. Hames and D.M. Glover); Proudfoot (1989) *Trends Biochem.*
25 *Sci.* 14:105]. These sequences direct the transcription of an mRNA which can be translated into the polypeptide encoded by the DNA. Examples of transcription terminator/polyadenylation signals include those derived from SV40 [Sambrook et al (1989) "Expression of cloned genes in cultured mammalian cells." In *Molecular Cloning: A Laboratory Manual*].

Usually, the above described components, comprising a promoter, polyadenylation signal, and transcription termination sequence are put together into expression constructs. Enhancers, introns with functional splice donor and acceptor sites, and leader sequences may also be included in an expression construct, if desired. Expression constructs are often maintained in a replicon, such as an extrachromosomal element (eg. plasmids) capable of stable maintenance in a host, such as mammalian cells or bacteria. Mammalian replication systems include those derived from animal viruses, which require trans-acting factors to replicate. For example, plasmids containing the replication systems of papovaviruses, such as SV40 [Gluzman (1981) *Cell* 23:175] or polyomavirus, replicate to extremely high copy number in the presence of the appropriate viral T antigen. Additional examples of mammalian replicons include those derived from bovine papillomavirus and Epstein-Barr virus. Additionally, the replicon may have two replicaton systems, thus allowing it to be maintained, for example, in mammalian cells for expression and in a prokaryotic host for cloning and amplification. Examples of such mammalian-bacteria shuttle vectors include pMT2 [Kaufman et al. (1989) *Mol. Cell. Biol.* 9:946] and pHEBO [Shimizu et al. (1986) *Mol. Cell. Biol.* 6:1074].

The transformation procedure used depends upon the host to be transformed. Methods for introduction of heterologous polynucleotides into mammalian cells are known in the art and include dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei.

Mammalian cell lines available as hosts for expression are known in the art and include many immortalized cell lines available from the American Type Culture Collection (ATCC), including but not limited to, Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (eg. Hep G2), and a number of other cell lines.

ii. Baculovirus Systems

The polynucleotide encoding the protein can also be inserted into a suitable insect expression vector, and is operably linked to the control elements within that vector. Vector construction employs techniques which are known in the art. Generally, the components of the expression system include a transfer vector, usually a bacterial plasmid, which contains both a fragment of the baculovirus

genome, and a convenient restriction site for insertion of the heterologous gene or genes to be expressed; a wild type baculovirus with a sequence homologous to the baculovirus-specific fragment in the transfer vector (this allows for the homologous recombination of the heterologous gene in to the baculovirus genome); and appropriate insect host cells and growth media.

- 5 After inserting the DNA sequence encoding the protein into the transfer vector, the vector and the wild type viral genome are transfected into an insect host cell where the vector and viral genome are allowed to recombine. The packaged recombinant virus is expressed and recombinant plaques are identified and purified. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, *inter alia*, Invitrogen, San Diego CA ("MaxBac" kit).
- 10 These techniques are generally known to those skilled in the art and fully described in Summers and Smith, *Texas Agricultural Experiment Station Bulletin No. 1555* (1987) (hereinafter "Summers and Smith").

- Prior to inserting the DNA sequence encoding the protein into the baculovirus genome, the above described components, comprising a promoter, leader (if desired), coding sequence of interest, and
- 15 transcription termination sequence, are usually assembled into an intermediate transplacement construct (transfer vector). This construct may contain a single gene and operably linked regulatory elements; multiple genes, each with its owned set of operably linked regulatory elements; or multiple genes, regulated by the same set of regulatory elements. Intermediate transplacement constructs are often maintained in a replicon, such as an extrachromosomal element (*eg.* plasmids) capable of stable
- 20 maintenance in a host, such as a bacterium. The replicon will have a replication system, thus allowing it to be maintained in a suitable host for cloning and amplification.

- Currently, the most commonly used transfer vector for introducing foreign genes into AcNPV is pAc373. Many other vectors, known to those of skill in the art, have also been designed. These include, for example, pVL985 (which alters the polyhedrin start codon from ATG to ATT, and
- 25 which introduces a BamHI cloning site 32 basepairs downstream from the ATT; see Luckow and Summers, *Virology* (1989) 17:31.

The plasmid usually also contains the polyhedrin polyadenylation signal (Miller et al. (1988) *Ann. Rev. Microbiol.*, 42:177) and a prokaryotic ampicillin-resistance (*amp*) gene and origin of replication for selection and propagation in *E. coli*.

Baculovirus transfer vectors usually contain a baculovirus promoter. A baculovirus promoter is any DNA sequence capable of binding a baculovirus RNA polymerase and initiating the downstream (5' to 3') transcription of a coding sequence (eg. structural gene) into mRNA. A promoter will have a transcription initiation region which is usually placed proximal to the 5' end of the coding sequence. This transcription initiation region usually includes an RNA polymerase binding site and a transcription initiation site. A baculovirus transfer vector may also have a second domain called an enhancer, which, if present, is usually distal to the structural gene. Expression may be either regulated or constitutive.

Structural genes, abundantly transcribed at late times in a viral infection cycle, provide particularly useful promoter sequences. Examples include sequences derived from the gene encoding the viral polyhedron protein, Friesen et al., (1986) "The Regulation of Baculovirus Gene Expression," in: *The Molecular Biology of Baculoviruses* (ed. Walter Doerfler); EPO Publ. Nos. 127 839 and 155 476; and the gene encoding the p10 protein, Vlak et al., (1988), *J. Gen. Virol.* 69:765.

DNA encoding suitable signal sequences can be derived from genes for secreted insect or baculovirus proteins, such as the baculovirus polyhedrin gene (Carbonell et al. (1988) *Gene*, 73:409). Alternatively, since the signals for mammalian cell posttranslational modifications (such as signal peptide cleavage, proteolytic cleavage, and phosphorylation) appear to be recognized by insect cells, and the signals required for secretion and nuclear accumulation also appear to be conserved between the invertebrate cells and vertebrate cells, leaders of non-insect origin, such as those derived from genes encoding human α -interferon, Maeda et al., (1985), *Nature* 315:592; human gastrin-releasing peptide, Lebacqz-Verheyden et al., (1988), *Molec. Cell. Biol.* 8:3129; human IL-2, Smith et al., (1985) *Proc. Nat'l Acad. Sci. USA*, 82:8404; mouse IL-3, (Miyajima et al., (1987) *Gene* 58:273; and human glucocerebrosidase, Martin et al. (1988) *DNA*, 7:99, can also be used to provide for secretion in insects.

A recombinant polypeptide or polyprotein may be expressed intracellularly or, if it is expressed with the proper regulatory sequences, it can be secreted. Good intracellular expression of nonfused foreign proteins usually requires heterologous genes that ideally have a short leader sequence containing suitable translation initiation signals preceding an ATG start signal. If desired, methionine at the N-terminus may be cleaved from the mature protein by *in vitro* incubation with cyanogen bromide.

Alternatively, recombinant polyproteins or proteins which are not naturally secreted can be secreted from the insect cell by creating chimeric DNA molecules that encode a fusion protein comprised of a leader sequence fragment that provides for secretion of the foreign protein in insects. The leader sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the translocation of the protein into the endoplasmic reticulum.

After insertion of the DNA sequence and/or the gene encoding the expression product precursor of the protein, an insect cell host is co-transformed with the heterologous DNA of the transfer vector and the genomic DNA of wild type baculovirus -- usually by co-transfection. The promoter and transcription termination sequence of the construct will usually comprise a 2-5kb section of the baculovirus genome. Methods for introducing heterologous DNA into the desired site in the baculovirus virus are known in the art. (See Summers and Smith *supra*; Ju et al. (1987); Smith et al., *Mol. Cell. Biol.* (1983) 3:2156; and Luckow and Summers (1989)). For example, the insertion can be into a gene such as the polyhedrin gene, by homologous double crossover recombination; insertion can also be into a restriction enzyme site engineered into the desired baculovirus gene. Miller et al., (1989), *Bioessays* 4:91. The DNA sequence, when cloned in place of the polyhedrin gene in the expression vector, is flanked both 5' and 3' by polyhedrin-specific sequences and is positioned downstream of the polyhedrin promoter.

The newly formed baculovirus expression vector is subsequently packaged into an infectious recombinant baculovirus. Homologous recombination occurs at low frequency (between about 1% and about 5%); thus, the majority of the virus produced after cotransfection is still wild-type virus. Therefore, a method is necessary to identify recombinant viruses. An advantage of the expression system is a visual screen allowing recombinant viruses to be distinguished. The polyhedrin protein, which is produced by the native virus, is produced at very high levels in the nuclei of infected cells at late times after viral infection. Accumulated polyhedrin protein forms occlusion bodies that also contain embedded particles. These occlusion bodies, up to 15 μ m in size, are highly refractile, giving them a bright shiny appearance that is readily visualized under the light microscope. Cells infected with recombinant viruses lack occlusion bodies. To distinguish recombinant virus from wild-type virus, the transfection supernatant is plaqued onto a monolayer of insect cells by techniques known to those skilled in the art. Namely, the plaques are screened under the light microscope for the presence (indicative of wild-type virus) or absence (indicative of recombinant

virus) of occlusion bodies. "Current Protocols in Microbiology" Vol. 2 (Ausubel et al. eds) at 16.8 (Supp. 10, 1990); Summers and Smith, *supra*; Miller et al. (1989).

Recombinant baculovirus expression vectors have been developed for infection into several insect cells. For example, recombinant baculoviruses have been developed for, *inter alia*: *Aedes aegypti*,
5 *Autographa californica*, *Bombyx mori*, *Drosophila melanogaster*, *Spodoptera frugiperda*, and *Trichoplusia ni* (WO 89/046699; Carbonell et al., (1985) *J. Virol.* 56:153; Wright (1986) *Nature* 321:718; Smith et al., (1983) *Mol. Cell. Biol.* 3:2156; and see generally, Fraser, *et al.* (1989) *In Vitro Cell. Dev. Biol.* 25:225).

Cells and cell culture media are commercially available for both direct and fusion expression of
10 heterologous polypeptides in a baculovirus/expression system; cell culture technology is generally known to those skilled in the art. *See, eg.* Summers and Smith *supra*.

The modified insect cells may then be grown in an appropriate nutrient medium, which allows for stable maintenance of the plasmid(s) present in the modified insect host. Where the expression product gene is under inducible control, the host may be grown to high density, and expression induced.
15 Alternatively, where expression is constitutive, the product will be continuously expressed into the medium and the nutrient medium must be continuously circulated, while removing the product of interest and augmenting depleted nutrients. The product may be purified by such techniques as chromatography, *eg.* HPLC, affinity chromatography, ion exchange chromatography, etc.; electrophoresis; density gradient centrifugation; solvent extraction, or the like. As appropriate, the
20 product may be further purified, as required, so as to remove substantially any insect proteins which are also secreted in the medium or result from lysis of insect cells, so as to provide a product which is at least substantially free of host debris, *eg.* proteins, lipids and polysaccharides.

In order to obtain protein expression, recombinant host cells derived from the transformants are incubated under conditions which allow expression of the recombinant protein encoding sequence.
25 These conditions will vary, dependent upon the host cell selected. However, the conditions are readily ascertainable to those of ordinary skill in the art, based upon what is known in the art.

iii. Plant Systems

There are many plant cell culture and whole plant genetic expression systems known in the art. Exemplary plant cellular genetic expression systems include those described in patents, such as:

US 5,693,506; US 5,659,122; and US 5,608,143. Additional examples of genetic expression in plant cell culture has been described by Zenk, *Phytochemistry* 30:3861-3863 (1991). Descriptions of plant protein signal peptides may be found in addition to the references described above in Vaulcombe et al., *Mol. Gen. Genet.* 209:33-40 (1987); Chandler et al., *Plant Molecular Biology* 3:407-418 (1984); Rogers, *J. Biol. Chem.* 260:3731-3738 (1985); Rothstein et al., *Gene* 55:353-356 (1987); Whittier et al., *Nucleic Acids Research* 15:2515-2535 (1987); Wirsal et al., *Molecular Microbiology* 3:3-14 (1989); Yu et al., *Gene* 122:247-253 (1992). A description of the regulation of plant gene expression by the phytohormone, gibberellic acid and secreted enzymes induced by gibberellic acid can be found in R.L. Jones and J. MacMillin, *Gibberellins*: in: *Advanced Plant Physiology*, Malcolm B. Wilkins, ed., 1984 Pitman Publishing Limited, London, pp. 21-52. References that describe other metabolically-regulated genes: Sheen, *Plant Cell*, 2:1027-1038(1990); Maas et al., *EMBO J.* 9:3447-3452 (1990); Benkel and Hickey, *Proc. Natl. Acad. Sci.* 84:1337-1339 (1987)

Typically, using techniques known in the art, a desired polynucleotide sequence is inserted into an expression cassette comprising genetic regulatory elements designed for operation in plants. The expression cassette is inserted into a desired expression vector with companion sequences upstream and downstream from the expression cassette suitable for expression in a plant host. The companion sequences will be of plasmid or viral origin and provide necessary characteristics to the vector to permit the vectors to move DNA from an original cloning host, such as bacteria, to the desired plant host. The basic bacterial/plant vector construct will preferably provide a broad host range prokaryote replication origin; a prokaryote selectable marker; and, for *Agrobacterium* transformations, T DNA sequences for *Agrobacterium*-mediated transfer to plant chromosomes. Where the heterologous gene is not readily amenable to detection, the construct will preferably also have a selectable marker gene suitable for determining if a plant cell has been transformed. A general review of suitable markers, for example for the members of the grass family, is found in Wilmink and Dons, 1993, *Plant Mol. Biol. Repr.* 11(2):165-185.

Sequences suitable for permitting integration of the heterologous sequence into the plant genome are also recommended. These might include transposon sequences and the like for homologous recombination as well as Ti sequences which permit random insertion of a heterologous expression cassette into a plant genome. Suitable prokaryote selectable markers include resistance toward

antibiotics such as ampicillin or tetracycline. Other DNA sequences encoding additional functions may also be present in the vector, as is known in the art.

The nucleic acid molecules of the subject invention may be included into an expression cassette for expression of the protein(s) of interest. Usually, there will be only one expression cassette, although two or more are feasible. The recombinant expression cassette will contain in addition to the heterologous protein encoding sequence the following elements, a promoter region, plant 5' untranslated sequences, initiation codon depending upon whether or not the structural gene comes equipped with one, and a transcription and translation termination sequence. Unique restriction enzyme sites at the 5' and 3' ends of the cassette allow for easy insertion into a pre-existing vector.

10 A heterologous coding sequence may be for any protein relating to the present invention. The sequence encoding the protein of interest will encode a signal peptide which allows processing and translocation of the protein, as appropriate, and will usually lack any sequence which might result in the binding of the desired protein of the invention to a membrane. Since, for the most part, the transcriptional initiation region will be for a gene which is expressed and translocated during
15 germination, by employing the signal peptide which provides for translocation, one may also provide for translocation of the protein of interest. In this way, the protein(s) of interest will be translocated from the cells in which they are expressed and may be efficiently harvested. Typically secretion in seeds are across the aleurone or scutellar epithelium layer into the endosperm of the seed. While it is not required that the protein be secreted from the cells in which the protein is
20 produced, this facilitates the isolation and purification of the recombinant protein.

Since the ultimate expression of the desired gene product will be in a eucaryotic cell it is desirable to determine whether any portion of the cloned gene contains sequences which will be processed out as introns by the host's splicosome machinery. If so, site-directed mutagenesis of the "intron" region may be conducted to prevent losing a portion of the genetic message as a false intron code,
25 Reed and Maniatis, *Cell* 41:95-105, 1985.

The vector can be microinjected directly into plant cells by use of micropipettes to mechanically transfer the recombinant DNA. Crossway, *Mol. Gen. Genet*, 202:179-185, 1985. The genetic material may also be transferred into the plant cell by using polyethylene glycol, Krens, et al., *Nature*, 296, 72-74, 1982. Another method of introduction of nucleic acid segments is high

velocity ballistic penetration by small particles with the nucleic acid either within the matrix of small beads or particles, or on the surface, Klein, et al., *Nature*, 327, 70-73, 1987 and Knudsen and Muller, 1991, *Planta*, 185:330-336 teaching particle bombardment of barley endosperm to create transgenic barley. Yet another method of introduction would be fusion of protoplasts with other entities, either minicells, cells, lysosomes or other fusible lipid-surfaced bodies, Fraley, et al., *Proc. Natl. Acad. Sci. USA*, 79, 1859-1863, 1982.

The vector may also be introduced into the plant cells by electroporation. (Fromm et al., *Proc. Natl. Acad. Sci. USA* 82:5824, 1985). In this technique, plant protoplasts are electroporated in the presence of plasmids containing the gene construct. Electrical impulses of high field strength reversibly permeabilize biomembranes allowing the introduction of the plasmids. Electroporated plant protoplasts reform the cell wall, divide, and form plant callus.

All plants from which protoplasts can be isolated and cultured to give whole regenerated plants can be transformed by the present invention so that whole plants are recovered which contain the transferred gene. It is known that practically all plants can be regenerated from cultured cells or tissues, including but not limited to all major species of sugarcane, sugar beet, cotton, fruit and other trees, legumes and vegetables. Some suitable plants include, for example, species from the genera *Fragaria*, *Lotus*, *Medicago*, *Onobrychis*, *Trifolium*, *Trigonella*, *Vigna*, *Citrus*, *Linum*, *Geranium*, *Manihot*, *Daucus*, *Arabidopsis*, *Brassica*, *Raphanus*, *Sinapis*, *Atropa*, *Capsicum*, *Datura*, *Hyoscyamus*, *Lycopersion*, *Nicotiana*, *Solanum*, *Petunia*, *Digitalis*, *Majorana*, *Cichorium*, *Helianthus*, *Lactuca*, *Bromus*, *Asparagus*, *Antirrhinum*, *Hererocallis*, *Nemesia*, *Pelargonium*, *Panicum*, *Pennisetum*, *Ramunculus*, *Senecio*, *Salpiglossis*, *Cucumis*, *Browaalia*, *Glycine*, *Lolium*, *Zea*, *Triticum*, *Sorghum*, and *Datura*.

Means for regeneration vary from species to species of plants, but generally a suspension of transformed protoplasts containing copies of the heterologous gene is first provided. Callus tissue is formed and shoots may be induced from callus and subsequently rooted. Alternatively, embryo formation can be induced from the protoplast suspension. These embryos germinate as natural embryos to form plants. The culture media will generally contain various amino acids and hormones, such as auxin and cytokinins. It is also advantageous to add glutamic acid and proline to the medium, especially for such species as corn and alfalfa. Shoots and roots normally develop simultaneously. Efficient regeneration will depend on the medium, on the genotype, and on the

history of the culture. If these three variables are controlled, then regeneration is fully reproducible and repeatable.

In some plant cell culture systems, the desired protein of the invention may be excreted or alternatively, the protein may be extracted from the whole plant. Where the desired protein of the invention is secreted into the medium, it may be collected. Alternatively, the embryos and embryoless-half seeds or other plant tissue may be mechanically disrupted to release any secreted protein between cells and tissues. The mixture may be suspended in a buffer solution to retrieve soluble proteins. Conventional protein isolation and purification methods will be then used to purify the recombinant protein. Parameters of time, temperature pH, oxygen, and volumes will be adjusted through routine methods to optimize expression and recovery of heterologous protein.

iv. Bacterial Systems

Bacterial expression techniques are known in the art. A bacterial promoter is any DNA sequence capable of binding bacterial RNA polymerase and initiating the downstream (3') transcription of a coding sequence (eg. structural gene) into mRNA. A promoter will have a transcription initiation region which is usually placed proximal to the 5' end of the coding sequence. This transcription initiation region usually includes an RNA polymerase binding site and a transcription initiation site. A bacterial promoter may also have a second domain called an operator, that may overlap an adjacent RNA polymerase binding site at which RNA synthesis begins. The operator permits negative regulated (inducible) transcription, as a gene repressor protein may bind the operator and thereby inhibit transcription of a specific gene. Constitutive expression may occur in the absence of negative regulatory elements, such as the operator. In addition, positive regulation may be achieved by a gene activator protein binding sequence, which, if present is usually proximal (5') to the RNA polymerase binding sequence. An example of a gene activator protein is the catabolite activator protein (CAP), which helps initiate transcription of the lac operon in *Escherichia coli* (*E. coli*) [Raibaud *et al.* (1984) *Annu. Rev. Genet.* 18:173]. Regulated expression may therefore be either positive or negative, thereby either enhancing or reducing transcription.

Sequences encoding metabolic pathway enzymes provide particularly useful promoter sequences. Examples include promoter sequences derived from sugar metabolizing enzymes, such as galactose, lactose (*lac*) [Chang *et al.* (1977) *Nature* 198:1056], and maltose. Additional examples include promoter sequences derived from biosynthetic enzymes such as tryptophan (*trp*) [Goeddel *et al.*

(1980) *Nuc. Acids Res.* 8:4057; Yelverton *et al.* (1981) *Nucl. Acids Res.* 9:731; US patent 4,738,921; EP-A-0036776 and EP-A-0121775]. The g-laotamase (*bla*) promoter system [Weissmann (1981) "The cloning of interferon and other mistakes." In *Interferon 3* (ed. I. Gresser)], bacteriophage lambda PL [Shimatake *et al.* (1981) *Nature* 292:128] and T5 [US patent 4,689,406]
5 promoter systems also provide useful promoter sequences.

In addition, synthetic promoters which do not occur in nature also function as bacterial promoters. For example, transcription activation sequences of one bacterial or bacteriophage promoter may be joined with the operon sequences of another bacterial or bacteriophage promoter, creating a synthetic hybrid promoter [US patent 4,551,433]. For example, the *tac* promoter is a hybrid *trp-lac*
10 promoter comprised of both *trp* promoter and *lac* operon sequences that is regulated by the *lac* repressor [Amann *et al.* (1983) *Gene* 25:167; de Boer *et al.* (1983) *Proc. Natl. Acad. Sci.* 80:21]. Furthermore, a bacterial promoter can include naturally occurring promoters of non-bacterial origin that have the ability to bind bacterial RNA polymerase and initiate transcription. A naturally occurring promoter of non-bacterial origin can also be coupled with a compatible RNA polymerase
15 to produce high levels of expression of some genes in prokaryotes. The bacteriophage T7 RNA polymerase/promoter system is an example of a coupled promoter system [Studier *et al.* (1986) *J. Mol. Biol.* 189:113; Tabor *et al.* (1985) *Proc Natl. Acad. Sci.* 82:1074]. In addition, a hybrid promoter can also be comprised of a bacteriophage promoter and an *E. coli* operator region (EPO-A-0 267 851).

20 In addition to a functioning promoter sequence, an efficient ribosome binding site is also useful for the expression of foreign genes in prokaryotes. In *E. coli*, the ribosome binding site is called the Shine-Dalgarno (SD) sequence and includes an initiation codon (ATG) and a sequence 3-9 nucleotides in length located 3-11 nucleotides upstream of the initiation codon [Shine *et al.* (1975) *Nature* 254:34]. The SD sequence is thought to promote binding of mRNA to the ribosome by the
25 pairing of bases between the SD sequence and the 3' end of *E. coli* 16S rRNA [Steitz *et al.* (1979) "Genetic signals and nucleotide sequences in messenger RNA." In *Biological Regulation and Development: Gene Expression* (ed. R.F. Goldberger)]. To express eukaryotic genes and prokaryotic genes with weak ribosome-binding site [Sambrook *et al.* (1989) "Expression of cloned genes in Escherichia coli." In *Molecular Cloning: A Laboratory Manual*].

A DNA molecule may be expressed intracellularly. A promoter sequence may be directly linked with the DNA molecule, in which case the first amino acid at the N-terminus will always be a methionine, which is encoded by the ATG start codon. If desired, methionine at the N-terminus may be cleaved from the protein by *in vitro* incubation with cyanogen bromide or by either *in vivo* on *in vitro* incubation with a bacterial methionine N-terminal peptidase (EPO-A-0 219 237).

Fusion proteins provide an alternative to direct expression. Usually, a DNA sequence encoding the N-terminal portion of an endogenous bacterial protein, or other stable protein, is fused to the 5' end of heterologous coding sequences. Upon expression, this construct will provide a fusion of the two amino acid sequences. For example, the bacteriophage lambda cell gene can be linked at the 5' terminus of a foreign gene and expressed in bacteria. The resulting fusion protein preferably retains a site for a processing enzyme (factor Xa) to cleave the bacteriophage protein from the foreign gene [Nagai *et al.* (1984) *Nature* 309:810]. Fusion proteins can also be made with sequences from the *lacZ* [Jia *et al.* (1987) *Gene* 60:197], *trpE* [Allen *et al.* (1987) *J. Biotechnol.* 5:93; Makoff *et al.* (1989) *J. Gen. Microbiol.* 135:11], and *Chey* [EP-A-0 324 647] genes. The DNA sequence at the junction of the two amino acid sequences may or may not encode a cleavable site. Another example is a ubiquitin fusion protein. Such a fusion protein is made with the ubiquitin region that preferably retains a site for a processing enzyme (*eg.* ubiquitin specific processing-protease) to cleave the ubiquitin from the foreign protein. Through this method, native foreign protein can be isolated [Miller *et al.* (1989) *Bio/Technology* 7:698].

Alternatively, foreign proteins can also be secreted from the cell by creating chimeric DNA molecules that encode a fusion protein comprised of a signal peptide sequence fragment that provides for secretion of the foreign protein in bacteria [US patent 4,336,336]. The signal sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the secretion of the protein from the cell. The protein is either secreted into the growth media (gram-positive bacteria) or into the periplasmic space, located between the inner and outer membrane of the cell (gram-negative bacteria). Preferably there are processing sites, which can be cleaved either *in vivo* or *in vitro* encoded between the signal peptide fragment and the foreign gene.

DNA encoding suitable signal sequences can be derived from genes for secreted bacterial proteins, such as the *E. coli* outer membrane protein gene (*ompA*) [Masui *et al.* (1983), in: *Experimental Manipulation of Gene Expression*; Ghrayeb *et al.* (1984) *EMBO J.* 3:2437] and the *E. coli* alkaline

phosphatase signal sequence (*phoA*) [Oka *et al.* (1985) *Proc. Natl. Acad. Sci.* 82:7212]. As an additional example, the signal sequence of the alpha-amylase gene from various *Bacillus* strains can be used to secrete heterologous proteins from *B. subtilis* [Palva *et al.* (1982) *Proc. Natl. Acad. Sci. USA* 79:5582; EP-A-0 244 042].

- 5 Usually, transcription termination sequences recognized by bacteria are regulatory regions located 3' to the translation stop codon, and thus together with the promoter flank the coding sequence. These sequences direct the transcription of an mRNA which can be translated into the polypeptide encoded by the DNA. Transcription termination sequences frequently include DNA sequences of about 50 nucleotides capable of forming stem loop structures that aid in terminating transcription.
- 10 Examples include transcription termination sequences derived from genes with strong promoters, such as the *trp* gene in *E. coli* as well as other biosynthetic genes.

- Usually, the above described components, comprising a promoter, signal sequence (if desired), coding sequence of interest, and transcription termination sequence, are put together into expression constructs. Expression constructs are often maintained in a replicon, such as an extrachromosomal
- 15 element (*eg.* plasmids) capable of stable maintenance in a host, such as bacteria. The replicon will have a replication system, thus allowing it to be maintained in a prokaryotic host either for expression or for cloning and amplification. In addition, a replicon may be either a high or low copy number plasmid. A high copy number plasmid will generally have a copy number ranging from about 5 to about 200, and usually about 10 to about 150. A host containing a high copy
- 20 number plasmid will preferably contain at least about 10, and more preferably at least about 20 plasmids. Either a high or low copy number vector may be selected, depending upon the effect of the vector and the foreign protein on the host.

- Alternatively, the expression constructs can be integrated into the bacterial genome with an integrating vector. Integrating vectors usually contain at least one sequence homologous to the
- 25 bacterial chromosome that allows the vector to integrate. Integrations appear to result from recombinations between homologous DNA in the vector and the bacterial chromosome. For example, integrating vectors constructed with DNA from various *Bacillus* strains integrate into the *Bacillus* chromosome (EP-A- 0 127 328). Integrating vectors may also be comprised of bacteriophage or transposon sequences.

Usually, extrachromosomal and integrating expression constructs may contain selectable markers to allow for the selection of bacterial strains that have been transformed. Selectable markers can be expressed in the bacterial host and may include genes which render bacteria resistant to drugs such as ampicillin, chloramphenicol, erythromycin, kanamycin (neomycin), and tetracycline [Davies *et al.* (1978) *Annu. Rev. Microbiol.* 32:469]. Selectable markers may also include biosynthetic genes, such as those in the histidine, tryptophan, and leucine biosynthetic pathways.

Alternatively, some of the above described components can be put together in transformation vectors. Transformation vectors are usually comprised of a selectable market that is either maintained in a replicon or developed into an integrating vector, as described above.

Expression and transformation vectors, either extra-chromosomal replicons or integrating vectors, have been developed for transformation into many bacteria. For example, expression vectors have been developed for, *inter alia*, the following bacteria: *Bacillus subtilis* [Palva *et al.* (1982) *Proc. Natl. Acad. Sci. USA* 79:5582; EP-A-0 036 259 and EP-A-0 063 953; WO 84/04541], *Escherichia coli* [Shimatake *et al.* (1981) *Nature* 292:128; Amann *et al.* (1985) *Gene* 40:183; Studier *et al.* (1986) *J. Mol. Biol.* 189:113; EP-A-0 036 776, EP-A-0 136 829 and EP-A-0 136 907], *Streptococcus cremoris* [Powell *et al.* (1988) *Appl. Environ. Microbiol.* 54:655]; *Streptococcus lividans* [Powell *et al.* (1988) *Appl. Environ. Microbiol.* 54:655], *Streptomyces lividans* [US patent 4,745,056].

Methods of introducing exogenous DNA into bacterial hosts are well-known in the art, and usually include either the transformation of bacteria treated with CaCl_2 or other agents, such as divalent cations and DMSO. DNA can also be introduced into bacterial cells by electroporation. Transformation procedures usually vary with the bacterial species to be transformed. See *eg.* [Masson *et al.* (1989) *FEMS Microbiol. Lett.* 60:273; Palva *et al.* (1982) *Proc. Natl. Acad. Sci. USA* 79:5582; EP-A-0 036 259 and EP-A-0 063 953; WO 84/04541, *Bacillus*], [Miller *et al.* (1988) *Proc. Natl. Acad. Sci.* 85:856; Wang *et al.* (1990) *J. Bacteriol.* 172:949, *Campylobacter*], [Cohen *et al.* (1973) *Proc. Natl. Acad. Sci.* 69:2110; Dower *et al.* (1988) *Nucleic Acids Res.* 16:6127; Kushner (1978) "An improved method for transformation of *Escherichia coli* with ColE1-derived plasmids. In *Genetic Engineering: Proceedings of the International Symposium on Genetic Engineering* (eds. H.W. Boyer and S. Nicosia); Mandel *et al.* (1970) *J. Mol. Biol.* 53:159; Taketo (1988) *Biochim. Biophys. Acta* 949:318; *Escherichia*], [Chassy *et al.* (1987) *FEMS Microbiol. Lett.*

- 44:173 *Lactobacillus*]; [Fiedler *et al.* (1988) *Anal. Biochem* 170:38, *Pseudomonas*]; [Augustin *et al.* (1990) *FEMS Microbiol. Lett.* 66:203, *Staphylococcus*], [Barany *et al.* (1980) *J. Bacteriol.* 144:698; Harlander (1987) "Transformation of *Streptococcus lactis* by electroporation, in: *Streptococcal Genetics* (ed. J. Ferretti and R. Curtiss III); Perry *et al.* (1981) *Infect. Immun.* 32:1295; Powell *et al.* (1988) *Appl. Environ. Microbiol.* 54:655; Somkuti *et al.* (1987) *Proc. 4th Evr. Cong. Biotechnology* 1:412, *Streptococcus*].

v. Yeast Expression

- Yeast expression systems are also known to one of ordinary skill in the art. A yeast promoter is any DNA sequence capable of binding yeast RNA polymerase and initiating the downstream (3') transcription of a coding sequence (*eg.* structural gene) into mRNA. A promoter will have a transcription initiation region which is usually placed proximal to the 5' end of the coding sequence. This transcription initiation region usually includes an RNA polymerase binding site (the "TATA Box") and a transcription initiation site. A yeast promoter may also have a second domain called an upstream activator sequence (UAS), which, if present, is usually distal to the structural gene. The UAS permits regulated (inducible) expression. Constitutive expression occurs in the absence of a UAS. Regulated expression may be either positive or negative, thereby either enhancing or reducing transcription.

- Yeast is a fermenting organism with an active metabolic pathway, therefore sequences encoding enzymes in the metabolic pathway provide particularly useful promoter sequences. Examples include alcohol dehydrogenase (ADH) (EP-A-0 284 044), enolase, glucokinase, glucose-6-phosphate isomerase, glyceraldehyde-3-phosphate-dehydrogenase (GAP or GAPDH), hexokinase, phosphofructokinase, 3-phosphoglycerate mutase, and pyruvate kinase (PyK) (EPO-A-0 329 203). The yeast *PHO5* gene, encoding acid phosphatase, also provides useful promoter sequences [Myanohara *et al.* (1983) *Proc. Natl. Acad. Sci. USA* 80:1].

- In addition, synthetic promoters which do not occur in nature also function as yeast promoters. For example, UAS sequences of one yeast promoter may be joined with the transcription activation region of another yeast promoter, creating a synthetic hybrid promoter. Examples of such hybrid promoters include the ADH regulatory sequence linked to the GAP transcription activation region (US Patent Nos. 4,876,197 and 4,880,734). Other examples of hybrid promoters include promoters which consist of the regulatory sequences of either the *ADH2*, *GAL4*, *GAL10*, OR *PHO5* genes,

combined with the transcriptional activation region of a glycolytic enzyme gene such as GAP or PyK (EP-A-0 164 556). Furthermore, a yeast promoter can include naturally occurring promoters of non-yeast origin that have the ability to bind yeast RNA polymerase and initiate transcription. Examples of such promoters include, *inter alia*, [Cohen *et al.* (1980) *Proc. Natl. Acad. Sci. USA* 77:1078; Henikoff *et al.* (1981) *Nature* 283:835; Hollenberg *et al.* (1981) *Curr. Topics Microbiol. Immunol.* 96:119; Hollenberg *et al.* (1979) "The Expression of Bacterial Antibiotic Resistance Genes in the Yeast *Saccharomyces cerevisiae*," in: *Plasmids of Medical, Environmental and Commercial Importance* (eds. K.N. Timmis and A. Puhler); Mercerau-Puigalon *et al.* (1980) *Gene* 11:163; Panthier *et al.* (1980) *Curr. Genet.* 2:109;].

- 10 A DNA molecule may be expressed intracellularly in yeast. A promoter sequence may be directly linked with the DNA molecule, in which case the first amino acid at the N-terminus of the recombinant protein will always be a methionine, which is encoded by the ATG start codon. If desired, methionine at the N-terminus may be cleaved from the protein by *in vitro* incubation with cyanogen bromide.
- 15 Fusion proteins provide an alternative for yeast expression systems, as well as in mammalian, baculovirus, and bacterial expression systems. Usually, a DNA sequence encoding the N-terminal portion of an endogenous yeast protein, or other stable protein, is fused to the 5' end of heterologous coding sequences. Upon expression, this construct will provide a fusion of the two amino acid sequences. For example, the yeast or human superoxide dismutase (SOD) gene, can be
- 20 linked at the 5' terminus of a foreign gene and expressed in yeast. The DNA sequence at the junction of the two amino acid sequences may or may not encode a cleavable site. See *eg.* EP-A-0 196 056. Another example is a ubiquitin fusion protein. Such a fusion protein is made with the ubiquitin region that preferably retains a site for a processing enzyme (*eg.* ubiquitin-specific processing protease) to cleave the ubiquitin from the foreign protein. Through this method,
- 25 therefore, native foreign protein can be isolated (*eg.* WO88/024066).

Alternatively, foreign proteins can also be secreted from the cell into the growth media by creating chimeric DNA molecules that encode a fusion protein comprised of a leader sequence fragment that provide for secretion in yeast of the foreign protein. Preferably, there are processing sites encoded between the leader fragment and the foreign gene that can be cleaved either *in vivo* or *in vitro*. The

leader sequence fragment usually encodes a signal peptide comprised of hydrophobic amino acids which direct the secretion of the protein from the cell.

DNA encoding suitable signal sequences can be derived from genes for secreted yeast proteins, such as the yeast invertase gene (EP-A-0 012 873; JPO. 62,096,086) and the A-factor gene (US
5 patent 4,588,684). Alternatively, leaders of non-yeast origin, such as an interferon leader, exist that also provide for secretion in yeast (EP-A-0 060 057).

A preferred class of secretion leaders are those that employ a fragment of the yeast alpha-factor gene, which contains both a "pre" signal sequence, and a "pro" region. The types of alpha-factor fragments that can be employed include the full-length pre-pro alpha factor leader (about 83 amino
10 acid residues) as well as truncated alpha-factor leaders (usually about 25 to about 50 amino acid residues) (US Patents 4,546,083 and 4,870,008; EP-A-0 324 274). Additional leaders employing an alpha-factor leader fragment that provides for secretion include hybrid alpha-factor leaders made with a presequence of a first yeast, but a pro-region from a second yeast alphafactor. (eg. see WO 89/02463.)

15 Usually, transcription termination sequences recognized by yeast are regulatory regions located 3' to the translation stop codon, and thus together with the promoter flank the coding sequence. These sequences direct the transcription of an mRNA which can be translated into the polypeptide encoded by the DNA. Examples of transcription terminator sequence and other yeast-recognized termination sequences, such as those coding for glycolytic enzymes.

20 Usually, the above described components, comprising a promoter, leader (if desired), coding sequence of interest, and transcription termination sequence, are put together into expression constructs. Expression constructs are often maintained in a replicon, such as an extrachromosomal element (eg. plasmids) capable of stable maintenance in a host, such as yeast or bacteria. The replicon may have two replication systems, thus allowing it to be maintained, for example, in yeast
25 for expression and in a prokaryotic host for cloning and amplification. Examples of such yeast-bacteria shuttle vectors include YEp24 [Botstein *et al.* (1979) *Gene* 8:17-24], pCl/1 [Brake *et al.* (1984) *Proc. Natl. Acad. Sci USA* 81:4642-4646], and YRp17 [Stinchcomb *et al.* (1982) *J. Mol. Biol.* 158:157]. In addition, a replicon may be either a high or low copy number plasmid. A high copy number plasmid will generally have a copy number ranging from about 5 to about 200, and

usually about 10 to about 150. A host containing a high copy number plasmid will preferably have at least about 10, and more preferably at least about 20. Either a high or low copy number vector may be selected, depending upon the effect of the vector and the foreign protein on the host. See *eg. Brake et al., supra*.

- 5 Alternatively, the expression constructs can be integrated into the yeast genome with an integrating vector. Integrating vectors usually contain at least one sequence homologous to a yeast chromosome that allows the vector to integrate, and preferably contain two homologous sequences flanking the expression construct. Integrations appear to result from recombinations between homologous DNA in the vector and the yeast chromosome [Orr-Weaver *et al.* (1983) *Methods in*
10 *Enzymol.* 101:228-245]. An integrating vector may be directed to a specific locus in yeast by selecting the appropriate homologous sequence for inclusion in the vector. See Orr-Weaver *et al., supra*. One or more expression construct may integrate, possibly affecting levels of recombinant protein produced [Rine *et al.* (1983) *Proc. Natl. Acad. Sci. USA* 80:6750]. The chromosomal sequences included in the vector can occur either as a single segment in the vector, which results
15 in the integration of the entire vector, or two segments homologous to adjacent segments in the chromosome and flanking the expression construct in the vector, which can result in the stable integration of only the expression construct.

- Usually, extrachromosomal and integrating expression constructs may contain selectable markers to allow for the selection of yeast strains that have been transformed. Selectable markers may
20 include biosynthetic genes that can be expressed in the yeast host, such as *ADE2*, *HIS4*, *LEU2*, *TRP1*, and *ALG7*, and the G418 resistance gene, which confer resistance in yeast cells to tunicamycin and G418, respectively. In addition, a suitable selectable marker may also provide yeast with the ability to grow in the presence of toxic compounds, such as metal. For example, the presence of *CUP1* allows yeast to grow in the presence of copper ions [Butt *et al.* (1987) *Microbiol.*
25 *Rev.* 51:351].

Alternatively, some of the above described components can be put together into transformation vectors. Transformation vectors are usually comprised of a selectable marker that is either maintained in a replicon or developed into an integrating vector, as described above.

Expression and transformation vectors, either extrachromosomal replicons or integrating vectors, have been developed for transformation into many yeasts. For example, expression vectors have been developed for, *inter alia*, the following yeasts: *Candida albicans* [Kurtz, *et al.* (1986) *Mol. Cell. Biol.* 6:142], *Candida maltosa* [Kunze, *et al.* (1985) *J. Basic Microbiol.* 25:141], *Hansenula polymorpha* [Gleeson, *et al.* (1986) *J. Gen. Microbiol.* 132:3459; Roggenkamp *et al.* (1986) *Mol. Gen. Genet.* 202:302], *Kluyveromyces fragilis* [Das, *et al.* (1984) *J. Bacteriol.* 158:1165], *Kluyveromyces lactis* [De Louvencourt *et al.* (1983) *J. Bacteriol.* 154:737; Van den Berg *et al.* (1990) *Bio/Technology* 8:135], *Pichia guilliermondii* [Kunze *et al.* (1985) *J. Basic Microbiol.* 25:141], *Pichia pastoris* [Cregg, *et al.* (1985) *Mol. Cell. Biol.* 5:3376; US Patent Nos. 4,837,148 and 4,929,555], *Saccharomyces cerevisiae* [Hinnen *et al.* (1978) *Proc. Natl. Acad. Sci. USA* 75:1929; Ito *et al.* (1983) *J. Bacteriol.* 153:163], *Schizosaccharomyces pombe* [Beach and Nurse (1981) *Nature* 300:706], and *Yarrowia lipolytica* [Davidow, *et al.* (1985) *Curr. Genet.* 10:380471 Gaillardin, *et al.* (1985) *Curr. Genet.* 10:49].

Methods of introducing exogenous DNA into yeast hosts are well-known in the art, and usually include either the transformation of spheroplasts or of intact yeast cells treated with alkali cations. Transformation procedures usually vary with the yeast species to be transformed. See *eg.* [Kurtz *et al.* (1986) *Mol. Cell. Biol.* 6:142; Kunze *et al.* (1985) *J. Basic Microbiol.* 25:141; *Candida*]; [Gleeson *et al.* (1986) *J. Gen. Microbiol.* 132:3459; Roggenkamp *et al.* (1986) *Mol. Gen. Genet.* 202:302; *Hansenula*]; [Das *et al.* (1984) *J. Bacteriol.* 158:1165; De Louvencourt *et al.* (1983) *J. Bacteriol.* 154:1165; Van den Berg *et al.* (1990) *Bio/Technology* 8:135; *Kluyveromyces*]; [Cregg *et al.* (1985) *Mol. Cell. Biol.* 5:3376; Kunze *et al.* (1985) *J. Basic Microbiol.* 25:141; US Patent Nos. 4,837,148 and 4,929,555; *Pichia*]; [Hinnen *et al.* (1978) *Proc. Natl. Acad. Sci. USA* 75:1929; Ito *et al.* (1983) *J. Bacteriol.* 153:163 *Saccharomyces*]; [Beach and Nurse (1981) *Nature* 300:706; *Schizosaccharomyces*]; [Davidow *et al.* (1985) *Curr. Genet.* 10:39; Gaillardin *et al.* (1985) *Curr. Genet.* 10:49; *Yarrowia*].

Antibodies

As used herein, the term "antibody" refers to a polypeptide or group of polypeptides composed of at least one antibody combining site. An "antibody combining site" is the three-dimensional binding space with an internal surface shape and charge distribution complementary to the features of an epitope of an antigen, which allows a binding of the antibody with the antigen. "Antibody"

includes, for example, vertebrate antibodies, hybrid antibodies, chimeric antibodies, humanised antibodies, altered antibodies, univalent antibodies, Fab proteins, and single domain antibodies.

Antibodies against the proteins of the invention are useful for affinity chromatography, immunoassays, and distinguishing/identifying Neisserial proteins.

- 5 Antibodies to the proteins of the invention, both polyclonal and monoclonal, may be prepared by conventional methods. In general, the protein is first used to immunize a suitable animal, preferably a mouse, rat, rabbit or goat. Rabbits and goats are preferred for the preparation of polyclonal sera due to the volume of serum obtainable, and the availability of labeled anti-rabbit and anti-goat antibodies. Immunization is generally performed by mixing or emulsifying the protein in saline,
- 10 preferably in an adjuvant such as Freund's complete adjuvant, and injecting the mixture or emulsion parenterally (generally subcutaneously or intramuscularly). A dose of 50-200 µg/injection is typically sufficient. Immunization is generally boosted 2-6 weeks later with one or more injections of the protein in saline, preferably using Freund's incomplete adjuvant. One may alternatively generate antibodies by *in vitro* immunization using methods known in the art, which
- 15 for the purposes of this invention is considered equivalent to *in vivo* immunization. Polyclonal antisera is obtained by bleeding the immunized animal into a glass or plastic container, incubating the blood at 25°C for one hour, followed by incubating at 4°C for 2-18 hours. The serum is recovered by centrifugation (*eg.* 1,000g for 10 minutes). About 20-50 ml per bleed may be obtained from rabbits.
- 20 Monoclonal antibodies are prepared using the standard method of Kohler & Milstein [*Nature* (1975) 256:495-96], or a modification thereof. Typically, a mouse or rat is immunized as described above. However, rather than bleeding the animal to extract serum, the spleen (and optionally several large lymph nodes) is removed and dissociated into single cells. If desired, the spleen cells may be screened (after removal of nonspecifically adherent cells) by applying a cell suspension to
- 25 a plate or well coated with the protein antigen. B-cells expressing membrane-bound immunoglobulin specific for the antigen bind to the plate, and are not rinsed away with the rest of the suspension. Resulting B-cells, or all dissociated spleen cells, are then induced to fuse with myeloma cells to form hybridomas, and are cultured in a selective medium (*eg.* hypoxanthine, aminopterin, thymidine medium, "HAT"). The resulting hybridomas are plated by limiting dilution,
- 30 and are assayed for the production of antibodies which bind specifically to the immunizing antigen

(and which do not bind to unrelated antigens). The selected MAb-secreting hybridomas are then cultured either *in vitro* (eg. in tissue culture bottles or hollow fiber reactors), or *in vivo* (as ascites in mice).

If desired, the antibodies (whether polyclonal or monoclonal) may be labeled using conventional techniques. Suitable labels include fluorophores, chromophores, radioactive atoms (particularly ^{32}P and ^{125}I), electron-dense reagents, enzymes, and ligands having specific binding partners. Enzymes are typically detected by their activity. For example, horseradish peroxidase is usually detected by its ability to convert 3,3',5,5'-tetramethylbenzidine (TMB) to a blue pigment, quantifiable with a spectrophotometer. "Specific binding partner" refers to a protein capable of binding a ligand molecule with high specificity, as for example in the case of an antigen and a monoclonal antibody specific therefor. Other specific binding partners include biotin and avidin or streptavidin, IgG and protein A, and the numerous receptor-ligand couples known in the art. It should be understood that the above description is not meant to categorize the various labels into distinct classes, as the same label may serve in several different modes. For example, ^{125}I may serve as a radioactive label or as an electron-dense reagent. HRP may serve as enzyme or as antigen for a MAb. Further, one may combine various labels for desired effect. For example, MAbs and avidin also require labels in the practice of this invention: thus, one might label a MAb with biotin, and detect its presence with avidin labeled with ^{125}I , or with an anti-biotin MAb labeled with HRP. Other permutations and possibilities will be readily apparent to those of ordinary skill in the art, and are considered as equivalents within the scope of the instant invention.

Pharmaceutical Compositions

Pharmaceutical compositions can comprise either polypeptides, antibodies, or nucleic acid of the invention. The pharmaceutical compositions will comprise a therapeutically effective amount of either polypeptides, antibodies, or polynucleotides of the claimed invention.

The term "therapeutically effective amount" as used herein refers to an amount of a therapeutic agent to treat, ameliorate, or prevent a desired disease or condition, or to exhibit a detectable therapeutic or preventative effect. The effect can be detected by, for example, chemical markers or antigen levels. Therapeutic effects also include reduction in physical symptoms, such as decreased body temperature. The precise effective amount for a subject will depend upon the subject's size and health, the nature and extent of the condition, and the therapeutics or combination of

therapeutics selected for administration. Thus, it is not useful to specify an exact effective amount in advance. However, the effective amount for a given situation can be determined by routine experimentation and is within the judgement of the clinician.

For purposes of the present invention, an effective dose will be from about 0.01 mg/ kg to 50 mg/kg or 0.05 mg/kg to about 10 mg/kg of the DNA constructs in the individual to which it is administered.

A pharmaceutical composition can also contain a pharmaceutically acceptable carrier. The term "pharmaceutically acceptable carrier" refers to a carrier for administration of a therapeutic agent, such as antibodies or a polypeptide, genes, and other therapeutic agents. The term refers to any pharmaceutical carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition, and which may be administered without undue toxicity. Suitable carriers may be large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, and inactive virus particles. Such carriers are well known to those of ordinary skill in the art.

Pharmaceutically acceptable salts can be used therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. A thorough discussion of pharmaceutically acceptable excipients is available in Remington's Pharmaceutical Sciences (Mack Pub. Co., N.J. 1991).

Pharmaceutically acceptable carriers in therapeutic compositions may contain liquids such as water, saline, glycerol and ethanol. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles. Typically, the therapeutic compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared. Liposomes are included within the definition of a pharmaceutically acceptable carrier.

Delivery Methods

Once formulated, the compositions of the invention can be administered directly to the subject. The subjects to be treated can be animals; in particular, human subjects can be treated.

Direct delivery of the compositions will generally be accomplished by injection, either subcutaneously, intraperitoneally, intravenously or intramuscularly or delivered to the interstitial space of a tissue. The compositions can also be administered into a lesion. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal or transcutaneous applications (eg. see WO98/20734), needles, and gene guns or hyposprays. Dosage treatment may be a single dose schedule or a multiple dose schedule.

Vaccines

Vaccines according to the invention may either be prophylactic (*ie.* to prevent infection) or therapeutic (*ie.* to treat disease after infection).

Such vaccines comprise immunising antigen(s), immunogen(s), polypeptide(s), protein(s) or nucleic acid, usually in combination with "pharmaceutically acceptable carriers," which include any carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, lipid aggregates (such as oil droplets or liposomes), and inactive virus particles. Such carriers are well known to those of ordinary skill in the art. Additionally, these carriers may function as immunostimulating agents ("adjuvants"). Furthermore, the antigen or immunogen may be conjugated to a bacterial toxoid, such as a toxoid from diphtheria, tetanus, cholera, *H. pylori*, etc. pathogens.

Preferred adjuvants to enhance effectiveness of the composition include, but are not limited to: (1) aluminum salts (alum), such as aluminum hydroxide, aluminum phosphate, aluminum sulfate, etc; (2) oil-in-water emulsion formulations (with or without other specific immunostimulating agents such as muramyl peptides (see below) or bacterial cell wall components), such as for example (a) MF59™ (WO 90/14837; Chapter 10 in *Vaccine design: the subunit and adjuvant approach*, eds. Powell & Newman, Plenum Press 1995), containing 5% Squalene, 0.5% Tween 80, and 0.5% Span 85 (optionally containing various amounts of MTP-PE (see below), although not required) formulated into submicron particles using a microfluidizer such as Model 110Y microfluidizer (Microfluidics, Newton, MA), (b) SAF, containing 10% Squalene, 0.4% Tween 80, 5% pluronic-blocked polymer L121, and thr-MDP (see below) either microfluidized into a submicron emulsion or vortexed to generate a larger particle size emulsion, and (c) Ribit™ adjuvant system (RAS), (Ribi Immunochem, Hamilton, MT) containing 2% Squalene, 0.2% Tween 80, and one or more bacterial

- cell wall components from the group consisting of monophosphorylipid A (MPL), trehalose dimycolate (TDM), and cell wall skeleton (CWS), preferably MPL + CWS (Detox™); (3) saponin adjuvants, such as Stimulon™ (Cambridge Bioscience, Worcester, MA) may be used or particles generated therefrom such as ISCOMs (immunostimulating complexes); (4) Complete Freund's Adjuvant (CFA) and Incomplete Freund's Adjuvant (IFA); (5) cytokines, such as interleukins (*eg.* IL-1, IL-2, IL-4, IL-5, IL-6, IL-7, IL-12, *etc.*), interferons (*eg.* gamma interferon), macrophage colony stimulating factor (M-CSF), tumor necrosis factor (TNF), *etc.*; and (6) other substances that act as immunostimulating agents to enhance the effectiveness of the composition. Alum and MF59™ are preferred.
- 10 As mentioned above, muramyl peptides include, but are not limited to, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-*sn*-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE), *etc.*

- 15 The immunogenic compositions (*eg.* the immunising antigen/immunogen/polypeptide/protein/nucleic acid, pharmaceutically acceptable carrier, and adjuvant) typically will contain diluents, such as water, saline, glycerol, ethanol, *etc.* Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles.

- Typically, the immunogenic compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared. The preparation also may be emulsified or encapsulated in liposomes for enhanced adjuvant effect, as discussed above under pharmaceutically acceptable carriers.
- 20

- Immunogenic compositions used as vaccines comprise an immunologically effective amount of the antigenic or immunogenic polypeptides, as well as any other of the above-mentioned components, as needed. By "immunologically effective amount", it is meant that the administration of that amount to an individual, either in a single dose or as part of a series, is effective for treatment or prevention. This amount varies depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated (*eg.* nonhuman primate, primate, *etc.*), the capacity of the individual's immune system to synthesize antibodies, the degree of protection desired, the formulation of the vaccine, the treating doctor's assessment of the medical situation,
- 25

and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials.

The immunogenic compositions are conventionally administered parenterally, *eg.* by injection, either subcutaneously, intramuscularly, or transdermally/transcutaneously (*eg.* WO98/20734).

- 5 Additional formulations suitable for other modes of administration include oral and pulmonary formulations, suppositories, and transdermal applications. Dosage treatment may be a single dose schedule or a multiple dose schedule. The vaccine may be administered in conjunction with other immunoregulatory agents.

- 10 As an alternative to protein-based vaccines, DNA vaccination may be employed [*eg.* Robinson & Torres (1997) *Seminars in Immunology* 9:271-283; Donnelly *et al.* (1997) *Annu Rev Immunol* 15:617-648; see later herein].

Gene Delivery Vehicles

- 15 Gene therapy vehicles for delivery of constructs including a coding sequence of a therapeutic of the invention, to be delivered to the mammal for expression in the mammal, can be administered either locally or systemically. These constructs can utilize viral or non-viral vector approaches in *in vivo* or *ex vivo* modality. Expression of such coding sequence can be induced using endogenous mammalian or heterologous promoters. Expression of the coding sequence *in vivo* can be either constitutive or regulated.

- 20 The invention includes gene delivery vehicles capable of expressing the contemplated nucleic acid sequences. The gene delivery vehicle is preferably a viral vector and, more preferably, a retroviral, adenoviral, adeno-associated viral (AAV), herpes viral, or alphavirus vector. The viral vector can also be an astrovirus, coronavirus, orthomyxovirus, papovavirus, paramyxovirus, parvovirus, picornavirus, poxvirus, or togavirus viral vector. See generally, Jolly (1994) *Cancer Gene Therapy* 1:51-64; Kimura (1994) *Human Gene Therapy* 5:845-852; Connelly (1995) *Human Gene Therapy* 25 6:185-193; and Kaplitt (1994) *Nature Genetics* 6:148-153.

Retroviral vectors are well known in the art and we contemplate that any retroviral gene therapy vector is employable in the invention, including B, C and D type retroviruses, xenotropic retroviruses (for example, NZB-X1, NZB-X2 and NZB9-1 (see O'Neill (1985) *J. Virol.* 53:160) polytropic retroviruses

eg. MCF and MCF-MLV (see Kelly (1983) *J. Virol.* 45:291), spumaviruses and lentiviruses. See RNA Tumor Viruses, Second Edition, Cold Spring Harbor Laboratory, 1985.

Portions of the retroviral gene therapy vector may be derived from different retroviruses. For example, retrovector LTRs may be derived from a Murine Sarcoma Virus, a tRNA binding site
5 from a Rous Sarcoma Virus, a packaging signal from a Murine Leukemia Virus, and an origin of second strand synthesis from an Avian Leukosis Virus.

These recombinant retroviral vectors may be used to generate transduction competent retroviral vector particles by introducing them into appropriate packaging cell lines (see US patent 5,591,624). Retrovirus vectors can be constructed for site-specific integration into host cell DNA
10 by incorporation of a chimeric integrase enzyme into the retroviral particle (see WO96/37626). It is preferable that the recombinant viral vector is a replication defective recombinant virus.

Packaging cell lines suitable for use with the above-described retrovirus vectors are well known in the art, are readily prepared (see WO95/30763 and WO92/05266), and can be used to create producer cell lines (also termed vector cell lines or "VCLs") for the production of recombinant
15 vector particles. Preferably, the packaging cell lines are made from human parent cells (eg. HT1080 cells) or mink parent cell lines, which eliminates inactivation in human serum.

Preferred retroviruses for the construction of retroviral gene therapy vectors include Avian Leukosis Virus, Bovine Leukemia, Virus, Murine Leukemia Virus, Mink-Cell Focus-Inducing Virus, Murine Sarcoma Virus, Reticuloendotheliosis Virus and Rous Sarcoma Virus. Particularly
20 preferred Murine Leukemia Viruses include 4070A and 1504A (Hartley and Rowe (1976) *J Virol* 19:19-25), Abelson (ATCC No. VR-999), Friend (ATCC No. VR-245), Graffi, Gross (ATCC No. VR-590), Kirsten, Harvey Sarcoma Virus and Rauscher (ATCC No. VR-998) and Moloney Murine Leukemia Virus (ATCC No. VR-190). Such retroviruses may be obtained from depositories or collections such as the American Type Culture Collection ("ATCC") in Rockville, Maryland or
25 isolated from known sources using commonly available techniques.

Exemplary known retroviral gene therapy vectors employable in this invention include those described in patent applications GB2200651, EP0415731, EP0345242, EP0334301, WO89/02468; WO89/05349, WO89/09271, WO90/02806, WO90/07936, WO94/03622, WO93/25698,

WO93/25234, WO93/11230, WO93/10218, WO91/02805, WO91/02825, WO95/07994, US 5,219,740, US 4,405,712, US 4,861,719, US 4,980,289, US 4,777,127, US 5,591,624. See also Vile (1993) *Cancer Res* 53:3860-3864; Vile (1993) *Cancer Res* 53:962-967; Ram (1993) *Cancer Res* 53 (1993) 83-88; Takamiya (1992) *J Neurosci Res* 33:493-503; Baba (1993) *J Neurosurg* 79:729-735; Mann (1983) *Cell* 33:153; Cane (1984) *Proc Natl Acad Sci* 81:6349; and Miller (1990) *Human Gene Therapy* 1.

Human adenoviral gene therapy vectors are also known in the art and employable in this invention. See, for example, Berkner (1988) *Biotechniques* 6:616 and Rosenfeld (1991) *Science* 252:431, and WO93/07283, WO93/06223, and WO93/07282. Exemplary known adenoviral gene therapy vectors employable in this invention include those described in the above referenced documents and in WO94/12649, WO93/03769, WO93/19191, WO94/28938, WO95/11984, WO95/00655, WO95/27071, WO95/29993, WO95/34671, WO96/05320, WO94/08026, WO94/11506, WO93/06223, WO94/24299, WO95/14102, WO95/24297, WO95/02697, WO94/28152, WO94/24299, WO95/09241, WO95/25807, WO95/05835, WO94/18922 and WO95/09654.

Alternatively, administration of DNA linked to killed adenovirus as described in Curiel (1992) *Hum. Gene Ther.* 3:147-154 may be employed. The gene delivery vehicles of the invention also include adenovirus associated virus (AAV) vectors. Leading and preferred examples of such vectors for use in this invention are the AAV-2 based vectors disclosed in Srivastava, WO93/09239. Most preferred AAV vectors comprise the two AAV inverted terminal repeats in which the native D-sequences are modified by substitution of nucleotides, such that at least 5 native nucleotides and up to 18 native nucleotides, preferably at least 10 native nucleotides up to 18 native nucleotides, most preferably 10 native nucleotides are retained and the remaining nucleotides of the D-sequence are deleted or replaced with non-native nucleotides. The native D-sequences of the AAV inverted terminal repeats are sequences of 20 consecutive nucleotides in each AAV inverted terminal repeat (*ie.* there is one sequence at each end) which are not involved in HP formation. The non-native replacement nucleotide may be any nucleotide other than the nucleotide found in the native D-sequence in the same position. Other employable exemplary AAV vectors are pWP-19, pWN-1, both of which are disclosed in Nahreini (1993) *Gene* 124:257-262. Another example of such an AAV vector is psub201 (see Samulski (1987) *J. Virol.* 61:3096). Another exemplary AAV vector is the Double-D ITR vector. Construction of the Double-D ITR vector is disclosed in US Patent 5,478,745. Still other vectors are those disclosed in Carter US Patent 4,797,368 and

Muzyczka US Patent 5,139,941, Chartejee US Patent 5,474,935, and Kotin WO94/288157. Yet a further example of an AAV vector employable in this invention is SSV9AFABTKneo, which contains the AFP enhancer and albumin promoter and directs expression predominantly in the liver. Its structure and construction are disclosed in Su (1996) *Human Gene Therapy* 7:463-470.

5 Additional AAV gene therapy vectors are described in US 5,354,678, US 5,173,414, US 5,139,941, and US 5,252,479.

The gene therapy vectors of the invention also include herpes vectors. Leading and preferred examples are herpes simplex virus vectors containing a sequence encoding a thymidine kinase polypeptide such as those disclosed in US 5,288,641 and EP0176170 (Roizman). Additional

10 exemplary herpes simplex virus vectors include HFEM/ICP6-LacZ disclosed in WO95/04139 (Wistar Institute), pHSVlac described in Geller (1988) *Science* 241:1667-1669 and in WO90/09441 and WO92/07945, HSV Us3::pgC-lacZ described in Fink (1992) *Human Gene Therapy* 3:11-19 and HSV 7134, 2 RH105 and GAL4 described in EP 0453242 (Breakefield), and those deposited with the ATCC as accession numbers ATCC VR-977 and ATCC VR-260.

15 Also contemplated are alpha virus gene therapy vectors that can be employed in this invention. Preferred alpha virus vectors are Sindbis viruses vectors. Togaviruses, Semliki Forest virus (ATCC VR-67; ATCC VR-1247), Middleberg virus (ATCC VR-370), Ross River virus (ATCC VR-373; ATCC VR-1246), Venezuelan equine encephalitis virus (ATCC VR923; ATCC VR-1250; ATCC VR-1249; ATCC VR-532), and those described in US patents 5,091,309, 5,217,879, and

20 WO92/10578. More particularly, those alpha virus vectors described in US Serial No. 08/405,627, filed March 15, 1995, WO94/21792, WO92/10578, WO95/07994, US 5,091,309 and US 5,217,879 are employable. Such alpha viruses may be obtained from depositories or collections such as the ATCC in Rockville, Maryland or isolated from known sources using commonly available techniques. Preferably, alphavirus vectors with reduced cytotoxicity are used (see USSN

25 08/679640).

DNA vector systems such as eukaryotic layered expression systems are also useful for expressing the nucleic acids of the invention. See WO95/07994 for a detailed description of eukaryotic layered expression systems. Preferably, the eukaryotic layered expression systems of the invention are derived from alphavirus vectors and most preferably from Sindbis viral vectors.

Other viral vectors suitable for use in the present invention include those derived from poliovirus, for example ATCC VR-58 and those described in Evans, Nature 339 (1989) 385 and Sabin (1973) *J. Biol. Standardization* 1:115; rhinovirus, for example ATCC VR-1110 and those described in Arnold (1990) *J Cell Biochem* L401; pox viruses such as canary pox virus or vaccinia virus, for example ATCC VR-111 and ATCC VR-2010 and those described in Fisher-Hoch (1989) *Proc Natl Acad Sci* 86:317; Flexner (1989) *Ann NY Acad Sci* 569:86, Flexner (1990) *Vaccine* 8:17; in US 4,603,112 and US 4,769,330 and WO89/01973; SV40 virus, for example ATCC VR-305 and those described in Mulligan (1979) *Nature* 277:108 and Madzak (1992) *J Gen Virol* 73:1533; influenza virus, for example ATCC VR-797 and recombinant influenza viruses made employing reverse genetics techniques as described in US 5,166,057 and in Enami (1990) *Proc Natl Acad Sci* 87:3802-3805; Enami & Palese (1991) *J Virol* 65:2711-2713 and Luytjes (1989) *Cell* 59:110, (see also McMichael (1983) *NEJ Med* 309:13, and Yap (1978) *Nature* 273:238 and *Nature* (1979) 277:108); human immunodeficiency virus as described in EP-0386882 and in Buchschacher (1992) *J. Virol.* 66:2731; measles virus, for example ATCC VR-67 and VR-1247 and those described in EP-0440219; Aura virus, for example ATCC VR-368; Bebaru virus, for example ATCC VR-600 and ATCC VR-1240; Cabassou virus, for example ATCC VR-922; Chikungunya virus, for example ATCC VR-64 and ATCC VR-1241; Fort Morgan Virus, for example ATCC VR-924; Getah virus, for example ATCC VR-369 and ATCC VR-1243; Kyzylagach virus, for example ATCC VR-927; Mayaro virus, for example ATCC VR-66; Mucambo virus, for example ATCC VR-580 and ATCC VR-1244; Ndumu virus, for example ATCC VR-371; Pixuna virus, for example ATCC VR-372 and ATCC VR-1245; Tonate virus, for example ATCC VR-925; Trinita virus, for example ATCC VR-469; Una virus, for example ATCC VR-374; Whataroa virus, for example ATCC VR-926; Y-62-33 virus, for example ATCC VR-375; O'Nyong virus, Eastern encephalitis virus, for example ATCC VR-65 and ATCC VR-1242; Western encephalitis virus, for example ATCC VR-70, ATCC VR-1251, ATCC VR-622 and ATCC VR-1252; and coronavirus, for example ATCC VR-740 and those described in Hamre (1966) *Proc Soc Exp Biol Med* 121:190.

Delivery of the compositions of this invention into cells is not limited to the above mentioned viral vectors. Other delivery methods and media may be employed such as, for example, nucleic acid expression vectors, polycationic condensed DNA linked or unlinked to killed adenovirus alone, for example see US Serial No. 08/366,787, filed December 30, 1994 and Curiel (1992) *Hum Gene Ther* 3:147-154 ligand linked DNA, for example see Wu (1989) *J Biol Chem* 264:16985-16987, eucaryotic cell delivery vehicles cells, for example see US Serial No.08/240,030, filed May 9,

1994, and US Serial No. 08/404,796, deposition of photopolymerized hydrogel materials, hand-held gene transfer particle gun, as described in US Patent 5,149,655, ionizing radiation as described in US5,206,152 and in WO92/11033, nucleic charge neutralization or fusion with cell membranes. Additional approaches are described in Philip (1994) *Mol Cell Biol* 14:2411-2418 and
5 in Woffendin (1994) *Proc Natl Acad Sci* 91:1581-1585.

Particle mediated gene transfer may be employed, for example see US Serial No. 60/023,867. Briefly, the sequence can be inserted into conventional vectors that contain conventional control sequences for high level expression, and then incubated with synthetic gene transfer molecules such as polymeric DNA-binding cations like polylysine, protamine, and albumin, linked to cell targeting
10 ligands such as asialoorosomuroid, as described in Wu & Wu (1987) *J. Biol. Chem.* 262:4429-4432, insulin as described in Hucked (1990) *Biochem Pharmacol* 40:253-263, galactose as described in Plank (1992) *Bioconjugate Chem* 3:533-539, lactose or transferrin.

Naked DNA may also be employed. Exemplary naked DNA introduction methods are described in WO 90/11092 and US 5,580,859. Uptake efficiency may be improved using biodegradable latex
15 beads. DNA coated latex beads are efficiently transported into cells after endocytosis initiation by the beads. The method may be improved further by treatment of the beads to increase hydrophobicity and thereby facilitate disruption of the endosome and release of the DNA into the cytoplasm.

Liposomes that can act as gene delivery vehicles are described in US 5,422,120, WO95/13796, WO94/23697, WO91/14445 and EP-524,968. As described in USSN. 60/023,867, on non-viral
20 delivery, the nucleic acid sequences encoding a polypeptide can be inserted into conventional vectors that contain conventional control sequences for high level expression, and then be incubated with synthetic gene transfer molecules such as polymeric DNA-binding cations like polylysine, protamine, and albumin, linked to cell targeting ligands such as asialoorosomuroid, insulin, galactose, lactose, or transferrin. Other delivery systems include the use of liposomes to encapsulate
25 DNA comprising the gene under the control of a variety of tissue-specific or ubiquitously-active promoters. Further non-viral delivery suitable for use includes mechanical delivery systems such as the approach described in Woffendin *et al* (1994) *Proc. Natl. Acad. Sci. USA* 91(24):11581-11585. Moreover, the coding sequence and the product of expression of such can be delivered through deposition of photopolymerized hydrogel materials. Other conventional methods
30 for gene delivery that can be used for delivery of the coding sequence include, for example, use of

hand-held gene transfer particle gun, as described in US 5,149,655; use of ionizing radiation for activating transferred gene, as described in US 5,206,152 and WO92/11033

Exemplary liposome and polycationic gene delivery vehicles are those described in US 5,422,120 and 4,762,915; in WO 95/13796; WO94/23697; and WO91/14445; in EP-0524968; and in Stryer, 5 Biochemistry, pages 236-240 (1975) W.H. Freeman, San Francisco; Szoka (1980) *Biochem Biophys Acta* 600:1; Bayer (1979) *Biochem Biophys Acta* 550:464; Rivnay (1987) *Meth Enzymol* 149:119; Wang (1987) *Proc Natl Acad Sci* 84:7851; Plant (1989) *Anal Biochem* 176:420.

A polynucleotide composition can comprises therapeutically effective amount of a gene therapy vehicle, as the term is defined above. For purposes of the present invention, an effective dose will 10 be from about 0.01 mg/ kg to 50 mg/kg or 0.05 mg/kg to about 10 mg/kg of the DNA constructs in the individual to which it is administered.

Delivery Methods

Once formulated, the polynucleotide compositions of the invention can be administered (1) directly to the subject; (2) delivered *ex vivo*, to cells derived from the subject; or (3) *in vitro* for expression 15 of recombinant proteins. The subjects to be treated can be mammals or birds. Also, human subjects can be treated.

Direct delivery of the compositions will generally be accomplished by injection, either subcutaneously, intraperitoneally, intravenously or intramuscularly or delivered to the interstitial space of a tissue. The compositions can also be administered into a lesion. Other modes of 20 administration include oral and pulmonary administration, suppositories, and transdermal or transcutaneous applications (eg. see WO98/20734), needles, and gene guns or hyposprays. Dosage treatment may be a single dose schedule or a multiple dose schedule.

Methods for the *ex vivo* delivery and reimplantation of transformed cells into a subject are known in the art and described in eg. WO93/14778. Examples of cells useful in *ex vivo* applications 25 include, for example, stem cells, particularly hematopoietic, lymph cells, macrophages, dendritic cells, or tumor cells.

Generally, delivery of nucleic acids for both *ex vivo* and *in vitro* applications can be accomplished by the following procedures, for example, dextran-mediated transfection, calcium phosphate precipitation, polybrene mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei, all well known in the art.

Polynucleotide and polypeptide pharmaceutical compositions

In addition to the pharmaceutically acceptable carriers and salts described above, the following additional agents can be used with polynucleotide and/or polypeptide compositions.

A. Polypeptides

- One example are polypeptides which include, without limitation: asialoglycoproteins; antibodies; antibody fragments; ferritin; interleukins; interferons, granulocyte, macrophage colony stimulating factor (GM-CSF), granulocyte colony stimulating factor (G-CSF), macrophage colony stimulating factor (M-CSF), stem cell factor and erythropoietin. Viral antigens, such as envelope proteins, can also be used. Also, proteins from other invasive organisms, such as the 17 amino acid peptide from the circumsporozoite protein of plasmodium falciparum known as RII.

B. Hormones, Vitamins, etc.

Other groups that can be included are, for example: hormones, steroids, androgens, estrogens, thyroid hormone, or vitamins, folic acid.

C. Polyalkylenes, Polysaccharides, etc.

Also, polyalkylene glycol can be included with the desired polynucleotides/polypeptides. In a preferred embodiment, the polyalkylene glycol is polyethylene glycol. In addition, mono-, di-, or polysaccharides can be included. In a preferred embodiment of this aspect, the polysaccharide is dextran or DEAE-dextran. Also, chitosan and poly(lactide-co-glycolide)

D. Lipids, and Liposomes

The desired polynucleotide/polypeptide can also be encapsulated in lipids or packaged in liposomes prior to delivery to the subject or to cells derived therefrom.

Lipid encapsulation is generally accomplished using liposomes which are able to stably bind or entrap and retain nucleic acid. The ratio of condensed polynucleotide to lipid preparation can vary but will generally be around 1:1 (mg DNA:micromoles lipid), or more of lipid. For a review of the use of liposomes as carriers for delivery of nucleic acids, see, Hug and Sleight (1991) *Biochim.*

5 *Biophys. Acta.* 1097:1-17; Straubinger (1983) *Meth. Enzymol.* 101:512-527.

Liposomal preparations for use in the present invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. Cationic liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner (1987) *Proc. Natl. Acad. Sci. USA* 84:7413-7416); mRNA (Malone (1989) *Proc. Natl. Acad. Sci. USA* 86:6077-6081); and purified
10 transcription factors (Debs (1990) *J. Biol. Chem.* 265:10189-10192), in functional form.

Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, NY. (See, also, Felgner *supra*). Other commercially available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer). Other cationic liposomes can be
15 prepared from readily available materials using techniques well known in the art. See, *eg.* Szoka (1978) *Proc. Natl. Acad. Sci. USA* 75:4194-4198; WO90/11092 for a description of the synthesis of DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes.

Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, AL), or can be easily prepared using readily available materials. Such materials include
20 phosphatidyl choline, cholesterol, phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

The liposomes can comprise multilammellar vesicles (MLVs), small unilamellar vesicles (SUVs),
25 or large unilamellar vesicles (LUVs). The various liposome-nucleic acid complexes are prepared using methods known in the art. See *eg.* Straubinger (1983) *Meth. Immunol.* 101:512-527; Szoka (1978) *Proc. Natl. Acad. Sci. USA* 75:4194-4198; Papahadjopoulos (1975) *Biochim. Biophys. Acta* 394:483; Wilson (1979) *Cell* 17:77; Deamer & Bangham (1976) *Biochim. Biophys. Acta* 443:629; Ostro (1977) *Biochem. Biophys. Res. Commun.* 76:836; Fraley (1979) *Proc. Natl. Acad. Sci. USA*

76:3348); Enoch & Strittmatter (1979) *Proc. Natl. Acad. Sci. USA* 76:145; Fraley (1980) *J. Biol. Chem.* (1980) 255:10431; Szoka & Papahadjopoulos (1978) *Proc. Natl. Acad. Sci. USA* 75:145; and Schaefer-Ridder (1982) *Science* 215:166.

E.Lipoproteins

- 5 In addition, lipoproteins can be included with the polynucleotide/polypeptide to be delivered. Examples of lipoproteins to be utilized include: chylomicrons, HDL, IDL, LDL, and VLDL. Mutants, fragments, or fusions of these proteins can also be used. Also, modifications of naturally occurring lipoproteins can be used, such as acetylated LDL. These lipoproteins can target the delivery of polynucleotides to cells expressing lipoprotein receptors. Preferably, if lipoproteins are including with
10 the polynucleotide to be delivered, no other targeting ligand is included in the composition.

Naturally occurring lipoproteins comprise a lipid and a protein portion. The protein portion are known as apoproteins. At the present, apoproteins A, B, C, D, and E have been isolated and identified. At least two of these contain several proteins, designated by Roman numerals, AI, AII, AIV; CI, CII, CIII.

- 15 A lipoprotein can comprise more than one apoprotein. For example, naturally occurring chylomicrons comprises of A, B, C, and E, over time these lipoproteins lose A and acquire C and E apoproteins. VLDL comprises A, B, C, and E apoproteins, LDL comprises apoprotein B; and HDL comprises apoproteins A, C, and E.

- The amino acid of these apoproteins are known and are described in, for example, Breslow (1985)
20 *Annu Rev. Biochem* 54:699; Law (1986) *Adv. Exp Med. Biol.* 151:162; Chen (1986) *J Biol Chem* 261:12918; Kane (1980) *Proc Natl Acad Sci USA* 77:2465; and Utermann (1984) *Hum Genet* 65:232.

- Lipoproteins contain a variety of lipids including, triglycerides, cholesterol (free and esters), and phospholipids. The composition of the lipids varies in naturally occurring lipoproteins. For example, chylomicrons comprise mainly triglycerides. A more detailed description of the lipid
25 content of naturally occurring lipoproteins can be found, for example, in *Meth. Enzymol.* 128 (1986). The composition of the lipids are chosen to aid in conformation of the apoprotein for receptor binding activity. The composition of lipids can also be chosen to facilitate hydrophobic interaction and association with the polynucleotide binding molecule.

Naturally occurring lipoproteins can be isolated from serum by ultracentrifugation, for instance. Such methods are described in *Meth. Enzymol. (supra)*; Pitas (1980) *J. Biochem.* 255:5454-5460 and Mahey (1979) *J Clin. Invest* 64:743-750. Lipoproteins can also be produced by *in vitro* or recombinant methods by expression of the apoprotein genes in a desired host cell. See, for example, 5 Atkinson (1986) *Annu Rev Biophys Chem* 15:403 and Radding (1958) *Biochim Biophys Acta* 30: 443. Lipoproteins can also be purchased from commercial suppliers, such as Biomedical Technologies, Inc., Stoughton, Massachusetts, USA. Further description of lipoproteins can be found in Zuckermann *et al.* PCT/US97/14465.

F. Polycationic Agents

- 10 Polycationic agents can be included, with or without lipoprotein, in a composition with the desired polynucleotide/polypeptide to be delivered.

Polycationic agents, typically, exhibit a net positive charge at physiological relevant pH and are capable of neutralizing the electrical charge of nucleic acids to facilitate delivery to a desired location. These agents have both *in vitro*, *ex vivo*, and *in vivo* applications. Polycationic agents can 15 be used to deliver nucleic acids to a living subject either intramuscularly, subcutaneously, etc.

The following are examples of useful polypeptides as polycationic agents: polylysine, polyarginine, polyornithine, and protamine. Other examples include histones, protamines, human serum albumin, DNA binding proteins, non-histone chromosomal proteins, coat proteins from DNA viruses, such as (X174, transcriptional factors also contain domains that bind DNA and therefore may be useful 20 as nucleic acid condensing agents. Briefly, transcriptional factors such as C/CEBP, c-jun, c-fos, AP-1, AP-2, AP-3, CPF, Prot-1, Sp-1, Oct-1, Oct-2, CREP, and TFIID contain basic domains that bind DNA sequences.

Organic polycationic agents include: spermine, spermidine, and putrescine.

The dimensions and of the physical properties of a polycationic agent can be extrapolated from the 25 list above, to construct other polypeptide polycationic agents or to produce synthetic polycationic agents.

Synthetic polycationic agents which are useful include, for example, DEAE-dextran, polybrene. Lipofectin™, and lipofectAMINE™ are monomers that form polycationic complexes when combined with polynucleotides/polypeptides.

Immunodiagnostic Assays

5 Neisserial antigens of the invention can be used in immunoassays to detect antibody levels (or, conversely, anti-Neisserial antibodies can be used to detect antigen levels). Immunoassays based on well defined, recombinant antigens can be developed to replace invasive diagnostics methods. Antibodies to Neisserial proteins within biological samples, including for example, blood or serum samples, can be detected. Design of the immunoassays is subject to a great deal of variation, and
10 a variety of these are known in the art. Protocols for the immunoassay may be based, for example, upon competition, or direct reaction, or sandwich type assays. Protocols may also, for example, use solid supports, or may be by immunoprecipitation. Most assays involve the use of labeled antibody or polypeptide; the labels may be, for example, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the probe are also known; examples of which
15 are assays which utilize biotin and avidin, and enzyme-labeled and mediated immunoassays, such as ELISA assays.

Kits suitable for immunodiagnosis and containing the appropriate labeled reagents are constructed by packaging the appropriate materials, including the compositions of the invention, in suitable containers, along with the remaining reagents and materials (for example, suitable buffers, salt
20 solutions, *etc.*) required for the conduct of the assay, as well as suitable set of assay instructions.

Nucleic Acid Hybridisation

“Hybridization” refers to the association of two nucleic acid sequences to one another by hydrogen bonding. Typically, one sequence will be fixed to a solid support and the other will be free in solution. Then, the two sequences will be placed in contact with one another under conditions that favor
25 hydrogen bonding. Factors that affect this bonding include: the type and volume of solvent; reaction temperature; time of hybridization; agitation; agents to block the non-specific attachment of the liquid phase sequence to the solid support (Denhardt's reagent or BLOTTO); concentration of the sequences; use of compounds to increase the rate of association of sequences (dextran sulfate or polyethylene glycol); and the stringency of the washing conditions following hybridization. See Sambrook *et al.*
30 [*supra*] Volume 2, chapter 9, pages 9.47 to 9.57.

“Stringency” refers to conditions in a hybridization reaction that favor association of very similar sequences over sequences that differ. For example, the combination of temperature and salt concentration should be chosen that is approximately 120 to 200°C below the calculated T_m of the hybrid under study. The temperature and salt conditions can often be determined empirically in preliminary experiments in which samples of genomic DNA immobilized on filters are hybridized to the sequence of interest and then washed under conditions of different stringencies. See Sambrook *et al.* at page 9.50.

Variables to consider when performing, for example, a Southern blot are (1) the complexity of the DNA being blotted and (2) the homology between the probe and the sequences being detected. The total amount of the fragment(s) to be studied can vary a magnitude of 10, from 0.1 to 1 µg for a plasmid or phage digest to 10^{-9} to 10^{-8} g for a single copy gene in a highly complex eukaryotic genome. For lower complexity polynucleotides, substantially shorter blotting, hybridization, and exposure times, a smaller amount of starting polynucleotides, and lower specific activity of probes can be used. For example, a single-copy yeast gene can be detected with an exposure time of only 1 hour starting with 1 µg of yeast DNA, blotting for two hours, and hybridizing for 4-8 hours with a probe of 10^8 cpm/µg. For a single-copy mammalian gene a conservative approach would start with 10 µg of DNA, blot overnight, and hybridize overnight in the presence of 10% dextran sulfate using a probe of greater than 10^8 cpm/µg, resulting in an exposure time of ~24 hours.

Several factors can affect the melting temperature (T_m) of a DNA-DNA hybrid between the probe and the fragment of interest, and consequently, the appropriate conditions for hybridization and washing. In many cases the probe is not 100% homologous to the fragment. Other commonly encountered variables include the length and total G+C content of the hybridizing sequences and the ionic strength and formamide content of the hybridization buffer. The effects of all of these factors can be approximated by a single equation:

$$T_m = 81 + 16.6(\log_{10} C_i) + 0.4[\%(G + C)] - 0.6(\%\text{formamide}) - 600/n - 1.5(\%\text{mismatch}).$$

where C_i is the salt concentration (monovalent ions) and n is the length of the hybrid in base pairs (slightly modified from Meinkoth & Wahl (1984) *Anal. Biochem.* 138: 267-284).

In designing a hybridization experiment, some factors affecting nucleic acid hybridization can be conveniently altered. The temperature of the hybridization and washes and the salt concentration during the washes are the simplest to adjust. As the temperature of the hybridization increases (*ie.* stringency), it becomes less likely for hybridization to occur between strands that are nonhomologous, and as a result, background decreases. If the radiolabeled probe is not completely homologous with the immobilized fragment (as is frequently the case in gene family and interspecies hybridization experiments), the hybridization temperature must be reduced, and background will increase. The temperature of the washes affects the intensity of the hybridizing band and the degree of background in a similar manner. The stringency of the washes is also increased with decreasing salt concentrations.

In general, convenient hybridization temperatures in the presence of 50% formamide are 42°C for a probe with is 95% to 100% homologous to the target fragment, 37°C for 90% to 95% homology, and 32°C for 85% to 90% homology. For lower homologies, formamide content should be lowered and temperature adjusted accordingly, using the equation above. If the homology between the probe and the target fragment are not known, the simplest approach is to start with both hybridization and wash conditions which are nonstringent. If non-specific bands or high background are observed after autoradiography, the filter can be washed at high stringency and reexposed. If the time required for exposure makes this approach impractical, several hybridization and/or washing stringencies should be tested in parallel.

20 Nucleic Acid Probe Assays

Methods such as PCR, branched DNA probe assays, or blotting techniques utilizing nucleic acid probes according to the invention can determine the presence of cDNA or mRNA. A probe is said to "hybridize" with a sequence of the invention if it can form a duplex or double stranded complex, which is stable enough to be detected.

25 The nucleic acid probes will hybridize to the Neisserial nucleotide sequences of the invention (including both sense and antisense strands). Though many different nucleotide sequences will encode the amino acid sequence, the native Neisserial sequence is preferred because it is the actual sequence present in cells. mRNA represents a coding sequence and so a probe should be complementary to the coding sequence; single-stranded cDNA is complementary to mRNA, and
30 so a cDNA probe should be complementary to the non-coding sequence.

The probe sequence need not be identical to the Neisserial sequence (or its complement) — some variation in the sequence and length can lead to increased assay sensitivity if the nucleic acid probe can form a duplex with target nucleotides, which can be detected. Also, the nucleic acid probe can include additional nucleotides to stabilize the formed duplex. Additional Neisserial sequence may
5 also be helpful as a label to detect the formed duplex. For example, a non-complementary nucleotide sequence may be attached to the 5' end of the probe, with the remainder of the probe sequence being complementary to a Neisserial sequence. Alternatively, non-complementary bases or longer sequences can be interspersed into the probe, provided that the probe sequence has sufficient complementarity with the a Neisserial sequence in order to hybridize therewith and
10 thereby form a duplex which can be detected.

The exact length and sequence of the probe will depend on the hybridization conditions, such as temperature, salt condition and the like. For example, for diagnostic applications, depending on the complexity of the analyte sequence, the nucleic acid probe typically contains at least 10-20 nucleotides, preferably 15-25, and more preferably at least 30 nucleotides, although it may be
15 shorter than this. Short primers generally require cooler temperatures to form sufficiently stable hybrid complexes with the template.

Probes may be produced by synthetic procedures, such as the triester method of Matteucci *et al.* [*J. Am. Chem. Soc.* (1981) 103:3185], or according to Urdea *et al.* [*Proc. Natl. Acad. Sci. USA* (1983) 80: 7461], or using commercially available automated oligonucleotide synthesizers.

20 The chemical nature of the probe can be selected according to preference. For certain applications, DNA or RNA are appropriate. For other applications, modifications may be incorporated *eg.* backbone modifications, such as phosphorothioates or methylphosphonates, can be used to increase *in vivo* half-life, alter RNA affinity, increase nuclease resistance *etc.* [*eg.* see Agrawal & Iyer (1995) *Curr Opin Biotechnol* 6:12-19; Agrawal (1996) *TIBTECH* 14:376-387]; analogues such as
25 peptide nucleic acids may also be used [*eg.* see Corey (1997) *TIBTECH* 15:224-229; Buchardt *et al.* (1993) *TIBTECH* 11:384-386].

Alternatively, the polymerase chain reaction (PCR) is another well-known means for detecting small amounts of target nucleic acids. The assay is described in: Mullis *et al.* [*Meth. Enzymol.* (1987) 155: 335-350]; US patents 4,683,195 and 4,683,202. Two "primer" nucleotides hybridize

with the target nucleic acids and are used to prime the reaction. The primers can comprise sequence that does not hybridize to the sequence of the amplification target (or its complement) to aid with duplex stability or, for example, to incorporate a convenient restriction site. Typically, such sequence will flank the desired Neisserial sequence.

- 5 A thermostable polymerase creates copies of target nucleic acids from the primers using the original target nucleic acids as a template. After a threshold amount of target nucleic acids are generated by the polymerase, they can be detected by more traditional methods, such as Southern blots. When using the Southern blot method, the labelled probe will hybridize to the Neisserial sequence (or its complement).
- 10 Also, mRNA or cDNA can be detected by traditional blotting techniques described in Sambrook *et al [supra]*. mRNA, or cDNA generated from mRNA using a polymerase enzyme, can be purified and separated using gel electrophoresis. The nucleic acids on the gel are then blotted onto a solid support, such as nitrocellulose. The solid support is exposed to a labelled probe and then washed to remove any unhybridized probe. Next, the duplexes containing the labeled probe are detected.
- 15 Typically, the probe is labelled with a radioactive moiety.

BRIEF DESCRIPTION OF THE DRAWINGS

- Figures 1-7 show biochemical data and sequence analysis pertaining to Examples 1, 2, 3, 7, 13, 16 and 19, respectively, with ORFs 40, 38, 44, 52, 114, 41 and 124.. M1 and M2 are molecular weight markers. Arrows indicate the position of the main recombinant product or, in Western blots, the position of the main *N.meningitidis* immunoreactive band. TP indicates *N.meningitidis* total protein extract; OMV indicates *N.meningitidis* outer membrane vesicle preparation. In bactericidal assay results: a diamond (♦) shows preimmune data; a triangle (▲) shows GST control data; a circle (●) shows data with recombinant *N.meningitidis* protein. Computer analyses show a hydrophilicity plot (upper), an antigenic index plot (middle), and an AMPHI analysis (lower). The
- 20 AMPHI program has been used to predict T-cell epitopes [Gao *et al.* (1989) *J. Immunol.* 143:3007; Roberts *et al.* (1996) *AIDS Res Hum Retrovir* 12:593; Quakyi *et al.* (1992) *Scand J Immunol* suppl.11:9) and is available in the Protean package of DNASTAR, Inc. (1228 South Park Street, Madison, Wisconsin 53715 USA).
- 25

EXAMPLES

The examples describe nucleic acid sequences which have been identified in *N.meningitidis*, along with their putative translation products. Not all of the nucleic acid sequences are complete *ie.* they encode less than the full-length wild-type protein. It is believed at present that none of the DNA
5 sequences described herein have significant homologs in *N.gonorrhoeae*.

The examples are generally in the following format:

- a nucleotide sequence which has been identified in *N.meningitidis* (strain B)
- the putative translation product of this sequence
- a computer analysis of the translation product based on database comparisons
- 10 • a corresponding gene and protein sequence identified in *N.meningitidis* (strain A)
- a description of the characteristics of the proteins which indicates that they might be suitably antigenic
- results of biochemical analysis (expression, purification, ELISA, FACS *etc.*)

The examples typically include details of sequence homology between species and strains. Proteins
15 that are similar in sequence are generally similar in both structure and function, and the homology often indicates a common evolutionary origin. Comparison with sequences of proteins of known function is widely used as a guide for the assignment of putative protein function to a new sequence and has proved particularly useful in whole-genome analyses.

Sequence comparisons were performed at NCBI (<http://www.ncbi.nlm.nih.gov>) using the
20 algorithms BLAST, BLAST2, BLASTn, BLASTp, tBLASTn, BLASTx, & tBLASTx [*eg.* see also Altschul *et al.* (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* 25:2289-3402]. Searches were performed against the following databases: non-redundant GenBank+EMBL+DDBJ+PDB sequences and non-redundant GenBank CDS translations+PDB+SwissProt+SPupdate+PIR sequences.

25 Dots within nucleotide sequences (*eg.* position 288 in Example 12) represent nucleotides which have been arbitrarily introduced in order to maintain a reading frame. In the same way, double-underlined nucleotides were removed. Lower case letters (*eg.* position 589 in Example 12) represent ambiguities which arose during alignment of independent sequencing reactions (some of

the nucleotide sequences in the examples are derived from combining the results of two or more experiments).

5 Nucleotide sequences were scanned in all six reading frames to predict the presence of hydrophobic domains using an algorithm based on the statistical studies of Esposti *et al.* [Critical evaluation of the hydropathy of membrane proteins (1990) *Eur J Biochem* 190:207-219]. These domains represent potential transmembrane regions or hydrophobic leader sequences.

Open reading frames were predicted from fragmented nucleotide sequences using the program ORFFINDER (NCBI).

10 Underlined amino acid sequences indicate possible transmembrane domains or leader sequences in the ORFs, as predicted by the PSORT algorithm (<http://www.psort.nibb.ac.jp>). Functional domains were also predicted using the MOTIFS program (GCG Wisconsin & PROSITE).

15 Various tests can be used to assess the *in vivo* immunogenicity of the proteins identified in the examples. For example, the proteins can be expressed recombinantly and used to screen patient sera by immunoblot. A positive reaction between the protein and patient serum indicates that the patient has previously mounted an immune response to the protein in question *ie.* the protein is an immunogen. This method can also be used to identify immunodominant proteins.

20 The recombinant protein can also be conveniently used to prepare antibodies *eg.* in a mouse. These can be used for direct confirmation that a protein is located on the cell-surface. Labelled antibody (*eg.* fluorescent labelling for FACS) can be incubated with intact bacteria and the presence of label on the bacterial surface confirms the location of the protein.

In particular, the following methods (A) to (S) were used to express, purify and biochemically characterise the proteins of the invention:

A) Chromosomal DNA preparation

25 *N.meningitidis* strain 2996 was grown to exponential phase in 100ml of GC medium, harvested by centrifugation, and resuspended in 5ml buffer (20% Sucrose, 50mM Tris-HCl, 50mM EDTA, pH8). After 10 minutes incubation on ice, the bacteria were lysed by adding 10ml lysis solution (50mM NaCl, 1% Na-Sarkosyl, 50µg/ml Proteinase K), and the suspension was incubated at 37°C for 2

hours. Two phenol extractions (equilibrated to pH 8) and one ChCl_3 /isoamylalcohol (24:1) extraction were performed. DNA was precipitated by addition of 0.3M sodium acetate and 2 volumes ethanol, and was collected by centrifugation. The pellet was washed once with 70% ethanol and redissolved in 4ml buffer (10mM Tris-HCl, 1mM EDTA, pH 8). The DNA
5 concentration was measured by reading the OD at 260 nm.

B) Oligonucleotide design

Synthetic oligonucleotide primers were designed on the basis of the coding sequence of each ORF, using (a) the meningococcus B sequence when available, or (b) the gonococcus/meningococcus A sequence, adapted to the codon preference usage of meningococcus as necessary. Any predicted
10 signal peptides were omitted, by deducing the 5'-end amplification primer sequence immediately downstream from the predicted leader sequence.

The 5' primers included two restriction enzyme recognition sites (*Bam*HI-*Nde*I, *Bam*HI-*Nhe*I, or *Eco*RI-*Nhe*I, depending on the gene's own restriction pattern); the 3' primers included a *Xho*I restriction site. This procedure was established in order to direct the cloning of each amplification
15 product (corresponding to each ORF) into two different expression systems: pGEX-KG (using either *Bam*HI-*Xho*I or *Eco*RI-*Xho*I), and pET21b+ (using either *Nde*I-*Xho*I or *Nhe*I-*Xho*I).

5'-end primer tail: CGCGGATCCCATATG (*Bam*HI-*Nde*I)

CGCGGATCCGCTAGC (*Bam*HI-*Nhe*I)

CCGGAATTCTAGCTAGC (*Eco*RI-*Nhe*I)

20 3'-end primer tail: CCCGCTCGAG (*Xho*I)

As well as containing the restriction enzyme recognition sequences, the primers included nucleotides which hybridised to the sequence to be amplified. The number of hybridizing nucleotides depended on the melting temperature of the whole primer, and was determined for each primer using the formulae:

25 $T_m = 4 (G+C) + 2 (A+T)$ (tail excluded)

$T_m = 64.9 + 0.41 (\% \text{ GC}) - 600/N$ (whole primer)

The average melting temperature of the selected oligos were 65-70°C for the whole oligo and 50-55°C for the hybridising region alone.

Table I shows the forward and reverse primers used for each amplification. Oligos were synthesized by a Perkin Elmer 394 DNA/RNA Synthesizer, eluted from the columns in 2ml NH_4OH , and deprotected by 5 hours incubation at 56°C . The oligos were precipitated by addition of 0.3M Na-Acetate and 2 volumes ethanol. The samples were then centrifuged and the pellets resuspended in either 100 μl or 1ml of water. OD_{260} was determined using a Perkin Elmer Lambda Bio spectrophotometer and the concentration was determined and adjusted to 2-10pmol/ μl .

C) Amplification

The standard PCR protocol was as follows: 50-200ng of genomic DNA were used as a template in the presence of 20-40 μM of each oligo, 400-800 μM dNTPs solution, 1x PCR buffer (including 1.5mM MgCl_2), 2.5 units *TaqI* DNA polymerase (using Perkin-Elmer AmpliTaq, GIBCO Platinum, Pwo DNA polymerase, or Tahara Shuzo Taq polymerase).

In some cases, PCR was optimised by the addition of 10 μl DMSO or 50 μl 2M betaine.

After a hot start (adding the polymerase during a preliminary 3 minute incubation of the whole mix at 95°C), each sample underwent a double-step amplification: the first 5 cycles were performed using as the hybridization temperature the one of the oligos excluding the restriction enzymes tail, followed by 30 cycles performed according to the hybridization temperature of the whole length oligos. The cycles were followed by a final 10 minute extension step at 72°C .

The standard cycles were as follows:

	Denaturation	Hybridisation	Elongation
First 5 cycles	30 seconds 95°C	30 seconds $50-55^\circ\text{C}$	30-60 seconds 72°C
Last 30 cycles	30 seconds 95°C	30 seconds $65-70^\circ\text{C}$	30-60 seconds 72°C

The elongation time varied according to the length of the ORF to be amplified.

The amplifications were performed using either a 9600 or a 2400 Perkin Elmer GeneAmp PCR System. To check the results, 1/10 of the amplification volume was loaded onto a 1-1.5% agarose gel and the size of each amplified fragment compared with a DNA molecular weight marker.

5 The amplified DNA was either loaded directly on a 1% agarose gel or first precipitated with ethanol and resuspended in a suitable volume to be loaded on a 1% agarose gel. The DNA fragment corresponding to the right size band was then eluted and purified from gel, using the Qiagen Gel Extraction Kit, following the instructions of the manufacturer. The final volume of the DNA fragment was 30µl or 50µl of either water or 10mM Tris, pH 8.5.

D) Digestion of PCR fragments

10 The purified DNA corresponding to the amplified fragment was split into 2 aliquots and double-digested with:

- *NdeI/XhoI* or *NheI/XhoI* for cloning into pET-21b+ and further expression of the protein as a C-terminus His-tag fusion
- *BamHI/XhoI* or *EcoRI/XhoI* for cloning into pGEX-KG and further expression of the
15 protein as N-terminus GST fusion.
- *EcoRI/PstI*, *EcoRI/SalI*, *SalI/PstI* for cloning into pGex-His and further expression of the protein as N-terminus His-tag fusion

20 Each purified DNA fragment was incubated (37°C for 3 hours to overnight) with 20 units of each restriction enzyme (New England Biolabs) in a either 30 or 40µl final volume in the presence of the appropriate buffer. The digestion product was then purified using the QIAquick PCR purification kit, following the manufacturer's instructions, and eluted in a final volume of 30 or 50µl of either water or 10mM Tris-HCl, pH 8.5. The final DNA concentration was determined by 1% agarose gel electrophoresis in the presence of titrated molecular weight marker.

E) Digestion of the cloning vectors (pET22B, pGEX-KG, pTRC-His A, and pGex-His)

25 10µg plasmid was double-digested with 50 units of each restriction enzyme in 200µl reaction volume in the presence of appropriate buffer by overnight incubation at 37°C. After loading the

whole digestion on a 1% agarose gel, the band corresponding to the digested vector was purified from the gel using the Qiagen QIAquick Gel Extraction Kit and the DNA was eluted in 50µl of 10mM Tris-HCl, pH 8.5. The DNA concentration was evaluated by measuring OD₂₆₀ of the sample, and adjusted to 50µg/µl. 1µl of plasmid was used for each cloning procedure.

- 5 The vector pGEX-His is a modified pGEX-2T vector carrying a region encoding six histidine residues upstream to the thrombin cleavage site and containing the multiple cloning site of the vector pTRC99 (Pharmacia).

F) Cloning

- 10 The fragments corresponding to each ORF, previously digested and purified, were ligated in both pET22b and pGEX-KG. In a final volume of 20µl, a molar ratio of 3:1 fragment/vector was ligated using 0.5µl of NEB T4 DNA ligase (400 units/µl), in the presence of the buffer supplied by the manufacturer. The reaction was incubated at room temperature for 3 hours. In some experiments, ligation was performed using the Boehringer "Rapid Ligation Kit", following the manufacturer's instructions.

- 15 In order to introduce the recombinant plasmid in a suitable strain, 100µl *E. coli* DH5 competent cells were incubated with the ligase reaction solution for 40 minutes on ice, then at 37°C for 3 minutes, then, after adding 800µl LB broth, again at 37°C for 20 minutes. The cells were then centrifuged at maximum speed in an Eppendorf microfuge and resuspended in approximately 200µl of the supernatant. The suspension was then plated on LB ampicillin (100mg/ml).

- 20 The screening of the recombinant clones was performed by growing 5 randomly-chosen colonies overnight at 37°C in either 2ml (pGEX or pTC clones) or 5ml (pET clones) LB broth + 100µg/ml ampicillin. The cells were then pelleted and the DNA extracted using the Qiagen QIAprep Spin Miniprep Kit, following the manufacturer's instructions, to a final volume of 30µl. 5µl of each individual miniprep (approximately 1g) were digested with either *NdeI/XhoI* or *BamHI/XhoI* and the whole digestion loaded onto a 1-1.5% agarose gel (depending on the expected insert size), in parallel with the molecular weight marker (1Kb DNA Ladder, GIBCO). The screening of the positive clones was made on the base of the correct insert size.
- 25

G) Expression

Each ORF cloned into the expression vector was transformed into the strain suitable for expression of the recombinant protein product. 1µl of each construct was used to transform 30µl of *E. coli* BL21 (pGEX vector), *E. coli* TOP 10 (pTRC vector) or *E. coli* BL21-DE3 (pET vector), as described
5 above. In the case of the pGEX-His vector, the same *E. coli* strain (W3110) was used for initial cloning and expression. Single recombinant colonies were inoculated into 2ml LB+Amp (100µg/ml), incubated at 37°C overnight, then diluted 1:30 in 20ml of LB+Amp (100µg/ml) in 100ml flasks, making sure that the OD₆₀₀ ranged between 0.1 and 0.15. The flasks were incubated at 30°C into gyratory water bath shakers until OD indicated exponential growth suitable for
10 induction of expression (0.4-0.8 OD for pET and pTRC vectors; 0.8-1 OD for pGEX and pGEX-His vectors). For the pET, pTRC and pGEX-His vectors, the protein expression was induced by addition of 1mM IPTG, whereas in the case of pGEX system the final concentration of IPTG was 0.2mM. After 3 hours incubation at 30°C, the final concentration of the sample was checked by OD. In order to check expression, 1ml of each sample was removed, centrifuged in a microfuge,
15 the pellet resuspended in PBS, and analysed by 12% SDS-PAGE with Coomassie Blue staining. The whole sample was centrifuged at 6000g and the pellet resuspended in PBS for further use.

H) GST-fusion proteins large-scale purification.

A single colony was grown overnight at 37°C on LB+Amp agar plate. The bacteria were inoculated into 20ml of LB+Amp liquid culture in a water bath shaker and grown overnight. Bacteria were
20 diluted 1:30 into 600ml of fresh medium and allowed to grow at the optimal temperature (20-37°C) to OD₅₅₀ 0.8-1. Protein expression was induced with 0.2mM IPTG followed by three hours incubation. The culture was centrifuged at 8000rpm at 4°C. The supernatant was discarded and the bacterial pellet was resuspended in 7.5ml cold PBS. The cells were disrupted by sonication on ice for 30 sec at 40W using a Branson sonifier B-15, frozen and thawed twice and centrifuged again.
25 The supernatant was collected and mixed with 150µl Glutathione-Sepharose 4B resin (Pharmacia) (previously washed with PBS) and incubated at room temperature for 30 minutes. The sample was centrifuged at 700g for 5 minutes at 4°C. The resin was washed twice with 10ml cold PBS for 10 minutes, resuspended in 1ml cold PBS, and loaded on a disposable column. The resin was washed twice with 2ml cold PBS until the flow-through reached OD₂₈₀ of 0.02-0.06. The GST-fusion
30 protein was eluted by addition of 700µl cold Glutathione elution buffer (10mM reduced

glutathione, 50mM Tris-HCl) and fractions collected until the OD_{280} was 0.1. 21 μ l of each fraction were loaded on a 12% SDS gel using either Biorad SDS-PAGE Molecular weight standard broad range (M1) (200, 116.25, 97.4, 66.2, 45, 31, 21.5, 14.4, 6.5 kDa) or Amersham Rainbow Marker (M2) (220, 66, 46, 30, 21.5, 14.3 kDa) as standards. As the MW of GST is 26kDa, this value must
5 be added to the MW of each GST-fusion protein.

I) His-fusion solubility analysis

To analyse the solubility of the His-fusion expression products, pellets of 3ml cultures were resuspended in buffer M1 [500 μ l PBS pH 7.2]. 25 μ l lysozyme (10mg/ml) was added and the bacteria were incubated for 15 min at 4°C. The pellets were sonicated for 30 sec at 40W using a
10 Branson sonifier B-15, frozen and thawed twice and then separated again into pellet and supernatant by a centrifugation step. The supernatant was collected and the pellet was resuspended in buffer M2 [8M urea, 0.5M NaCl, 20mM imidazole and 0.1M NaH_2PO_4] and incubated for 3 to 4 hours at 4°C. After centrifugation, the supernatant was collected and the pellet was resuspended
15 in buffer M3 [6M guanidinium-HCl, 0.5M NaCl, 20mM imidazole and 0.1M NaH_2PO_4] overnight at 4°C. The supernatants from all steps were analysed by SDS-PAGE.

J) His-fusion large-scale purification.

A single colony was grown overnight at 37°C on a LB + Amp agar plate. The bacteria were inoculated into 20ml of LB+Amp liquid culture and incubated overnight in a water bath shaker. Bacteria were diluted 1:30 into 600ml fresh medium and allowed to grow at the optimal
20 temperature (20-37°C) to OD_{550} 0.6-0.8. Protein expression was induced by addition of 1mM IPTG and the culture further incubated for three hours. The culture was centrifuged at 8000rpm at 4°C, the supernatant was discarded and the bacterial pellet was resuspended in 7.5ml of either (i) cold buffer A (300mM NaCl, 50mM phosphate buffer, 10mM imidazole, pH 8) for soluble proteins or (ii) buffer B (urea 8M, 10mM Tris-HCl, 100mM phosphate buffer, pH 8.8) for insoluble proteins.

25 The cells were disrupted by sonication on ice for 30 sec at 40W using a Branson sonifier B-15, frozen and thawed two times and centrifuged again.

For insoluble proteins, the supernatant was stored at -20°C, while the pellets were resuspended in 2ml buffer C (6M guanidine hydrochloride, 100mM phosphate buffer, 10mM Tris-HCl, pH 7.5) and treated in a homogenizer for 10 cycles. The product was centrifuged at 13000rpm for 40 minutes.

Supernatants were collected and mixed with 150µl Ni²⁺-resin (Pharmacia) (previously washed with either buffer A or buffer B, as appropriate) and incubated at room temperature with gentle agitation for 30 minutes. The sample was centrifuged at 700g for 5 minutes at 4°C. The resin was washed twice with 10ml buffer A or B for 10 minutes, resuspended in 1ml buffer A or B and loaded on a disposable column. The resin was washed at either (i) 4°C with 2ml cold buffer A or (ii) room temperature with 2ml buffer B, until the flow-through reached OD₂₈₀ of 0.02-0.06.

The resin was washed with either (i) 2ml cold 20mM imidazole buffer (300mM NaCl, 50mM phosphate buffer, 20mM imidazole, pH 8) or (ii) buffer D (urea 8M, 10mM Tris-HCl, 100mM phosphate buffer, pH 6.3) until the flow-through reached the O.D₂₈₀ of 0.02-0.06. The His-fusion protein was eluted by addition of 700µl of either (i) cold elution buffer A (300mM NaCl, 50mM phosphate buffer, 250mM imidazole, pH 8) or (ii) elution buffer B (urea 8M, 10mM Tris-HCl, 100mM phosphate buffer, pH 4.5) and fractions collected until the O.D₂₈₀ was 0.1. 21µl of each fraction were loaded on a 12% SDS gel.

K) His-fusion proteins renaturation

10% glycerol was added to the denatured proteins. The proteins were then diluted to 20µg/ml using dialysis buffer I (10% glycerol, 0.5M arginine, 50mM phosphate buffer, 5mM reduced glutathione, 0.5mM oxidised glutathione, 2M urea, pH 8.8) and dialysed against the same buffer at 4°C for 12-14 hours. The protein was further dialysed against dialysis buffer II (10% glycerol, 0.5M arginine, 50mM phosphate buffer, 5mM reduced glutathione, 0.5mM oxidised glutathione, pH 8.8) for 12-14 hours at 4°C. Protein concentration was evaluated using the formula:

$$\text{Protein (mg/ml)} = (1.55 \times \text{OD}_{280}) - (0.76 \times \text{OD}_{260})$$

L) His-fusion large-scale purification

500ml of bacterial cultures were induced and the fusion proteins were obtained soluble in buffer M1, M2 or M3 using the procedure described above. The crude extract of the bacteria was loaded

onto a Ni-NTA superflow column (Qiagen) equilibrated with buffer M1, M2 or M3 depending on the solubilization buffer of the fusion proteins. Unbound material was eluted by washing the column with the same buffer. The specific protein was eluted with the corresponding buffer containing 500mM imidazole and dialysed against the corresponding buffer without imidazole.

5 After each run the columns were sanitized by washing with at least two column volumes of 0.5 M sodium hydroxide and reequilibrated before the next use.

M) Mice immunisations

20µg of each purified protein were used to immunise mice intraperitoneally. In the case of ORF 44, CD1 mice were immunised with Al(OH)₃ as adjuvant on days 1, 21 and 42, and immune response

10 was monitored in samples taken on day 56. For ORF 40, CD1 mice were immunised using Freund's adjuvant, rather than Al(OH)₃, and the same immunisation protocol was used, except that the immune response was measured on day 42, rather than 56. Similarly, for ORF 38, CD1 mice were immunised with Freund's adjuvant, but the immune response was measured on day 49.

N) ELISA assay (sera analysis)

15 The acapsulated MenB M7 strain was plated on chocolate agar plates and incubated overnight at 37°C. Bacterial colonies were collected from the agar plates using a sterile dracon swab and inoculated into 7ml of Mueller-Hinton Broth (Difco) containing 0.25% Glucose. Bacterial growth was monitored every 30 minutes by following OD₆₂₀. The bacteria were let to grow until the OD reached the value of 0.3-0.4. The culture was centrifuged for 10 minutes at 10000rpm. The

20 supernatant was discarded and bacteria were washed once with PBS, resuspended in PBS containing 0.025% formaldehyde, and incubated for 2 hours at room temperature and then overnight at 4°C with stirring. 100µl bacterial cells were added to each well of a 96 well Greiner plate and incubated overnight at 4°C. The wells were then washed three times with PBT washing buffer (0.1% Tween-20 in PBS). 200µl of saturation buffer (2.7% Polyvinylpyrrolidone 10 in

25 water) was added to each well and the plates incubated for 2 hours at 37°C. Wells were washed three times with PBT. 200µl of diluted sera (Dilution buffer: 1% BSA, 0.1% Tween-20, 0.1% NaN₃ in PBS) were added to each well and the plates incubated for 90 minutes at 37°C. Wells were washed three times with PBT. 100µl of HRP-conjugated rabbit anti-mouse (Dako) serum diluted 1:2000 in dilution buffer were added to each well and the plates were incubated for 90 minutes at

37°C. Wells were washed three times with PBT buffer. 100µl of substrate buffer for HRP (25ml of citrate buffer pH5, 10mg of O-phenildiamine and 10µl of H₂O) were added to each well and the plates were left at room temperature for 20 minutes. 100µl H₂SO₄ was added to each well and OD₄₉₀ was followed. The ELISA was considered positive when OD₄₉₀ was 2.5 times the respective pre-immune sera.

O) FACScan bacteria Binding Assay procedure.

The acapsulated MenB M7 strain was plated on chocolate agar plates and incubated overnight at 37°C. Bacterial colonies were collected from the agar plates using a sterile dracon swab and inoculated into 4 tubes containing 8ml each Mueller-Hinton Broth (Difco) containing 0.25% glucose. Bacterial growth was monitored every 30 minutes by following OD₆₂₀. The bacteria were let to grow until the OD reached the value of 0.35-0.5. The culture was centrifuged for 10 minutes at 4000rpm. The supernatant was discarded and the pellet was resuspended in blocking buffer (1% BSA, 0.4% NaN₃) and centrifuged for 5 minutes at 4000rpm. Cells were resuspended in blocking buffer to reach OD₆₂₀ of 0.07. 100µl bacterial cells were added to each well of a Costar 96 well plate. 100µl of diluted (1:200) sera (in blocking buffer) were added to each well and plates incubated for 2 hours at 4°C. Cells were centrifuged for 5 minutes at 4000rpm, the supernatant aspirated and cells washed by addition of 200µl/well of blocking buffer in each well. 100µl of R-Phicoerytrin conjugated F(ab)₂ goat anti-mouse, diluted 1:100, was added to each well and plates incubated for 1 hour at 4°C. Cells were spun down by centrifugation at 4000rpm for 5 minutes and washed by addition of 200µl/well of blocking buffer. The supernatant was aspirated and cells resuspended in 200µl/well of PBS, 0.25% formaldehyde. Samples were transferred to FACScan tubes and read. The condition for FACScan setting were: FL1 on, FL2 and FL3 off; FSC-H threshold:92; FSC PMT Voltage: E 02; SSC PMT: 474; Amp. Gains 7.1; FL-2 PMT: 539; compensation values: 0.

25 P) OMV preparations

Bacteria were grown overnight on 5 GC plates, harvested with a loop and resuspended in 10 ml 20mM Tris-HCl. Heat inactivation was performed at 56°C for 30 minutes and the bacteria disrupted by sonication for 10 minutes on ice (50% duty cycle, 50% output). Unbroken cells were removed by centrifugation at 5000g for 10 minutes and the total cell envelope fraction recovered by centrifugation

at 50000g at 4°C for 75 minutes. To extract cytoplasmic membrane proteins from the crude outer membranes, the whole fraction was resuspended in 2% sarkosyl (Sigma) and incubated at room temperature for 20 minutes. The suspension was centrifuged at 10000g for 10 minutes to remove aggregates, and the supernatant further ultracentrifuged at 50000g for 75 minutes to pellet the outer membranes. The outer membranes were resuspended in 10mM Tris-HCl, pH8 and the protein concentration measured by the Bio-Rad Protein assay, using BSA as a standard.

Q) Whole Extracts preparation

Bacteria were grown overnight on a GC plate, harvested with a loop and resuspended in 1ml of 20mM Tris-HCl. Heat inactivation was performed at 56°C for 30 minutes.

10 R) Western blotting

Purified proteins (500ng/lane), outer membrane vesicles (5µg) and total cell extracts (25µg) derived from MenB strain 2996 were loaded on 15% SDS-PAGE and transferred to a nitrocellulose membrane. The transfer was performed for 2 hours at 150mA at 4°C, in transferring buffer (0.3 % Tris base, 1.44 % glycine, 20% methanol). The membrane was saturated by overnight incubation at 4°C in saturation buffer (10% skimmed milk, 0.1% Triton X100 in PBS). The membrane was washed twice with washing buffer (3% skimmed milk, 0.1% Triton X100 in PBS) and incubated for 2 hours at 37°C with mice sera diluted 1:200 in washing buffer. The membrane was washed twice and incubated for 90 minutes with a 1:2000 dilution of horseradish peroxidase labelled anti-mouse Ig. The membrane was washed twice with 0.1% Triton X100 in PBS and developed with the Opti-4CN Substrate Kit (Bio-Rad). The reaction was stopped by adding water.

S) Bactericidal assay

MC58 strain was grown overnight at 37°C on chocolate agar plates. 5-7 colonies were collected and used to inoculate 7ml Mueller-Hinton broth. The suspension was incubated at 37°C on a nutator and let to grow until OD₆₂₀ was 0.5-0.8. The culture was aliquoted into sterile 1.5ml Eppendorf tubes and centrifuged for 20 minutes at maximum speed in a microfuge. The pellet was washed once in Gey's buffer (Gibco) and resuspended in the same buffer to an OD₆₂₀ of 0.5, diluted 1:20000 in Gey's buffer and stored at 25°C.

50µl of Gey's buffer/1% BSA was added to each well of a 96-well tissue culture plate. 25µl of diluted mice sera (1:100 in Gey's buffer/0.2% BSA) were added to each well and the plate incubated at 4°C. 25µl of the previously described bacterial suspension were added to each well. 25µl of either heat-inactivated (56°C waterbath for 30 minutes) or normal baby rabbit complement were added to each well. Immediately after the addition of the baby rabbit complement, 22µl of each sample/well were plated on Mueller-Hinton agar plates (time 0). The 96-well plate was incubated for 1 hour at 37°C with rotation and then 22µl of each sample/well were plated on Mueller-Hinton agar plates (time 1). After overnight incubation the colonies corresponding to time 0 and time 1 hour were counted.

10 **Table II** gives a summary of the cloning, expression and purification results.

Example 1

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 1>:

```

1  ..ACACTGTTGT TTGCAACGGT TCAGGCAAGT GCTAACCAAT GAAGAGCAAG
51  AAGAAGATTT ATATTTAGAC CCCGTACAAC GCACTGTTGC CGTGTGATA
15  101  GTCAATTCCG ATAAAGAAGG CACGGGAGAA AAAGAAAAAG TAGAAGAAAA
151  TTCAGATTGG GCAGTATATT TCAACGAGAA AGGAGTACTA ACAGCCAGAG
201  AAATCACCyT CAAAGCCGGC GACAACCTGA AAATCAAACA AAACGGCACA
251  AACTTCACCT ACTCGCTGAA AAAAGACCTC AcAGATCTGA CCAGTGTGG
20  301  AACTGAAAAA TTATCGTTTA GCGCAAACGG CAATAAAGTC AACATcACAA
351  GCGACACCAA AGGCTTGAAT TTTGCGAAAG AAACGGCTGG sACGAACGgC
401  GACACCACGG TTCATCTGAA CGGTATTGGT TCGACTTTGA CCGATACGCT
451  GCTGAATACC GGAGCGACCA CAAACGTAAC CAACGACAAC GTTACCGATG
501  ACGAGAAAAA ACGTGCGGCA AGCGTTAAAG ACGTATTAAA CGCTGGCTGG
551  AACATTAAAG GCGTTAAACC CGGTACAACA GCTTCCGATA ACGTTGATTT
25  601  CGTCCGCACT TACGACACAG TCGAGTTCTT GAGCGCAGAT ACGAAAACAA
651  CGACTGTTAA TGTGAAAGC AAAGACAACG GCAAGAAAAAC CGAAGTTAAA
701  ATCGGTGCGA AGACTTCTGT TATTAAAGAA AAAGAC...
```

This corresponds to the amino acid sequence <SEQ ID 2; ORF40>:

```

1  ..TLLFATVQAS ANQEEQEEDL YLDPVQRTVA VLIVNSDKEG TGEKEKVEEN
30  51  SDWAVYFNEK GVLTAIREITX KAGDNLKIKQ NGTNFTYSLK KDLDLTLSVG
101  TEKLSFSANG NKNVITS DTK GLNFAKETAG TNGDITVHLN GIGSTLTDTL
151  LNTGATTNVT NDNVTDDEKK RAASVKDVLN AGWNIKVVP GTTASDNVDF
201  VRTYDTVEFL SADTKTTTVN VESKDNGKKT EVKIGAKTSV IKEKD...
```

Further work revealed the complete DNA sequence <SEQ ID 3>:

```

35  1  ATGAACAAAA TATACCGCAT CATTGGAAT AGTGCCCTCA ATGCCTGGGT
51  CGTCGTATCC GAGCTCACAC GCAACCACAC CAAACGCGCC TCCGCAACCG
101  TGAAGACCGC CGTATTGGCG AACTGTTGT TTGCAACGGT TCAGGCAAGT
151  GCTAACCAATG AAGAGCAAGA AGAAGATTTA TATTTAGACC CGTACAACG
40  201  CACTGTTGCC GTGTGTAGT TCAATTCCGA TAAAGAAGGC ACGGAGAGAA
251  AAGAAAAAGT AGAAGAAAT TCAGATTGGG CAGTATATTT CAACGAGAAA
301  GGAGTACTAA CAGCCAGAGA AATCACCCTC AAAGCCGGCG ACAACCTGAA
351  AATCAAACAA AACGGCACA AACTCACCTA CTCGCTGAAA AAAGACCTCA
401  CAGATCTGAC CAGTGTGGA ACTGAAAAAT TATCGTTTAG CGCAACGGC
45  451  AATAAAGTCA ACATCACAAG CGACACCAA GGCTTGAATT TTGCGAAAGA
501  AACGGCTGGG ACGAACGGCG ACACCACGGT TCATCTGAAC GGTATTGGTT
551  CGACTTTGAC CGATACGCTG CTGAATACCG GAGCGACCAC AAACGTAACC
601  AACGACAACG TTACCGATGA CGAGAAAAAA CGTGCGGCAA GCGTTAAAGA
```


5 651 CGTATTAAAC GCTGGCTGGA ACATTAAAGG CGTTAAACCC GGTACAACAG
 701 CTTCCGATAA CGTTGATTTC GTCCGCACTT ACGACACAGT CGAGTTCTTG
 751 AGCGCAGATA CGAAAACAAC GACTGTTAAT GTGGAAAGCA AAGACAACGG
 801 CAAGAAAACC GAAGTTAAAA TCGGTGCGAA GACTTCTGTT ATTAAGAAAA
 851 AAGACGGTAA GTTGGTTACT GGTAAAGACA AAGGCGAGAA TGGTTCTTCT
 901 ACAGACGAAG GCGAAGGCTT AGTGACTGCA AAAGAAGTGA TTGATGCAGT
 951 AAACAAGGCT GGTGAGAGAA TGAAAACAAC AACCGCTAAT GGTCAAACAG
 1001 GTCAAGCTGA CAAGTTTGAA ACCGTTACAT CAGGCACAAA TGTAACCTTT
 1051 GCTAGTGGTA AAGGTACAAC TGCGACTGTA AGTAAAGATG ATCAAGGCAA
 1101 CATCACTGTT ATGTATGATG TAAATGTCGG CGATGCCCTA AACGTCAATC
 1151 AGCTGCAAAA CAGCGGTTGG AATTTGGATT CCAAAGCGGT TGCAGGTTCT
 1201 TCGGGCAAAG TCATCAGCGG CAATGTTTCG CCGAGCAAGG GAAAGATGGA
 1251 TGAAACCGTC AACATTAATG CCGGCAACAA CATCGAGATT ACCCGCAACG
 1301 GTAAAAATAT CGACATCGCC ACTTCGATGA CCGGCGAGTT TTCCAGCGTT
 1351 TCGCTCGGCG CGGGGGCGGA TGCGCCCACT TTGAGCGTGG ATGGGCGGCG
 1401 ATTGAATGTC GGCAGCAAGA AGGACAACAA ACCCGTCCGC ATTACCAATG
 1451 TCGCCCCGGG CGTTAAAGAG GGGGATGTTA CAAACGTCGC ACAACTTAAA
 1501 GCGGTGGGCG AAAACTTGAA CAACCGCATC GACAATGTGG ACGGCAACGC
 1551 GCGTGGGCGC ATCGCCCAAG CGATTGCAAC CGCAGGTCGT GTTCAGGCGT
 20 1601 ATTTGCCCCG CAAGAGTATG ATGGCGATCG GCGGCGGCAC TTATCGCGGC
 1651 GAAGCCGGTT ACGCCATCGG CTACTCCAGT ATTTCCGACG GCGGAAATTG
 1701 GATTATCAAA GGCACGGCTT CCGGCAATTC GCGCGGCCAT TTCGGTGCTT
 1751 CCGCATCTGT CGGTTATCAG TGGTAA

This corresponds to the amino acid sequence <SEQ ID 4; ORF40-1>:

25 1 MNKIYRIIWN SALNAWVVVS ELTRNHTKRA SATVKTAFLA TLLFATVQAS
 51 ANNEEQEEDL YLDPVQRTVA VLVNSDKEG TGEKEKVEEN SDWAVYFNEK
 101 GVLTAIREITL KAGDNLKIKQ NGTNFTYSLK KDLTDLTSVG TEKLSFSANG
 151 NKVNITSDTK GLNFAKETAG TNGDITVHLN GIGSTLDTL LNTGATTNVT
 30 201 NDNVTDDEKK RAASVKDVLN AGWNIKGVKP GTTASDNVDF VRTYDTVEFL
 251 SADTKTTTTVN VESKDNKKKT EVKIGAKTSV IKEKDGKLVF GKDKGENGSS
 301 TDEGEGLVTA KEVIDAVNKA GWRMKTITAN GQTGQADKFE TVTSGTNVTF
 351 ASKGTTATV SKDDQGNITV MYDVNVGDAL NVNQLQNSGW NLDKAVAGS
 401 SGKVISGNVS PSKGKMDTV NINAGNNIEI TRNGKNIDIA TSMTPOFSSV
 451 SLGAGADAPT LSVGDALNV GSKKDNKPVR ITNVAPGVKE GDVTNVAQLK
 35 501 GVAQNLNRI DNVGNARAG IAQAIATAGL VQAYLPGKSM MAIGGGTYRG
 551 EAGYAIGYSS ISDGGNWIIC GTASGNSRGH FGASASVGYQ W*

Further work identified the corresponding gene in strain A of *N.meningitidis* <SEQ ID 5>:

40 1 ATGAACAAAA TATACCGCAT CATTGGAAT AGTGCCCTCA ATGCCTGNCT
 51 CGCCGTATCC GAGCTCACAC GCAACCACAC CAAACGCGCC TCCGCAACCG
 101 TGAAGACCGC CGATTGGCG ACACTGTTGT TTGCAACGGT TCAGGCGAAT
 151 GCTACCGATG AAGATGAAGA AGAAGAGTTA GAATCCGTAC AACGCTCTGT
 201 CGTAGGGAGC ATTCAAGCCA GTATGGAAGG CAGCGGCGAA TTGGAACGA
 251 TATCATTATC AATGACTAAC GACAGCAAGG AATTGTAGA CCCATACATA
 301 GTAGTTACCC TCAAGCCCG CGACAACCTG AAAATCAAAC AAAACACCAA
 45 351 TGAAACACCC AATGCCAGTA GCTTCACTA CTCGCTGAAA AAAGACCTCA
 401 CAGGCCTGAT CAATGTTGAN ACTGAAAAAT TATCGTTTGG CGCAACCGGC
 451 AAGAAAGTCA ACATCATAAG CGACACCAA GGCTTGAATT TCGCGAAAGA
 501 AACGGCTGGG ACGAACGGCG ACACCACGGT TCATCTGAAC GGTATCGGTT
 551 CGACTTTGAC CGATACGCTT GCGGGTTCTT CTGCTTCTCA CGTTGATGCG
 50 601 GGTAACCNAA GTACACATTA CACTCGTGCA GCAAGTATTA AGGATGTGTT
 651 GAATGCGGGT TGAATATTA AGGGTGTTAA ANNNGGCTCA ACAACTGGTC
 701 AATCAGAAAA TGTCGATTTC GTCCGCACTT ACGACACAGT CGAGTTCTTG
 751 AGCGCAGATA CGNAAACAAC GACNGTTAAT GTGGAAAGCA AAGACAACGG
 801 CAAGAGAACC GAAGTTAAAA TCGGTGCGAA GACTTCTGTT ATTAAGAAAA
 55 851 AAGACGGTAA GTTGGTTACT GGTAAAGGCA AAGGCGAGAA TGGTTCTTCT
 901 ACAGACGAAG GCGAAGGCTT AGTGACTGCA AAAGAAGTGA TTGATGCAGT
 951 AAACAAGGCT GGTGAGAGAA TGAAAACAAC AACCGCTAAT GGTCAAACAG
 1001 GTCAAGCTGA CAAGTTTGAA ACCGTTACAT CAGGCACAAA TGTAACCTTT
 1051 GCTAGTGGTA AAGGTACAAC TGCGACTGTA AGTAAAGATG ATCAAGGCAA
 60 1101 CATCACTGTT ATGTATGATG TAAATGTCGG CGATGCCCTA AACGTCAATC
 1151 AGCTGCAAAA CAGCGGTTGG AATTTGGATT CCAAAGCGGT TGCAGGTTCT
 1201 TCGGGCAAAG TCATCAGCGG CAATGTTTCG CCGAGCAAGG GAAAGATGGA
 1251 TGAAACCGTC AACATTAATG CCGGCAACAA CATCGAGATT AGCCGCAACG
 1301 GTAAAAATAT CGACATCGCC ACTTCGATGG CGCCGCGAGT TTCCAGCGTT
 65 1351 TCGCTCGGCG CGGGGGCAGA TGCGCCCACT TTAAGCGTGG ATGACGAGGG
 1401 CGCGTTGAAT GTCGGCAGCA AGGATGCCAA CAAACCCGTC CGCATTACCA

This encodes a protein having amino acid sequence <SEQ ID 6; ORF40a>:

10

15

20

1	MNKIYRIIWN	SALNAXVAVS	ELTRNHTKRA	SATVKTAVLA	TLLFATVQAN
51	ATDEDEEEEL	ESVQRSVGS	IQASMEGSGE	LETISLSMTN	DSKEFVDPII
101	VVTLKAGDNL	KIKQNTENT	NASSFTYSLK	KDLTGLINXV	TEKLSFGANG
151	KKVNIISDPT	GLNFAKETAG	TNGDFTYVHLN	GIGSTLVDFL	AGSSASHVDA
201	GNXSTHYTRA	ASIKDVLNAG	WNIKGVKXGS	TTGQSENVDF	VRTYDTEVEFL
251	SADTXTTTVN	VESKDNKRT	EVKIGAKTSV	IKEKDGKLVF	GKKGKENGSS
301	TDEGEGLTVA	KEVIDAVNKA	GWRMKTTTAN	GQTQADKFFE	TVTSGNTVTF
351	ASGKGTATV	SKDDQGNITV	MYDVNVGDAL	NVNQLQNSGW	NLDSKAVAGS
401	SGKVISGNVS	PSKGKMDETV	NINAGNNIEI	SRNGKNIDIA	TSMAPQFSSV
451	SLGAGADAPT	LSVDDEGALN	VGSKDANKPV	RITNVAPGVK	XGDVTNVXQL
501	KGVAQNLNNR	IDNVDDGNARA	GIAQAIATAG	LVQAYLPGKS	MMAIIGGGTYR
551	GEAGYATGYS	SISDGGNIWI	KGTASGNSRG	HFGASASVGY	OW*

The originally-identified partial strain B sequence (ORF40) shows 65.7% identity over a 254aa overlap with ORF40a:

[illegible]

The complete strain B sequence (ORF40-1) and ORF40a show 83.7% identity in 601 aa overlap:

```

55      orf40-1.pep      10      20      30      40      50      60
                        MNKIYRIIWNSALNAWVVVSELTRNHTKRASATVKTAVLATLLFATVQASANNEEQEEDL
                        |||||:|||||:|||||:|||||:|||||:|||||:|||||:|||||:|||||:|||||:|||||
      orf40a            MNKIYRIIWNSALNAXVAVSELTRNHTKRASATVKTAVLATLLFATVQANATDEDEEEE
                        10      20      30      40      50      60

60      orf40-1.pep      70      80      90      100     110     119
                        YLDPVQRTVAVLIVNSDKEGTGEKEKVEEN-SDWAVVFNEKGVLTAREITLKAGDNLKIK
                        :|||:|:||||:||||:||||:||||:||||:||||:||||:||||:||||:||||:||||:||||:

```

55 Computer analysis of these amino acid sequences gave the following results:

Homology with Hsf protein encoded by the type b surface fibrils locus of *H. influenzae*
(accession number U41852)

ORF40 and Hsf protein show 54% aa identity in 251 aa overlap:

60	Orf40	1	TLLFATVQASANQEEQEEDLYLDPVQRTVAVLIVNSDXXXXXXXXXXXXNSDWAVFYNEK	60
	Hsf	41	TLLFATVQA+A E++E LDPV RT VL +SD NS+W +YF+ K	95
65	Orf40	61	GVLTAAREITXKAGDNLKIKQN-----GTNFTYSLKKDLTDLTSVGTTEKLSFSANGNKVN	114
	Hsf	96	GVLKAGAITLKAGDNLKIKQNTDESTNASSFTYSLKKDLTDLTSVATEKLSFGANGDKVD	155

Orf40 115 ITSDTKGLNFAKETAGTNGDTTVHLNIGIGSTLTDTLLNTGAXXXXXXXXXXXXXEKKRAAS 174
 ITSD GL AK G+ VHLNG+ STL D + NTG EK RAA+
 Hsf 156 ITSDANGLKLAK-----TGNGNVHLNGLDSTLPDAVTNTGVLSSSFTPNVDV-EKTRAAT 209
 Orf40 175 VKDVLNAGWNIKGKPGTTASDNVDFVRTYDTVEFLSADTKTTTVNVESKDNGKKTEVKI 234
 VKDVLNAGWNIKG K ++VD V Y+ VEF++ D T V + +K+NGK TEVK
 Hsf 210 VKDVLNAGWNIKGAKTAGGNVESVDLVSAYNNVEFITGDKNTLDVVLTAKEKNGKTEVKF 269
 Orf40 235 GAKTSVIKEKD 245
 KTSVIKEKD
 Hsf 270 TPKTSVIKEKD 280

ORF40a also shows homology to Hsf:

gi|1666683 (U41852) hsf gene product [Haemophilus influenzae] Length = 2353
 Score = 153 (67.7 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 33/36 (91%), Positives = 34/36 (94%)

Query: 16 VAVSELTRNHTKRASATVKTAVLATLLFATVQANAT 51
 V VSELTR HTKRASATV+TAVLATLLFATVQANAT
 Sbjct: 17 VVSELTRTHTKRASATVETAVLATLLFATVQANAT 52

Score = 161 (71.2 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 32/38 (84%), Positives = 36/38 (94%)

Query: 101 VTLKAGDNLKIKQNTNENTNASSFTYSLKKDLTGLINV 138
 +TLKAGDNLKIKQNT+E+TNASSFTYSLKKDLT L +V
 Sbjct: 103 ITLKAGDNLKIKQNTDESTNASSFTYSLKKDLTDLTSV 140

Score = 110 (48.7 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 21/29 (72%), Positives = 25/29 (86%)

Query: 138 VTEKLSFGANGKKVNIISDTKGLNFAKET 166
 V++KLS G NG KVNI SDTKGLNFAK++
 Sbjct: 1439 VSDKLSLGTNGNKVNITSDTKGLNFAKDS 1467

Score = 85 (37.6 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 18/32 (56%), Positives = 20/32 (62%)

Query: 169 TNGDTTVHLNIGIGSTLTDTLAGSSASHVDAGN 200
 T D +HLNGI STLDTL S A+ GN
 Sbjct: 1469 TGDDANIHLNGIASTLTDTLLNSGATTNLGGN 1500

Score = 92 (40.7 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 16/19 (84%), Positives = 19/19 (100%)

Query: 206 RAASIKDVLNAGWNIKGK 224
 RAAS+KDVLNAGWN++GK
 Sbjct: 1509 RAASVKDVLNAGWNVRGK 1527

Score = 90 (39.8 bits), Expect = 1.5e-116, Sum P(11) = 1.5e-116
 Identities = 17/28 (60%), Positives = 20/28 (71%)

Query: 226 STTGQSENVDFVRTYDTVEFLSADTTTT 253
 S Q EN+DFV TYDTV+F+S D TT
 Sbjct: 1530 SANNQVENIDFVATYDTVDFVSGDKDTT 1557

Based on homology with Hsf, it was predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

ORF40-1 (61kDa) was cloned in pET and pGex vectors and expressed in *E.coli*, as described above. The products of protein expression and purification were analyzed by SDS-PAGE. Figure 1A shows the results of affinity purification of the His-fusion protein, and Figure 1B shows the

results of expression of the GST-fusion in *E.coli*. Purified His-fusion protein was used to immunise mice, whose sera were used for FACS analysis (Figure 1C), a bactericidal assay (Figure 1D), and ELISA (positive result). These experiments confirm that ORF40-1 is a surface-exposed protein, and that it is a useful immunogen.

5 Figure 1E shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF40-1.

Example 2

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 7>

```

10      1  ATGTTACGTT TGACTGCTTT AGCCGTATGC ACCGCCCTCG CTTTGGGCGC
      51  GTGTTGCGCG CAAAATTCCG ACTCTGCCCC ACAAGCCAAA GaACAGGCGG
     101  TTTCCGCGCG ACAAACCGAA GgCGCGTCCG TTACCGTCAA AACCgCGCGC
     151  GGCGACGTTT AAATACCGCA AAACCCCGAA CGCATCGCGG TTTACGATTT
     201  GGGTATGCTC GACACCTTGA GCAAACCTGGG CGTGAAAACC GGTTTGTCGG
     251  TCGATAAAAA CCGCCTGCGG TATTTAGAGG AATATTTCAA AACGACAAAA
     301  CCTGCCGGCA CTTTGTTCTGA GCCGGATTAC GAAACGCTCA ACGCTTACAA
     15  351  ACCGAGCTC ATCATCATCG GCAGCCGCGC CgCCAAGGCG TTTGACAAAT
     401  TGAACGAAAT CGCGCCGACC ATCGTmWTGA CCGCCGATAC CGCCAACCTC
     451  AAAGAAAGTG CCAArGAGGC ATCGACGCTG GCGCAAATCT TC..

```

This corresponds to the amino acid sequence <SEQ ID 8; ORF38>:

```

20      1  MLRLTALAVC TALALGACSP QNSDSAPQAK EQAVSAAQTE GASVTVKTKR
     51  GDVQIPQNPE RIAVYDLGML DTLskLGVKT GLSVDKNRLP YLEEFKTTK
     101  PAGTLFEPDY ETLNAYKPQL IIIGSRAAKA FDKLNEIAPT IXXTADTANL
     151  KESAKEASTL AQIF..

```

Further work revealed the complete nucleotide sequence <SEQ ID 9>:

```

25      1  ATGTTACGTT TGACTGCTTT AGCCGTATGC ACCGCCCTCG CTTTGGGCGC
     51  GTGTTGCGCG CAAAATTCCG ACTCTGCCCC ACAAGCCAAA GAACAGGCGG
     101  TTTCCGCGCG ACAAACCGAA GGCGCGTCCG TTACCGTCAA AACCgCGCGC
     151  GGCGACGTTT AAATACCGCA AAACCCCGAA CGCATCGCGG TTTACGATTT
     201  GGGTATGCTC GACACCTTGA GCAAACCTGGG CGTGAAAACC GGTTTGTCGG
     30  251  TCGATAAAAA CCGCCTGCGG TATTTAGAGG AATATTTCAA AACGACAAAA
     301  CCTGCCGGCA CTTTGTTCTGA GCCGGATTAC GAAACGCTCA ACGCTTACAA
     351  ACCGAGCTC ATCATCATCG GCAGCCGCGC CGCCAAGGCG TTTGACAAAT
     401  TGAACGAAAT CGCGCCGACC ATCGAAATGA CGCCGATAC CGCCAACCTC
     451  AAAGAAAGTG CCAAGAGCG CATCGACGCG CTGGCGCAAA TCTTCGCGCA
     501  ACAGGCGGAA GCCGACAAGC TGAAGGCGGA AATCGACGCG TCTTTTGAAG
     35  551  CCGCGAAAAC TGCCGACAAA GGTAAGGGCA AAGGTTTGGT GATTTTGCTC
     601  AACGGCGGCA AGATGTCGGC TTTCGGCCCG TCTTCACGCT TGGGCGGCTG
     651  GCTGCACAAA GACATCGGCG TTCCCGCTGT CGATGAATCA ATTAAGAAG
     701  GCAGCCACGG TCAGCCTATC AGCTTTGAAT ACCTGAAAGA GAAAAATCCC
     751  GACTGGCTGT TTGTCCTTGA CCGAAGCGCG GCCATCGCGG AAGAGGGTCA
     40  801  GGCGCGGAAA GACGTGTTGG ATAATCCGCT GGTGCGCGAA ACAACCGCTT
     851  GGAAAAAAGG ACAGGTCGTG TACCTCGTTC CTGAAACTTA TTTGGCAGCC
     901  GGTGGCGCGC AAGAGCTGCT GAATGCAAGC AAACAGGTTG CCGACGCTTT
     951  TAACGCGGCA AAATAA

```

This corresponds to the amino acid sequence <SEQ ID 10; ORF38-1>:

```

45      1  MLRLTALAVC TALALGACSP QNSDSAPQAK EQAVSAAQTE GASVTVKTKR
     51  GDVQIPQNPE RIAVYDLGML DTLskLGVKT GLSVDKNRLP YLEEFKTTK
     101  PAGTLFEPDY ETLNAYKPQL IIIGSRAAKA FDKLNEIAPT IEMTADTANL
     151  KESAKERIDA LAQIFGKQAE ADKLKAEIDA SFEEAKTAAQ GKKGGLVILV
     201  NGGKMSAFGP SSRLGGWLHK DIGVPAVDES IKGSHGQPI SFEYLKEKNP
     50  251  DWLFVLDRSA AIGEEGQAAK DVLNPLVAE TTAWKKGQVV YLVPETYLAA

```

301 GGAQELLNAS KQVADAFNAA K*

Computer analysis of this amino acid sequence reveals a putative prokaryotic membrane lipoprotein lipid attachment site (underlined).

Further work identified the corresponding gene in strain A of *N.meningitidis* <SEQ ID 11>:

```

5      1  ATGTTACGTT  TGA CTGCTTT  AGCCGTATGC  ACCGCCCTCG  CTTTGGGCGC
      51  GTGTTGCGCG  CAAAATTCCG  ACTCTGCCCC  ACAAGCCAAA  GAACAGGCGG
     101  TTTCCGCGCG  ACAATCCGAA  GGC GTGTCG  TTACCGTCAA  AACGGCGCGC
     151  GGCGATGTTT  AAATACCGCA  AAACCCCGAA  CGTATCGCCG  TTTACGATTT
     201  GGGTATGCTC  GACACCTTGA  GCAAACTGGG  CGTGAAAACC  GGTTTGTCCG
    10  251  TCGATAAAAA  CCGCCTGCCG  TATTTAGAGG  AATATTTCAA  AACGACAAAA
     301  CCTGCCGGAA  CTTTGTTCGA  GCCGGATTAC  GAAACGCTCA  ACGCTTACAA
     351  ACCGCAGCTC  ATCATCATCG  GCAGCCGCGC  AGCCAAAGCG  TTTGACAAAT
     401  TGAACGAAAT  CGCGCCGACC  ATCGAAATGA  CCGCCGATAC  CGCCAACCTC
     451  AAAGAAAGTG  CCAAAGAGCG  TATCGACGCG  CTGGCGCAAA  TCTTCGCAAA
    15  501  AAAGGCGGAA  GCCGACAAGC  TGAAGCGGGA  AATCGACGCG  TCTTTTGAAG
     551  CCGCGAAAAC  TGCCGCGCAA  GGCAAAGGCA  AGGGTTTGGT  GATTTTGGTC
     601  AACGGCGGCA  AGATGTCCCG  CTTCCGCCCG  TCTTCACGAC  TGGGCGGCTG
     651  GCTGCACAAA  GACATCGGCG  TTCCCGCTGT  TGACGAAGCC  ATCAAAGAAG
     701  GCAGCCACGG  TCAGCCTATC  AGCTTTGAAT  ACCTGAAAGA  GAAAAATCCC
    20  751  GACTGGCTGT  TTGTCCTTGA  CCGCAGCGCG  GCCATCGGCG  AAGAGGGTCA
     801  GGCGGCGAAA  GACGTGTTGA  ACAATCCGCT  GGTGCGCGAA  ACAACCGCTT
     851  GGAAAAAAGG  ACAAGTCGTT  TACCTTGTTT  CTGAAACTTA  TTTGGCAGCC
     901  GGTGGCGCGC  AAGAGCTACT  GAATGCAAGC  AAACAGGTTG  CCGACGCTTT
     951  TAACGCGGCA  AAATAA
  
```

25 This encodes a protein having amino acid sequence <SEQ ID 12; ORF38a>:

```

      1  MLRLTALAVC  TALALGACSP  QNSDSAPQAK  EQAVSAAQSE  GVS VTKTAR
     51  GDVQIPQNPE  RIAVYDLGML  DTL SKLGVT  GLSVDKNRLP  YLEEFKTTK
    101  PAGTLFEPDY  ETLNAYKPQL  IIIGSRAAKA  FDKLNEIAPT  IEMTADTANL
    151  KESAKERIDA  LAQIFGKKA  ADKLKAEIDA  SFEAAKTAQ  KGKGLVILV
    201  NGGKMSAFGP  SSRLGGWLHK  DIGVPAVDEA  IKEGSHGQPI  SFEYLKEKNP
    251  DWLFVLDRSA  AIGEEGQAAK  DVLNNPLVAE  TTAWKKQV  YLVPETYLAA
    301  GGAQELLNAS  KQVADAFNAA  K*
  
```

The originally-identified partial strain B sequence (ORF38) shows 95.2% identity over a 165aa overlap with ORF38a:

```

35      10      20      30      40      50      60
    orf38.pep  MLRLTALAVCTALALGACSPQNSDSAPQAKEQAVSAAQTEGASVTVKTARGDVQIPQNPE
    orf38a      MLRLTALAVCTALALGACSPQNSDSAPQAKEQAVSAAQSEGVSVTVKTARGDVQIPQNPE
      10      20      30      40      50      60
    40      70      80      90     100     110     120
    orf38.pep  RIAVYDLGMLDTLSKLGVTGLSVDKNRLPYLEEFKTTKPAGTLFEPDYETLNAYKPQL
    orf38a      RIAVYDLGMLDTLSKLGVTGLSVDKNRLPYLEEFKTTKPAGTLFEPDYETLNAYKPQL
      70      80      90     100     110     120
    45
      130     140     150     160
    orf38.pep  IIIGSRAAKAFDKLNEIAPTIXXTADTANLKESAKE-ASTLAQIF
    orf38a      IIIGSRAAKAFDKLNEIAPTIEMTADTANLKESAKERIDALAQIFGKKAADKLKAEIDA
      130     140     150     160     170     180
    50
    orf38a      SFEAAKTAQGGKGLVILVNGGKMSAFGPSSRLGGWLHKDIGVPAVDEAIKEGSHGQPI
      190     200     210     220     230     240
  
```

55 The complete strain B sequence (ORF38-1) and ORF38a show 98.4% identity in 321 aa overlap:

```

    orf38a.pep    MLRLTALAVCTALALGACSPQNSDSAPQAKEQAVSAAQSEGVSVTVKTTARGDVQIPQNPE
    orf38-1       MLRLTALAVCTALALGACSPQNSDSAPQAKEQAVSAAQTEGASVTVKTTARGDVQIPQNPE
5   orf38a.pep    RIAVYDLGMLDTLSKLGVKTTGLSVVDKNRLPYLEEFKTTKPGATLFEFDPYETLNAYKPQL
    orf38-1       RIAVYDLGMLDTLSKLGVKTTGLSVVDKNRLPYLEEFKTTKPGATLFEFDPYETLNAYKPQL
10  orf38a.pep    IIIGSRAAKAFDKLNEIAPTIEMTADTANLKESAKERIDALAQIFGKKAADKLKAEIDA
    orf38-1       IIIGSRAAKAFDKLNEIAPTIEMTADTANLKESAKERIDALAQIFGKQAEADKLKAEIDA
15  orf38a.pep    SFEAAKTAAQKGKGLVILVNGGKMSAFGPSSRLGGWLHKDIGVPAVDEAIKEGSHGQPI
    orf38-1       SFEAAKTAAQKGKGLVILVNGGKMSAFGPSSRLGGWLHKDIGVPAVDESIKEGSHGQPI
20  orf38a.pep    SFEYLKEKNPDWLFVLDRAAIGEEGQAAKDVLDNPLVAETTAWKKGQVVYLPETYLA
    orf38-1       SFEYLKEKNPDWLFVLDRAAIGEEGQAAKDVLDNPLVAETTAWKKGQVVYLPETYLA
30  orf38a.pep    GGAQELLNASKQVADAFNA
    orf38-1       GGAQELLNASKQVADAFNA

```

Computer analysis of these sequences revealed the following:

25 Homology with a lipoprotein (lipo) of *C.jejuni* (accession number X82427)

ORF38 and lipo show 38% aa identity in 96 aa overlap:

```

    Orf38: 40  EGASVTVKTTARGDVQIPQNPERIAVYDLGMLDTLSKLGVKTTGLS-VDKNRLPYLEEFKT 98
                EG S  VK + G + + P+NP ++ + DLG+LDT  L +  ++ V  LP  + FK
30  Lipo:  51  EGDSEFLVKDSLGENKTPKNPSKVILDLGILDTFDALKLNDKVAGVPAKNLPKYLQQFEKN 110
    Orf38: 99  TKPAGTLFEFDPYETLNAYKPQLIIIGSRAAKAFDKL 134
                G + + D+E +NA KP LIII  R +K +DKL
    Lipo:  111 KPSVGGVQQVDFAINALKPDLLIIISGRQSKFYDKL 146

```

Based on this analysis, it was predicted that this protein from *N.meningitidis*, and its epitopes, could
 35 be useful antigens for vaccines or diagnostics.

ORF38-1 (32kDa) was cloned in pET and pGex vectors and expressed in *E.coli*, as described above. The products of protein expression and purification were analyzed by SDS-PAGE. Figure 2A shows the results of affinity purification of the His-fusion protein, and Figure 2B shows the results of expression of the GST-fusion in *E.coli*. Purified His-fusion protein was used to immunise
 40 mice, whose sera were used for Western blot analysis (Figure 2C) and FACS analysis (Figure 2D). These experiments confirm that ORF38-1 is a surface-exposed protein, and that it is a useful immunogen.

Figure 2E shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF38-1.

Example 3

45 The following *N.meningitidis* DNA sequence was identified <SEQ ID 13>:

1 ATGAAACTTC TGACCACCGC AATCCTGTCT TCCGCAATCG CGCTCAGCAG
 51 TATGGCTGCC GCCGTGGCA CGGACAACCC CACTGTTGCA AAAAAAACCG
 101 TCAGCTACGT CTGCCAGCAA GGTAAAAAAG TCAAAGTAAC CTACGGCTTC
 5 151 AACAAACAGG GTCTGACCAC ATACGCTTCC GCCGTCATCA ACGGCAAACG
 201 CGTGCAAATG CCTGTCAATT TGGACAAATC CGACAATGTG GAAACATTCT
 251 ACGGCAAAGA AGGCGGTTAT GTTTTGGGTA CCGGCGTGAT GGATGGCAAA
 301 TCCTACCGCA AACAGCCCAT TATGATTACC GCACCTGACA ACCAAATCGT
 351 CTTCAAAGAC TGTCCCCAC GTTAA

This corresponds to the amino acid sequence <SEQ ID 14; ORF44>:

10 1 MKLLTTAILS SAIALSSMAA AAGTDNPTVA KKTVSYVCQQ GKVKVITYGF
 51 NKQGLTTYAS AVINGKRVQM PVNLDKSDNV ETFYGKEGGY VLGTGVMDGK
 101 SYRKQPIMIT APDNQIVFKD CSPR*

Computer analysis of this amino acid sequence predicted the leader peptide shown underlined.

Further work identified the corresponding gene in strain A of *N.meningitidis* <SEQ ID 15>:

15 1 ATGAAACTTC TGACCACCGC AATCCTGTCT TCCGCAATCG CGCTCAGCAG
 51 TATGGCTGCT GCTGCCGGCA CGAACAACCC CACCGTTGCC AAAAAAACCG
 101 TCAGCTACGT CTGCCAGCAA GGTAAAAAAG TCAAAGTAAC CTACGGCTTT
 151 AACAAACAGG GCCTGACCAC ATACGCTTCC GCCGTCATCA ACGGCAAACG
 20 201 TGTGCAAATG CCTGTCAATT TGGACAAATC CGACAATGTG GAAACATTCT
 251 ACGGCAAAGA AGGCGGTTAT GTTTTGGGTA CCGGCGTGAT GGATGGCAAA
 301 TCCTATCGCA AACAGCCTAT TATGATTACC GCACCTGACA ACCAAATCGT
 351 CTTCAAAGAC TGTCCCCAC GTTAA

This encodes a protein having amino acid sequence <SEQ ID 16; ORF44a>:

25 1 MKLLTTAILS SAIALSSMAA AAGTNNPTVA KKTVSYVCQQ GKVKVITYGF
 51 NKQGLTTYAS AVINGKRVQM PVNLDKSDNV ETFYGKEGGY VLGTGVMDGK
 101 SYRKQPIMIT APDNQIVFKD CSPR*

The strain B sequence (ORF44) shows 99.2% identity over a 124aa overlap with ORF44a:

		10	20	30	40	50	60
30	orf44.pep	<u>MKLLTTAILSSAIALSSMAAAAGTDNPTVA</u> KKTVSYVCQQGKKVKVITYGFNKQGLTTYAS					
	orf44a	MKLLTTAILSSAIALSSMAAAAGTNNPTVAKKTVSYVCQQGKKVKVITYGFNKQGLTTYAS					
		10	20	30	40	50	60
35	orf44.pep	AVINGKRVQMPVNLDKSDNVETFYGKEGGYVLGTGVMDGKSYRKQPIMITAPDNQIVFKD					
	orf44a	AVINGKRVQMPVNLDKSDNVETFYGKEGGYVLGTGVMDGKSYRKQPIMITAPDNQIVFKD					
		70	80	90	100	110	120
40	orf44.pep	CSPRX					
	orf44a	CSPRX					

Computer analysis gave the following results:

Homology with the LecA adhesin of *Eikenella corrodens* (accession number D78153)

45 ORF44 and LecA protein show 45% aa identity in 91 aa overlap:

Orf44 33 TVSYVCQQGKKVKVITYGFNKQGLTTYASAVINGKRVQMPVNLDKSDNVETFYGKEGGYVL 92
 +V+YVCQQG+++ V Y FN G+ T A +N + +++P NL SDNV+T + GY L
 LecA 135 SVAYVCQQRRLNVNRYRFSAGVPTSaelrvnnrnlrlpynlsasdnvdtvf-SANGYRL 193
 50 Orf44 93 GTGVMDGKSYRKQPIMITAPDNQIVFKDCSP 123
 T MD +YR Q I+++AP+ Q+++KDCSP

LecA 194 TTNAMDSANYRSQDIIVSAPNGQMLYKDCSP 224

Based on homology with the adhesin, it was predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

ORF44-1 (11.2kDa) was cloned in pET and pGex vectors and expressed in *E.coli*, as described above. The products of protein expression and purification were analyzed by SDS-PAGE. Figure 3A shows the results of affinity purification of the His-fusion protein, and Figure 3B shows the results of expression of the GST-fusion in *E.coli*. Purified His-fusion protein was used to immunise mice, whose sera were used for ELISA, which gave positive results, and for a bactericidal assay (Figure 3C). These experiments confirm that ORF44-1 is a surface-exposed protein, and that it is a useful immunogen.

Figure 3D shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF44-1.

Example 4

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 17>

```

15      1  ..GGCACC GAAT TCAAAACCAC CCTTTC CGGA GCCGACATAC AGGCAGGGGT
      51  GGGTGAAAAA GCCCGAGCCG ATGCGAAAAT TATCCTAAAA GGCATCGTTA
     101  ACCGCATCCA AACCGAAGAA AAGCTGGAAT CCAACTCGAC CGTATGGCAA
     151  AAGCAGGCCG GAAGCGGCAG CACGGTTGAA ACGCTGAAGC TACCGAGCTT
     201  TGAAGGGCCG GCACTGCCTA AGCTGACCGC TCCCGGCCGC TATATCGCCG
     251  ACATCCCCAA AGGCAACCTC AAAACCGAAA TCGAAAAGCT GGCCAAACAG
     301  CCCGAATATG CCTATCTGAA ACAGCTTCAG ACGGTCAAGG ACGTGAAGTG
     351  GAACCAAGTA CAGCTCGCTT ACGACAAATG GGAATAATAA CAGGAAGGCC
     401  TAACCGGAGC CGGAGCCGCA ATTANCGCAC TGGCCGTAC CGTGGTCACC
     451  TCAGGCGCAG GAACCGGAGC CGTATTGGGA TTAANACGNG TGGCCGCCGC
     501  CGCAACCGAT GCAGCATTT...
```

25 This corresponds to the amino acid sequence <SEQ ID 18; ORF49>:

```

      1  ..GTEFKTTL SG ADIQAGVGEK ARADAKIILK GIVNRIQTEE KLESNSTVWQ
     51  KQAGSGSTVE TLKLPSFEGP ALPKLTAPGG YIADIPKGNL KTEIEKLAKQ
    101  PEYAYLKQLQ TVKDVNWNQV QLAYDKWDYK QEGLTGAGAA IXALAVTVVT
    151  SGAGTGAVLG LXRVAATAAD AAF..
```

30 Further work revealed the complete nucleotide sequence <SEQ ID 19>:

```

      1  ATGCAACTGC TGGCAGCCGA AGGCATTAC CAACACCAAT TGAATGTTCA
     51  GAAAAGTACC CGTTTCATCG GCATCAAAGT GGGTAAAAGC AATTACAGCA
    101  AAAACGAGCT GAACGAAACC AAACGTGCCG TACGCGTTAT CGCCCAAACA
    151  GCCAAAACCC GTTCCGGCTG GGATACCGTA CTCGAAGGCA CGAATTCAA
    201  AACCACCCTT TCCGGAGCCG ACATACAGGC AGGGGTGGGT GAAAAGCCC
    251  GAGCCGATGC GAAAATTATC CTAAAAGGCA TCGTTAACCG CATCCAAACC
    301  GAAGAAAAGC TGGAAATCAA CTCGACCGTA TGGCAAAAGC AGGCCGGAAG
    351  CGGCAGCAGC GTTGAAACGC TGAAGCTACC GAGCTTTGAA GGGCCGGCAC
    401  TGCCTAAGCT GACCGCTCCC GCGGGCTATA TCGCCGACAT CCCCAGGCG
    451  AACCTCAAAA CCGAAATCGA AAAGCTGGCC AAACAGCCCG AATATGCCTA
    501  TCTGAAACAG CTTAGACGGT TCAAGGACGT GAACTGGAAC CAAGTACAGC
    551  TCGCTTACGA CAAATGGGAC TATAACAGG AAGGCCTAAC CGGAGCGGGA
    601  GCCGCAATTA TCGCACTGGC CGTTACCGTG GTCACCTCAG GCGCAGGAAC
    651  CGGAGCCGTA TTGGGATTAA ACGGTGCGGC CGCCGCCGCA ACCGATGCAG
```

5
10
15
20
25

```

701 CATTTGCCTC TTTGGCCAGC CAGGCTTCCG TATCGTTCAT CAACAACAAA
751 GGCAATATCG GTAACACCCT GAAAGAGCTG GGCAGAAGCA GCACGGTGAA
801 AAATCTGATG GTTGCCGTCG CTACCGCAGG CGTAGCCGAC AAAATCGGTG
851 CTTCCGGCACT GAACAATGTC AGCGATAAGC AGTGGATCAA CAACCTGACC
901 GTCAACCTGG CCAATGCGGG CAGTGCCGCA CTGATTAATA CCGCTGTCAA
951 CGGCGGCAGC CTGAAAGACA ATCTGGAAGC GAATATCCTT GCGGCTTTGG
1001 TGAATACTGC GCATGGAGAG GCAGCAAGTA AAATCAAACA GTTGGATCAG
1051 CACTACATTG CCCATAAGAT TGCCCATGCC ATAGCGGGCT GTGCGGCAGC
1101 GGCGGCGAAT AAGGGCAAGT GTCAGATGG TGCGATCGGT GCGGCGGTCTG
1151 GTGAAATCCT TGGCGAAACC CTACTGGACG GCAGAGACCC TGGCAGCCTG
1201 AATGTGAAGG ACAGGGCAAA AATCATTGCT AAGGCGAAGC TGGCAGCAGG
1251 GGCGGTTGCG GCGTTGAGTA AGGGGGATGT GAGTACGGCG GCGAATGCGG
1301 CTGCTGTGGC GGTAGAGAAT AATTCTTTAA ATGATATACA GGATCGTTTG
1351 TTGAGTGGAA ATTATGCTTT ATGTATGAGT GCAGGAGGAG CAGAAAGCTT
1401 TTGTGGTTT AAATGGGAC TGGGCTTGCC ACACTTTGTA AGTGTTTCAG
1451 GAGAAATGAA ATTACCTAAT AAATTCGGGA ATCGTATGGT TAATGGAAAA
1501 TTAATTATTA AACTAGAAA TGGCAATGTA TATTCTCTCT TAGGTAAAT
1551 ATGGAGTACT GTAAAATCAA CAAAATCAAA TATAAGTGGG GTATCTGTCTG
1601 GNTGGGTTT AAATGTTTCC CCTAATGATT ATTTAAAAGA AGCATCTATG
1651 AATGATTTC AAAATAGTAA TCAAAATAAA GCCTATGCAG AAATGATTTT
1701 CCAGACTTTG GTAGGTGAGA GTGTTGGTGG TAGTCTTTGT CTGACAAGAG
1751 CCTGCTTTTC GGTAAGTTCA ACAATATCTA AATCTAAATC TCCTTTTAAA
1801 GATTCAAAAA TTATTGGGGA AATCGGTTTG GGAAGTGGTG TTGCTGCAGG
1851 AGTAGAAAAA ACAATATACA TAGGTAACAT AAAAGATATT GATAAATTTA
1901 TTAGTGCAAA CATAAAAAA TAG

```

This corresponds to the amino acid sequence <SEQ ID 20; ORF49-1>:

30
35
40

```

1  MQLLAAEGIH QHQLNVQKST RFIGIKVGKS NYSKNELNET KLPVRVIAQT
51 AKTRSGWDTV LEGTEFKTTL SGADIQAGVG EKARADAKII LKGIVNRIQT
101 EEKLESNSTV WQKQAGSGST VETLKLPSFE GPALPKLTAP GGYIADIPKG
151 NLKTEIEKLA KOPEYAYLKQ LQTVKDVNWN QVQLAYDKWD YKQEGLTGAG
201 AAIIALAVTV VTSGAGTGAV LGLNGAAAAA TDAAFASLAS QASVSFINNK
251 GNIGNTLKEL GRSSTVKNLM VAVATAGVAD KIGASALNNV SDKQWINNLT
301 VNLANAGSAA LINTAVNGGS LKDNLEANIL AALVNTAHGE AASKIKQLDQ
351 HYIAHKIAHA IAGCAAAAAN KGKQDGAIG AAVGEILGET LLDGRDPGSL
401 NVKDRAKIIA KAKLAAGAVA ALSKGDVSTA ANAAAVAVEN NSLNDIQDRL
451 LSGNYALCMS AGGAESFCES YRPLGLPHFV SVSGEMKLPN KFGNRMVNGK
501 LIINTRNGNV YFSVGKIWST VKSTKSNISG VSVGWVLNVS PNDYLKEASM
551 NDFRNSNQNK AYAEMISQTL VGESVGGSLC LTRACFSVSS TISKSKSPFK
601 DSKIIGEIGL GSGVAAGVEK TIYIGNIKDI DKFISANIKK *

```

Computer analysis predicts a transmembrane domain and also indicates that ORF49 has no significant amino acid homology with known proteins. A corresponding ORF from *N.meningitidis* strain A was, however, identified:

ORF49 shows 86.1% identity over a 173aa overlap with an ORF (ORF49a) from strain A of *N.meningitidis*:

45
50
55

```

orf49.pep          10      20      30
                   GTEFKTTL SGADIQAGVGEKARADAKIILK
                   |||||:||||| |||:|||||
orf49a             SKNELNETKLPVRVVAQXAATRSGWDTVLEGTEFKTTLGADIQAGVXEKARVDAKIILK
                   40      50      60      70      80      90

orf49.pep          40      50      60      70      80      90
                   GIVNRIQTEEKLESNSTVWQKQAGSGSTVETLKLPSFEGPALPKLTAPGGYIADIPKGNL
                   |||||:||||| |||||:||||| |||:||||| |||:|||||
orf49a             GIVNRIQSEKLETNSTVWQKQAGRGSTIETLKLPSFESPTPPKLSAPGGYIVDIPKGNL
                   100     110     120     130     140     150

orf49.pep          100     110     120     130     140     150
                   KTEIEKLAKQPEYAYLKQLQTVKDVNWNQVQLAYDKWDYKQEGLTGAGAAIXALAVTVVT
                   |||||:||||| |||:||||| |||:||||| |||:|||||

```

```

orf49a      KTEIEKLSKQPEYAYLKLQVAKNINWNQVQLAYDRWDYKQEGLTEAGAAIIALAVTVVT
              160      170      180      190      200      210

              160      170
orf49.pep   SGAGTGAVLGLXRVAAAATDAAF
              ||||| : |||||
orf49a      SGAGTGAVLGLNGAXAAATDAAFASLASQASVSFINNKGDVGKTLKELGRSSTVKNLVA
              220      230      240      250      260      270

```

ORF49-1 and ORF49a show 83.2% identity in 457 aa overlap:

10	orf49a.pep	XQLLAAEGIGHKHELDVQKSRRFIGIKVGXSNYSKNELNKLPVRVVAQXAATRSQWDTV : : : : :
	orf49-1	MQLLAAEGIHQHQLNVQKSTRFIGIKVGKSNYSKNELNKLPVRVIAQTAKTRSGWDTV
15	orf49a.pep	LEGTEFKTTLGADIQAGVXEKARVDAKIILKGIVNRIQSEEKLETNSTVWQKQAGRGST : : : :
	orf49-1	LEGTEFKTTLGADIQAGVGEKARADAKIILKGIVNRIQTEEKLESNSTVWQKQAGSGST
20	orf49a.pep	IETLKLPSFESPTPPKLSAPGGYIVDIPKGNLKTEIEKLSKQPEYAYLKQLQVAKNINWN : : : : : : : :
	orf49-1	VETLKLPSFEGPALPKLTAPGGYIADIPKGNLKTEIEKLAQPEYAYLKQLQTVKDVNWN
25	orf49a.pep	QVQLAYDRWDYKQEGLTEAGAAI IALAVTVVTSAGTGAVLGLNGAXAAATDAAFASLAS :
	orf49-1	QVQLAYDKWDYKQEGLTGAGAAI IALAVTVVTSAGTGAVLGLNGAAAAATDAAFASLAS
30	orf49a.pep	QASVSFINNKGDVGKTLKELGRSSTVKNLVVAAATAGVADKIGASALXNVSDKQWINNLT : : : :
	orf49-1	QASVSFINNKGNI NT LKELGRSSTVKNLMVAVATAGVADKIGASALNNVSDKQWINNLT
35	orf49a.pep	VNLANAGS AALINTAVNGGSLKDXLEANILAALVNTAHGEAASKIKQLDQHYIVHKIAHA
	orf49-1	VNLANAGS AALINTAVNGGSLKDNLEANILAALVNTAHGEAASKIKQLDQHYIAHKIAHA
40	orf49a.pep	IAGCAAAAANKGKCQDGAIGA AVGEIVGEALTNGKNPDTLTAKEREQILAYSKLVAGTVS : : : : : : :
	orf49-1	IAGCAAAAANKGKCQDGAIGA AVGEILGETLLDGRDPGSLNVKDRAKI IAKAKLAAGAVA
45	orf49a.pep	GVVGGDVNAAANAAE VAVKNNQLSDXEGREFDNEMTACAKQNX PQLCRKNTVKKYQNVAD : : : : : : : : : : : : :
	orf49-1	ALSKGDVSTAANAAAVAVENNSLNDIQDRLLSGNYALCMSAGGAESFCESYRPLGLPHFV
50	orf49a.pep	KRLAASIAICTDISRSTECRTIRKQHLIDSRSLHSSWEAGLIGKDDWEYKLFSSKSYTQAD
	orf49-1	SVSGEMKLPNKFGRNMVNGKLIINTRNGNVYFSVGKIWSTVKSTKSNISGVSVGWVLNVS

45 The complete length ORF49a nucleotide sequence <SEQ ID 21> is:

	1	NTGCAACTGC	TGGCAGAAGA	AGGCATCCAC	AAGCACGAGT	TGGATGTCCA
	51	AAAAAGCCGC	CGCTTTATCG	GCATCAAGGT	AGGTNAGAGC	AATTACAGTA
	101	AAAACGAATC	GAAACGAAACC	AAATTGCCTG	TCCGCGTCGT	CGCCCAAANT
50	151	GCAGCAACCC	GTTCAGGCTG	GGTACCGCTG	CTCGAAGGTA	CCGAATTCAA
	201	AACCACGCTG	GCCGGTGCCG	ACATTGAGGC	AGGTGTANGC	GAAAAGAGCCC
	251	GTGTCGATGC	GAAAATTATC	CTCAAAGGCA	TTGTGAACCG	TATCCAGTCG
	301	GAAGAAAAAT	TAGAAACCAA	CTCAACCGTA	TGGCAGAAAC	AGGCCGGACG
	351	CGGCAGCACT	ATCGAAACGC	TAAAACCTGCC	CAGCTTCGAA	AGCCCTACTC
55	401	CGCCCAAATT	GTCCGCACCC	GGCGGNTATA	TCGTTCACAT	TCCGAAAGGC
	451	AATCTGAAAA	CCGAAATCGA	AAAGCTGTCC	AAACAGCCCG	AGTATGCCTA
	501	CTGAAACAG	CTCCAAGTAG	CGAAAAACAT	CAACTGGAAT	CAGGTGCAGC
	551	TTGCTTACGA	CAGATGGGAC	TACAAACAGG	AGGGCTTAAC	CGAAGCAGGT
	601	GCGGCGATTA	TCGCACCTGCG	CGTTACCGTG	GTCACTTCAG	GCGCAGGAAC
60	651	CGGAGCCGTA	TTGGGATTAA	ACGGTGCAGC	CGCCGCCGCA	ACCGATGCAG
	701	CATTGCGCTC	TTTGGCCAGC	CAGGCTTCGG	TATCGTTTCAT	CAACAACAAA
	751	GGCGATGTG	GCAAAACCC	GAAAGAGCTG	GGCAGAAGCA	GCACGGTGAA
	801	AAATCTGGTG	GTTGCCGCGC	CTACCGCAGG	CGTAGCCGAC	AAAAATCGGC
	851	CTTCGGCACT	GANCAATGTC	AGCGATAAGC	AGTGGATCAA	CAACCTGACC
	901	GTCACCTAG	CCAATGCGGG	CAGTGCOCGA	CTGATTAATC	CCGCTGTCAA
65	951	CGGCCGCGAG	CTGAAAGACA	NTCTGGAAGC	GAATATCCTT	GCGGCTTTGG
	1001	TCAATACCGC	GCATGGAGAA	GCAGCCAGTA	AAATCAAACA	GTTGGATCAG

```

1051  CACTACATAG TCCACAAGAT TGCCCATGCC ATAGCGGGCT GTGCGGCAGC
1101  GCGGCGCAAT AAGGGCAAGT GTCAGGATGG TCGGATAGGT GCGGCTGTGG
1151  GCGAGATAGT CGGGGAGGCT TTGACAAACG GCAAAAATCC TGACACTTTG
5    1201  ACAGCTAAAG AACGCGAACA GATTTTGGCA TACAGCAAAC TGGTTGCCGG
1251  TACGGTAAGC GGTGTGGTCG GCGGCGATGT AAATGCGGCG GCGAATGCGG
1301  CTGAGGTAGC GGTGAAAAAT AATCAGCTTA GCGACNAAGA GGGTAGAGAA
1351  TTTGATAACG AAATGACTGC ATGCGCCAAA CAGAATANTC CTCAACTGTG
1401  CAGAAAAAAT ACTGTAAAAA AGTATCAAAA TGTGTCTGAT AAAAGACTTG
1451  CTGCTTCGAT TGCAATATGT ACGGATATAT CCCGTAGTAC TGAATGTAGA
10   1501  ACAATCAGAA AACAACATTG GATCGATAGT AGAAGCCTTC ATTCATCTTG
1551  GGAAGCAGGT CTAATTGGTA AAGATGATGA ATGGTATAAA TTATTCAGCA
1601  AATCTTACAC CCAAGCAGAT TTGGCTTTAC AGTCTTATCA TTTGAATACT
1651  GCTGCTAAAT CTTGGCTTCA ATCGGGCAAT ACAAAGCCTT TATCCGAATG
1701  GATGTCCGAC CAAGGTTATA CACTTATTTT AGGAGTTAAT CCTAGATTCA
15   1751  TTCCAATACC AAGAGGTTT GTAAACAAA ATACACCTAT TACTAATGTA
1801  AAATACCCGG AAGGCATCAG TTTCGATACA AACCTANAAA GACATCTGGC
1851  AAATGCTGAT GGTTTTAGTC AAGAACAGGG CATTAAGGA GCCCATAACC
1901  GCACCAATNT TATGGCAGAA CTAAATTCAC GAGGAGGANG NGTAAATCT
1951  GAAACCCANA TAGTATTGA AGGCATTACC CGAATTAAAT ATGAGTCTCC
20   2001  TACACTAGAC AGGACAGGTA AACCTGATGG TGGATTTAAG GAAATTTCAA
2051  GTATAAAAAC TGTTTATAAT CCTAAAAANT TTNNGATGA TAAATACTT
2101  CAAATGGCTC AANATGCTGN TTCACAAGGA TATTCAAAG CCTCTAAAT
2151  TGCTCAAAAT GAAAGAACTA AATCAATATC GGAAAGAAAA AATGTCATTC
2201  AATTCTCAGA AACCTTTGAC GGAATCAAAT TTAGANNNTA TNTNGATGTA
25   2251  AATACAGGAA GAATTACAAA CATTACCCCA GAATAATTTA A

```

This encodes a protein having amino acid sequence <SEQ ID 22>:

```

1    XQLLAEEGIH KHELDVQKSR RFIGIKVGXS NYSKNELNET KLPVRVVAQX
51   AATRSWWDTV LEGTEFKTTL AGADIQAGVX EKARVDAKII LKGIVNRIQS
101  EEKLETNSTV WQKQAGRST IETLKLPSFE SPTPKLSAP GGYIVDIPKG
30   151  NLKTEIEKLS KOPEYAYLKQ LQVAKNINWN QVQLAYDRWD YKQEGLTEAG
201  AIIIALAVTV VTSGAGTGAV LGLNGAXAAA TDAAFASLAS QASVSFINNK
251  GDVGKTLKEL GRSSTVKNLV VAAATAGVAD KIGASALXNV SDKOWINNL
301  VNLNAGSAA LINTAVNGGS LKDXLEANIL AALVNTAHGE AASKIKQLDQ
35   351  HYIVHKIAHA IAGCAAAAN KGKQDGAIG AAVGEIVGEA LTNGKNPDTL
401  TAKEREQILA YSKLVAGTVS GVVGGDVNAA ANAAEVAVKN NQLSDXEGRE
451  FDNEMTCAK QNXPQLCRKN TVKKYQNVAD KRLAASIAIC TDISRSTECR
501  TIRKQHLIDS RSLHSSWEAG LIGKDDEWYK LFSKSYTQAD LALQSYHLNT
551  AAKSWLQSGN TKPLSEWMSD QGYTLISGVN PRFIPIPRGF VKONTPIITNV
601  KYPEGISFDT NLXRHLANAD GFSQEQGIK AHNRNTNXMAE LNSRGGXVKS
40   651  ETXTDIEGIT RIKYEIPTLD RTGKPDGGFK EISSIKTVYN PKXFXDDKIL
701  QMAQXAXSQG YSKASKIAQN ERTKSISERK NVIQFSETFD GIKFRXYXDV
751  NTGRITNIHP E*

```

Based on the presence of a putative transmembrane domain, it is predicted that these proteins from *N.meningitidis*, and their epitopes, could be useful antigens for vaccines or diagnostics.

45 Example 5

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 23>

```

1    ..CGGATCGTTG TAGGTTTGCG GATTTCTTGC GCCGTAGTCA CCGTAGTCCC
51   AAGTATAACC CAAGGCTTTG TCTTCGCCTT TCATTCCGAT AAGGGATATG
101  ACGCTTGGT CCGTATAGCC GTCTTGGGAA CCTTTGTCCA CCCAACGCAT
50   151  ATCTGCCTGC GGATTCTCAT TGCCGCTTCT TGGCTGCTGA TTTTCTGCC
201  TTCGCGTTTT TCAACTTCGC GCTTGAGGGC TTCGGCATAT TTGTCGGCCA
251  ACGCCATTTC TTTCGATGAC AGCTGCCTAT TGTTCCAATC TACATTGCGA
301  CCCACCACAG CACCACCACT ACCACCAGTT GCATAG

```

This corresponds to the amino acid sequence <SEQ ID 24; ORF50>:

```

55   1    ..RIVVGLRISC AVVTVVPSIT QGFVFAFHS D KGYDALVGIA VLGTFVHPTH
51   ICLRILIAAS WLLIFLPSRF STSRLRASAY LSANAISFGC SCLLFQSTFA
101  PTTAPPLPPV A*

```

Computer analysis predicts two transmembrane domains and also indicates that ORF50 has no significant amino acid homology with known proteins.

Based on the presence of a putative transmembrane domain, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

5 Example 6

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 25>

```

1    ..AAGTTTGA CT TACCTGGT TATTCGGCG GTAATCAAAT ACCGCCGGTT
51   GTTTTTTGAA GTATTGGTGG TGTCGGTGGT GTTGCAGCTG TTTGCGCTGA
101  TTACGCCCTCT GTTTTTCCTA GTGGTGATGG ACAAGGTGCT GGTACATCGG
151  GGATTCTCTA CTTTGGATGT GGTGTCGGTG GCTTTGTTGG TGGTGTGCGT
201  GTTTGAGATT GTGTTGGGCG GTTTGCGGAC GTATCTGTTT GCACATACGA
251  CTTACAGTAT TGATGTGGAA TTGGGCGCGC GTTGTTCCTG GCATCTGCTT
301  TCCCTGCCTT TATCCTATTT CGAGCACAGA CGAGTGGGTG ATACGGTGGC
351  TCGGGTGCGG GAATTGGAGC AGATTGCGAA TTTCTTGACC GGTGAGGCGC
401  TGAATTCCGT GTTGGATTG GCGTTTTCGT TTATCTTCTT GCGCGTGATG
451  TGGTATTACA GCTCCACTCT GACTTGGGTG GTATTGGCTT CGTTG.....

1451 .....
1501 ..... ..ATTGCGC
1551 CAACCGGACG GTGCTGATTA TCGCCACCG TCTGTCCACT GTTAAACGG
1601 CACACCGGAT CATTGCCATG GATAAAGGCA GGATTGTGGA AGCGGGAACA
1651 CAGCAGGAAT TGCTGGCGAA CG..AACGGA TATTACCGCT ATCTGTATGA
1701 TTTACAGAAC GGGTAG

```

This corresponds to the amino acid sequence <SEQ ID 26; ORF39>:

```

25   1    ..KFDFTWFIPA VIKYRRFFE VLVVSVVLQL FALITPLFFQ VVMDKVLVHR
51   51   GFSTLDVVSF ALLVVSLEFI VLGGRLTYLF AHTTSRIDVE LGARLFRHLL
101  101  SLPLSYFEHR RVGDTVARVR ELEQIRNFLT GQALTSVLDL AFSFIFLAVM
151  151  WYYSSTLTWV VLASL.....

30   501  ..... ..ICANRT VLIIAHRLST VKTAHRIAM DKGRIVEAGT
551  551  QQELLANXNG YYRYLYDLQN G*

```

Further work revealed the complete nucleotide sequence <SEQ ID 27>:

```

35   1    ATGTCTATCG TATCCGCACC GCTCCCGGCC CTTTCCGCC TCATCATCCT
51   51   CGCCCATTA CACGGCATTG CCGCCAATCC TGCCGATATA CAGCATGAAT
101  101  TTTGTACTTC CGCACAGAGC GATTTAAATG AAACGCAATG GCTGTAGCC
151  151  GCCAAATCTT TGGGATTGAA GGCAAAGGTA GTCCGCCAGC CTATTAAACG
201  201  TTTGGCTATG GCGACTTTAC CCGCATTGGT ATGGTGTGAT GACGGCAACC
251  251  ATTTCAATTT GGCCAAAACA GACGGTGAGG GTGAGCATGC CCAATTTTGT
301  301  ATACAGGATT TGGTTACGAA TAAGTCTGCG GTATTGTCTT TTGCGGAATT
351  351  TTCTAACAGA TATTCGGGCA AACTGATATT GGTGCTTCC CGCGCTTCGG
401  401  TATTGGGACG TTGGCAAAG TTTGACTTTA CCTGGTTTAT TCCGGCGGTA
451  451  ATCAAATACC GCGGTTGTT TTTTGAAGTA TTGGTGGTGT CGGTGGTGTG
501  501  GCAGCTGTTT GCGCTGATTA CGCCTCTGTT TTTCCAAGTG GTGATGGACA
551  551  AGGTGCTGGT ACATCGGGGA TTCTCTACTT TGGATGTGGT GTCGGTGGCT
601  601  TTTTGGTGG TGTCCTGTT TGAGATTGTG TTGGGCGGTT TGCGGACGTA
651  651  TCTGTTTGCA CACGACTT CACGTATTGA TGTGGAATTG GCGCGCGGTT
701  701  TGTTCCGGCA TCTGCTTCC CTGCTTTTAT CCTATTTCGA GCACAGACGA
751  751  GTGGGTGATA CGGTGGCTCG GGTGCGGGAA TTGGAGCAGA TTCGCAATTT
801  801  CTTGACCGGT CAGGCGCTGA CTTGCTGTT GGATTGGCG TTTTCTTTA
851  851  TCTTTCTGGC GGTGATGTGG TATTACAGCT CCACTCTGAC TTGGGTGGTA
901  901  TTGGCTTCGT TGCCTGCCTA TGCGTTTGG TCGGCATTTA TCAGTCCGAT
951  951  ACTGCGGACG CGTCTGAACG ATAAGTTCGC GCGCAATGCA GACAACCACT
1001 1001 CGTTTTTAGT AGAAAGCATC ACTGCGGTGG GTACGGTAAA GGCGATGGCG
1051 1051 GTGGAGCCGC AGATGACGCA GCGTTGGGAC AATCAGTTGG CGGCTTATGT

```

1101 GGCTTCGGGA TTTCGGGTAA CGAAGTTGGC GGTGGTCGGC CAGCAGGGGG
 1151 TGCAGCTGAT TCAGAAGCTG GTGACGGTGG CGACGTTGTG GATTGGCGCA
 1201 CGGCTGGTAA TTGAGAGCAA GCTGACGGTG GGGCAGCTGA TTGCGTTTAA
 1251 TATGCTCTCG GGACAGGTGG CGGCGCCTGT TATCCGTTTG GCGCAGTTGT
 1301 GGCAGGATTT CCAGCAGGTG GGGATTTTCG TGGCGCGTTT GGGGGATATT
 1351 CTGAATGCGC CGACCGAGAA TCGCTCTTCG CATTGCGCTT TGCCCCGATAT
 1401 CCGGGGGGAG ATTACGTTCG AACATGTCGA TTTCCGCTAT AAGGCGGACG
 1451 GCAGGCTGAT TTTGCAGGAT TTGAACCTGC GGATTCGGGC GGGGGAAGTG
 1501 CTGGGGATTG TGGGACGTTT GGGGTCGGGC AAATCCACAC TCACCAAATT
 1551 GGTGCAGCGT CTGTATGTAC CGGAGCAGGG ACGGGTGTG GTGGACGGCA
 1601 ACGATTGGC TTTGGCCGCT CCTGCCTGGC TGGCGGCGCA GGTCCGGCTG
 1651 GTCTTGCAGG AGAATGTGCT GCTCAACCGC AGCATAACGC ACAATATCGC
 1701 GCTGACGGAT ACGGGTATGC CGCTGGAACG CATTATCGAA GCAGCCAAAC
 1751 TGGCGGGCGC ACACGAGTTT ATTATGGAGC TGCCGGAAGG CTACGGCACC
 1801 GTGGTGGGCG AACACGGGCG CGGCTTGTCT GCGCGACAGC GGCAGCGTAT
 1851 TCGGATTGCC CGCGCGTTAA TCACCAATCC GCGCATTCTG ATTTTGTATG
 1901 AAGCCACCAG CGCGCTGGAT TATGAAAGTG AACGAGCGAT TATGCAGAAC
 1951 ATGACAGGCA TTGCGCCAA CCGGACGGTG CTGATTATCG CCCACCGTCT
 2001 GTCCACTGTT AAAACGGCAC ACCGGATCAT TGCCATGGAT AAGGCGAGGA
 2051 TTGTGGAAGC GGGAACACAG CAGGAATTGC TGGCGAAGCC GAACCGATAT
 2101 TACCGCTATC TGTATGATTT ACAGAACGGG TAG

This corresponds to the amino acid sequence <SEQ ID 28; ORF39-1>:

1 MSIVSAPLPA LSALIILAHY HGIAANPADI QHEFCTSAQS DLNETOWLLA
 51 AKSLGLKAKV VRQPIKRLAM ATLPALVWCD DGNHFILAKT DGEHEHAQFL
 101 IQDLVTNKSA VLSFAEFSNR YSGKLILVAS RASVLGSLAK FDFTFWIPAV
 151 IKYRRFFFEV LVVSVVLQLF ALITPLFFQV VMDKVLVHRG FSTLDVVSVA
 201 LLVVSLEIV LGGLRTYLFA HTTSRIDVEL GARLFRHLLS LPLSYFEHRR
 251 VGDTVARVRE LEQIRNFLTG QALTSVLDLA FSFIFLAVMW YYSSTLTWVV
 301 LASLPAYAEW SAFISPIRLT RLNDKFARNA DNQSFIVESI TAVGTVKAMA
 351 VEPQMTQRWD NQLAAYVASG FRVTKLAVVG QQGVQLIQKL VTVATLWIGA
 401 RLVIESKLT VQLIAFNMLS GQVAAPVIRL AQLWQDFQOV GISVARLGDI
 451 LNAPTENASS HLALPDIRGE ITFEHVDFRY KADGRLILQD LNLRIAGEV
 501 LGIVGRSGSG KSTLTCLKVQR LYVPEQGRVL VDGNDLALAA PAWLRRQVGV
 551 VLQENVLLNR SIRDNIATLD TGMPLERIE AAKLAGAHEF IMELPEGYGT
 601 VVGEQGAGLS GGQRQRIIA RALITNPRIL IFDEATSALD YESERAIMQN
 651 MQAICANRTV LIIAHLSTV KTAHRIIAMD KGRIVEAGTQ QELLAKPNGY
 701 YRYLYDLQNG *

Computer analysis of this amino acid sequence gave the following results:

Homology with a predicted ORF from *N.meningitidis* (strain A)

ORF39 shows 100% identity over a 165aa overlap with an ORF (ORF39a) from strain A of *N. meningitidis*:

orf39.pep 10 20 30
 KFDFTWFIPAVIKYRRLEFEVLVVSVVVLQ
 45 orf39a AVLSFAEFSNRYSGKLILVASRASVLGSLAKFDFTWFIPAVIKYRRLEFEVLVVSVVVLQ
 110 120 130 140 150 160
 orf39.pep 40 50 60 70 80 90
 FALITPLFFQVMDKVLVHRGFSTLDVVSVALLVVSLEIVLGGLRTYLFAHTTSRIDVE
 50 orf39a FALITPLFFQVMDKVLVHRGFSTLDVVSVALLVVSLEIVLGGLRTYLFAHTTSRIDVE
 170 180 190 200 210 220
 orf39.pep 100 110 120 130 140 150
 LGARLFRHLLSLPLSYFEHRRVGDTVARVRELEQIRNFLTGOALTSVLDLAFSFI FLAVM
 55 orf39a LGARLFRHLLSLPLSYFEHRRVGDTVARVRELEQIRNFLTGOALTSVLDLAFSFI FLAVM
 230 240 250 260 270 280
 60 orf39.pep 160 170 180 190 200 210
 WYYSSTLTWVVLASLXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXICANRTVLIIAHLSTV

orf39a |||||
 WYYSSTLTWVVLASLPAYAFWSAFISPILRTRLNDKFARNADNQSFIVESITAVGTVKAM
 290 300 310 320 330 340

ORF39-1 and ORF39a show 99.4% identity in 710 aa overlap:

5	orf39-1.pep	MSIVSAPLPALSALIILAHYHGIAANPADIQHEFCTSAQSDLNQWLLAAKSLGLKAKV
	orf39a	MSIVSAPLPALSALIILAHYHGIAANPADIQHEFCTSAQSDLNQWLLAAKSLGLKAKV
10	orf39-1.pep	VRQPIKRLAMATLPALVWCDDGNHFILAKTDGEGEHAQFLIQDLVTNKSAVLSFAEFSNR
	orf39a	VRQPIKRLAMATLPALVWCDDGNHFILAKTDGEGEHAQFLIQDLVTNKSAVLSFAEFSNR
15	orf39-1.pep	YSGKLILVASRASVLGSLAKFDTWFIPAVIKYRRLFFEVLVSVVLQLFALITPLFFQV
	orf39a	YSGKLILVASRASVLGSLAKFDTWFIPAVIKYRRLFFEVLVSVVLQLFALITPLFFQV
20	orf39-1.pep	VMDKVLVHRGFSTLDVVSVALLVVSLFEIVLGGRLTYLFAHTTSRIDVELGARLFRHLLS
	orf39a	VMDKVLVHRGFSTLDVVSVALLVVSLFEIVLGGRLTYLFAHTTSRIDVELGARLFRHLLS
25	orf39-1.pep	LPLSYFEHRRVGDTVARVRELEQIRNFLTQALTSVLDLAFSFI FLAVMWYYSSTLTWV
	orf39a	LPLSYFEHRRVGDTVARVRELEQIRNFLTQALTSVLDLAFSFI FLAVMWYYSSTLTWV
30	orf39-1.pep	NQLAAYVASGFRVTKLAVVGQGVQLIQKLVTVATLWIGARLVIESKLTVGQLIAFNMLS
	orf39a	NQLAAYVASGFRVTKLAVVGQGVQLIQKLVTVATLWIGARLVIESKLTVGQLIAFNMLS
35	orf39-1.pep	GQVAAPVIRLAQLWQDFQQVGISVARLGDILNAPTENASSHLALPDIRGEITFEHVDFRY
	orf39a	GQVAAPVIRLAQLWQDFQQVGISVARLGDILNAPTENASSHLALPDIRGEITFEHVDFRY
40	orf39-1.pep	KADGRLILQDLNLRIRAGEVLGIVGRSGSGKSTLTKLVQRLYVPEQGRVLVDGNDLALAA
	orf39a	KADGRLILQDLNLRIRAGEVLGIVGRSGSGKSTLTKLVQRLYVPAQGRVLVDGNDLALAA
45	orf39-1.pep	PAWLRRQVGVLQENVLLNRSIRDNIALTDGTMPLERIIEAAKLAGAHEFIMELPEGYGT
	orf39a	PAWLRRQVGVLQENVLLNRSIRDNIALTDGTMPLERIIEAAKLAGAHEFIMELPEGYGT
50	orf39-1.pep	VVGEQGAGLSGGQRQRIARALITNPRILIFDEATSALDYESERAIMQNMQAICANRTV
	orf39a	VVGEQGAGLSGGQRQRIARALITNPRILIFDEATSALDYESERAIMQNMQAICANRTV
55	orf39-1.pep	LIIAHRLSTVKTAHRIIAMDKGRIVEAGTQQELLAKPNGYYRYLYDLQNGX
	orf39a	LIIAHRLSTVKTAHRIIAMDKGRIVEAGTQQELLAKPNGYYRYLYDLQNGX

The complete length ORF39a nucleotide sequence <SEQ ID 29> is:

1	ATGTCTATCG	TATCCGCACC	GCTCCCCGCC	CTTCCGCC	TCATCATCCT
55	51	CGCCCATAC	CACGGCATTG	CCGCCAATCC	TGCCGATATA
	101	TTTGTAATC	CGCACAGAGC	GATTTAATG	AAACGCAATG
	151	GCCAAATCTT	TGGGATTGAA	GGCAAAGGTA	GTCGCCAGC
	201	TTTGGCTATG	GCGACTTTAC	CCGCATTGGT	ATGGTGTGAT
	251	ATTTTATTTT	GGCTAAAACA	GACGGTGGGG	GTGAGCATGC
	301	ATACAGGATT	TAACACGAA	TAAGTCTGCG	GTATTGTCTT
60	351	TTCTAACAGA	TATTCGGGCA	AACTGATATT	GGTTGCTTCC
	401	TATTGGGCAG	TTTGGCAAAG	TTTGACTTTA	CCTGGTTTAT
	451	ATCAAATACC	GCCGGTTGTT	TTTTGAAGTA	TTGGTGGTGT
	501	GCAGCTGTTT	GCGCTGATTA	CGCCTCTGTT	TTTCCAAGTG
	551	AGGTGCTGGT	ACATCGGGGA	TTCTCTACTT	TGGATGTGGT
65	601	TTGTTGGTGG	TGTCGCTGTT	TGAGATTGTG	TTGGGCGGTT
	651	TCTGTTTGCA	CATACGACTT	CACGTATTGA	TGTGGAATTG

701 TGTTCGGCA TCTGCTTTCC CTGCCTTTAT CCTATTTTCA GCACAGACGA
 751 GTGGGTGATA CGGTGGCTCG GGTGCGGGAA TTGGAGCAGA TTCGCAATTT
 801 CTTGACCGGT CAGGCGCTGA CTTCCGTGTT GGATTGTCG TTTTCGTTTA
 851 TCTTTCTGGC GGTGATGTGG TATTACAGCT CCACCTGAC TTGGGTGGTA
 901 TTGGCTTCGT TGCCTGCCTA TCGCTTTTGG TCGGCATTA TCAGTCCGAT
 951 ACTGCGGACG CGTCTGAACG ATAAGTTTCG GCGCAATGCA GACAACCACT
 1001 CGTTTTTAGT AGAAAGCATC ACTGCGGTGG GTACGGTAAA GGCATGGCG
 1051 GTGGAGCCGC AGATGACGCA GCGTTGGGAC AATCAGTTGG CGGCTTATGT
 1101 GGCTTCGGGA TTTCCGGTAA CGAAGTTGGC GGTGGTCGCG CAGCAGGGGG
 1151 TGCAGCTGAT TCAGAGCTG GTGACGGTGG CGACGTTGTG GATTGGCGCA
 1201 CGGCTGGTAA TTGAGAGCAA GCTGACGGTG GGGCAGCTGA TTGCGTTTAA
 1251 TATGCTCTCG GGACAGGTGG CGGCGCTGT TATCCGTTG GCGCAGTTGT
 1301 GGCAGGATTT CCAGCAGGTG GGGATTTCGG TGGCGCGTTT GGGGAGTAT
 1351 CTGAATGCGC CGACCGAGAA TCGCTCTTCG CATTTGGCTT TGCCCGATAT
 1401 CCGGGGGGAG ATTACGTTTC AACATGTCGA TTTCCGCTAT AAGGCGGACG
 1451 GCAGGCTGAT TTGACAGGAT TTGAACCTGC GGATTCGGGC GGGGGAAGTG
 1501 CTGGGGATTG TGGGACGTTT GGGGTTCGGC AAATCCACAC TCACCAAATT
 1551 GGTGCAGCGT CTGTATGTAC CGGCGCAGGG ACGGGTGTG GTGGACGGCA
 1601 ACGATTTGGC TTTGCCCGCT CCTGCTTGGC TGCGGCGGCA GGTTCGGCTG
 1651 GTCTTGACAG AGAATGTGCT GCTCAACCGC AGCATAACCG ACAATATCGC
 1701 GCTGACGGAT ACGGCTATGC CGCTGGAACG CATTATCGAA GCAGCCAAAC
 1751 TGGCGGGCGC ACACGAGTTT ATTATGGAGC TGCCGGAAGG CTACGGCACC
 1801 GTGGTGGGCG AACAGGGGCG CGGCTGTTCG GGGCGACAGC GGCAGCGTAT
 1851 TGCATTGCC CGCGGTTTAA TCACCAATCC GCGCATTCTG ATTTTGTATG
 1901 AAGCCACCAG CGCGCTGGAT TATGAAAGTG AACGAGCGAT TATGCAAGC
 1951 ATGCAGGCCA TTTGCGCCAA CCGGACGGTG CTGATTATCG CCCACCGTCT
 2001 GTCCACTGTT AAAACGGCAC ACCGGATCAT TGCCATGGAT AAAGGCAGGA
 2051 TTGTGGAAGC GGAACACAG CAGGAATTGC TGGCGAAGCC GAACGGATAT
 2101 TACCGCTATC TGTATGATT ACAGAACGGG TAG

30 This encodes a protein having amino acid sequence <SEQ ID 30>:

1 MSIVSAPLPA LSALIILAHY HGIAANPADI QHEFCTSAQS DLNETOWLLA
 51 AKSLGLKAKV VRQPIKRLAM ATLPALVWCD DGNHFILAKT DGGGEHAQYL
 101 IQDLTTNKS A VLSFAEFSNR YSGKLILVAS RASVLGSLAK FDFWTFIPAV
 151 IKYRRLFFEV LVSVVLQLF ALITPLFFQV VMDKVLVHRG FSTLDVVSVA
 201 LLVVSLEFIV LGLRTRYLFA HTTSRIDVEL GARLFRHLLS LPLSYFEHRR
 251 VGDTVARVRE LEQIRNFLTQ QALTSVLDLA FSFIFLAVMW YYSSTLTWVV
 301 LASLPAYAFW SAFISPIRLT RLNDKFARNA DNQSFIVESI TAVGTVKAMA
 351 VEPQMTQRWD NQLAAYVASG FRVTKLAVVG QQGVQLIQKL VTVATLWIGA
 401 RLVIKSLTV GQLIAFNMLS GOVAAPVIRL AOLWQDFQOV GISVARLQDI
 451 LNAPTENASS HLALPDIRGE ITFEHVDFRY KADGRILQD LNLIRAGEV
 501 LGIVGRSGSG KSTLTKLVR LYVPAQGRVL VDGNDLALAA PAWLRRQGV
 551 VLQENVLLNR SIRDNIATD TGMPLERIE AAKLAGAHEF IMELPEGYGT
 601 VVGEQGAGLS GGRQRIIAIA RALITNPRIL IFDEATSALD YESERAIMON
 651 MQAICANRTV LIIAHLSTV KTAHRIIAMD KGRIVEAGTQ QELLAKPNYG
 701 YRYLYDLQNG *

ORF39a is homologous to a cytolysin from *A. pleuropneumoniae*:

sp|P26760|RT1B_ACTPL RTX-I TOXIN DETERMINANT B (TOXIN RTX-I SECRETION ATP-
 BINDING PROTEIN) (APX-IB) (HLY-IB) (CYTOLYSIN IB) (CLY-IB)
 >gi|97137|pir||D43599 cytolysin IB - Actinobacillus pleuropneumoniae (serotype 9)
 50 >gi|38944 (X61112) ClyI-B protein [Actinobacillus pleuropneumoniae] Length = 707
 Score = 931 bits (2379), Expect = 0.0
 Identities = 472/690 (68%), Positives = 540/690 (77%), Gaps = 3/690 (0%)
 55 Query: 20 YHGIAANPADIQHEFCTSAQSDLNETOWXXXXXXXXXXXXVVRQPIKRLAMATLPALVWC 79
 YH IA NP +++H+F + L+ T W V++ I RLA LPALVW
 Sbjct: 20 YHNIAVNPEELKHKFDLEGKG-LDTAWLLAAKSLELKAKQVKKAIDRLAFIALPALVWR 78
 Query: 80 DDGNHFILAKTDGGGEHAQYLIQDLTTNKS AVLSFAEFSNRYSGLKILVASRASVLGSLA 139
 +DG HFIL K D E +YLI DL T+ +L AEF + Y GKLILVASRAS++G LA
 60 Sbjct: 79 EDGKHFILTKIDN--EAKKYLI FDLTHNPRILEQAEFESLYQGKILILVASRASIVGKLA 136
 Query: 140 KFDFTWFIPAVIKYRXXXXXXXXXXXXXXXXXITPLFFQVVMKVLVHRGFXXXXXXXXX 199
 KFDFTWFIPAVIKYR+ ITPLFFQVVMKVLVHRGF
 65 Sbjct: 137 KFDFTWFIPAVIKYRKIFETLIVSIFLQIFALITPLFFQVVMKVLVHRGFSTLNVITV 196
 Query: 200 XXXXXXXFEIVLGLRTRYLFAHTTSRIDVELGARLFRHLLSLPLSYFEHRRVGDVARVR 259

FEIVL GLR TY+FAH+TSRIDVELGARLFRHLL+LP+SYFE+RRVGD TVARVR
 Sbjct: 197 ALAIVVLFEIVLNGLR TYIFAHSTSRIDVELGARLFRHLLALPISYFENRRVGD TVARVR 256

5 Query: 260 ELEQIRNFLT GQALTSVLDLAFSFI FLAVM WYYSSTLTWVVLASLPAYAFWSAFISPILR 319
 EL+QIRNFLT GQALTSVLDL FSFIF AVMWYYS LT V+L SLP Y WS FISPI LR
 Sbjct: 257 ELDQIRNFLT GQALTSVLDL MFSFI FFAVMWYYS PKLTLVILGSLPFYMGWSIFISPILR 316

10 Query: 320 TRLNDKFARNADNQSF LVESITAVGTVKAMAVEPQMTQRWDNQLAAYVASGFRVTKLAVV 379
 RL++KFAR ADNQSFLVES+TA+ T+KA+AV PQMT WD QLA+YV++GFRVT LA +
 Sbjct: 317 RRLDEKFARGADNQSF LVESVTAINTIKALAVTPQMTNTWDKQLASYVSAGFRVTTLATI 376

15 Query: 380 GQQGVQLIQKLVTATLWIGARLVIESKLT V GQLIAFNMLSGQVAAPVIRLAQLWQDFQQ 439
 GQQGVQ IQK+V V TLW+GA LVI L++GQLIAFNMLSGQV APVIRLAQLWQDFQQ
 Sbjct: 377 GQQGVQFIQKVVMVITLWLG AHLVISGDL SIGQLIAFNMLSGQVIAPVIRLAQLWQDFQQ 436

20 Query: 440 VGISVARLGDILNAP TENASSHLALPDIRGEITFEHVDFRYKADGRLLIQDLNLRIRAGE 499
 VGISV RLGD+LN+PTE+ LALP+I+G+ITF ++ FRYK D +IL D+NL I+ GE
 Sbjct: 437 VGISVTRLGDV LNSPTESYQ GK LALPEIKGDITFRNIRFRYKPDAPVILNDVNLSIQQGE 496

25 Query: 500 VLGIVGRSGSGKSTLT KL VQR LYP AQGRVLVDGNDLALAAPAWLRRQVG VVLQENVLLN 559
 V+GIVGRSGSGKSTLT KL+QR Y+P G+VL+DG+DLALA P WLRQVG VVLQ+NVLLN
 Sbjct: 497 VIGIVGRSGSGKSTLT KL IQR FYI PENGQVLIDGHDLALADPNWLRQVG VVLQDNVLLN 556

30 Query: 560 RSIRDNI ALTDTGMPLERIIEAAKLAGAHEFIMELPEGYGT VVGEQAGLSGGQRQRIAI 619
 RSIRDNI AL D GMP+E+I+ AAKLAGAHEFI EL EGY T+VGEQAGLSGGQRQRIAI
 Sbjct: 557 RSIRDNI ALADPGMPMEKIVHA AKLAGAHEFISELREGYNTIVGEQAGLSGGQRQRIAI 616

35 Query: 620 ARALITNPRILIFDEATSALDYESERAI MQNMQAICANRTVLIIAHRLSTVKT AHRIIAM 679
 ARAL+ NP+ILIFDEATSALDYESE IM+NM IC RTV+IIAHRLSTVK A RII M
 Sbjct: 617 ARALVNNPKILIFDEATSALDYESEHI IMRNMHQICKGR TVIIIAHRLSTVKNADRIIVM 676

Query: 680 DKGRIVEAGTQQELLAKPNGY YRYLYDLQN 709
 +KG+IVE G +ELLA PNG Y YL+ LQ+
 Sbjct: 677 EKGQIVEQ GKHKELLADPNGLYHYLHQLQS 706

Homology with the HlyB leucotoxin secretion ATP-binding protein of *Haemophilus actinomycescomitans* (accession number X53955)

ORF39 and HlyB protein show 71% and 69% amino acid identity in 167 and 55 overlap at the N- and C-terminal regions, respectively:

40 Orf39 1 KFDFTWFIPAVIKYRXXXXXXXXXXXXXXXXXITPLFFQVVM D KVLVHRGFXXXXXXXXXX 60
 KFDFTWFIPAVIKYR+ ITPLFFQVVM D KVLVHRGF
 HlyB 137 KFDFTWFIPAVIKYRKIF IETLIVSIFLQIFALITPLFFQVVM D KVLVHRGFSTLNVITV 196

45 Orf39 61 XXXXXXXFEIVLGG LRTYLF AHTTSRIDVELGARLFRHLLSLPLSYFEHRRVGD TVARVR 120
 FEI+LGGLR TY+FAH+TSRIDVELGARLFRHLL+LP+SYFE RRVGD TVARVR
 HlyB 197 ALAIVVLFEIILGG LRTYVFAHSTSRIDVELGARLFRHLLALPISYFEARRVGD TVARVR 256

50 Orf39 121 ELEQIRNFLT GQALTSVLDLAFSFI FLAVM WYYSSTLTWVVLASLIC 167
 EL+QIRNFLT GQALTS+LDL FSFIF AVMWYYS LT VVL SL C
 HlyB 257 ELDQIRNFLT GQALTSILDLLFSFI FFAVMWYYS PKLTLVVLGSLPC 303

//

55 Orf39 166 ICANRTVLIIAHRLSTVKT AHRIIAMDKGRIVEAGTQQELLANXNGY YRYLYDLQ 220
 IC NRTVLIIAHRLSTVK A RII MDKG I+E G QELL + G Y YL+ LQ
 HlyB 651 ICQNRTVLIIAHRLSTVKNADRIIVMDKGEIIEQ GKHQELLKDEKGLSYLHQLQ 705

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 7

60 The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 31>

```

1  ATGAAATACT TGATCCGCAC CGCCTTACTC GCAGTCGCAG CCGCCGGCAT
51 CTACGCCTGC CAACCGCAAT CCGAAGCCGC AGTGCAAGTC AAGGCTGAAA
101 ACAGCCTGAC CGCTATGCGC TTAGCCGTCG CCGACAAACA GGCAGAGATT
151 GACGGGTGA ACGCCCAAAC sGACGCCGAA ATCAGA...

```

5 This corresponds to the amino acid sequence <SEQ ID 32; ORF52>:

```

1  MKYLIRTALL AVAAAGIYAC QPQSEAAVQV KAENSLTAMR LAVADKQAEI
51 DGLNAQXDAE IR..

```

Further work revealed the complete nucleotide sequence <SEQ ID 33>:

```

10      1  ATGAAATACT TGATCCGCAC CGCCTTACTC GCAGTCGCAG CCGCCGGCAT
        51 CTACGCCTGC CAACCGCAAT CCGAAGCCGC AGTGCAAGTC AAGGCTGAAA
        101 ACAGCCTGAC CGCTATGCGC TTAGCCGTCG CCGACAAACA GGCAGAGATT
        151 GACGGGTGA ACGCCCAAAT CGACGCCGAA ATCAGACAAC GCGAAGCCGA
        201 AGAATTGAAA GACTACCGAT GGATACACGG CGACGCCGAA GTGCCGGAGC
        251 TGGAAAAATG A

```

15 This corresponds to the amino acid sequence <SEQ ID 34; ORF52-1>:

```

1  MKYLIRTALL AVAAAGIYAC QPQSEAAVQV KAENSLTAMR LAVADKQAEI
51 DGLNAQIDAE IRQREAEELK DYRWIHGDAE VPELEK*

```

Computer analysis of this amino acid sequence predicts a prokaryotic membrane lipoprotein lipid attachment site (underlined).

20 ORF52-1 (7kDa) was cloned in the pGex vectors and expressed in *E.coli*, as described above. The products of protein expression and purification were analyzed by SDS-PAGE. Figure 4A shows the results of affinity purification of the GST-fusion. Figure 4B shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF52-1.

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could
25 be useful antigens for vaccines or diagnostics.

Example 8

The following DNA sequence was identified in *N.meningitidis* <SEQ ID 35>

```

30      1  ATGGTTATCG GAATATTACT CGCATCAAGC AAGCATGCTC TTGTCATTAC
        51 TCTATTGTTA AATCCCGTCT TCCATGCATC CAGTTGCGTA TCGCGTTsGG
        101 CAATACGGAA TAAAAtCTGC TGTTC TGCTT TGGCTAAATT TGCCAAATTG
        151 TTTATTGTTT CTTTAGGAGC AGCTTGCTTA GCCGCCTTCG CTTTCGACAA
        201 CGCCCCCACA GGCGCTTCCC AAGCgTTGCC TACCGTTACC GCACCCGTGG
        251 CGATTCCCGC GCCCGCTTCG GCAGCCTGA

```

This corresponds to the amino acid sequence <SEQ ID 36; ORF56>:

```

35      1  MVIGILLASS KHALVITLLL NPVFHASSCV SRXAIRNKIC CSALAKFAKL
        51 FIVSLGAACL AAFAFDNAPT GASQALPTVT APVAIPAPAS AA*

```

Further work revealed the complete nucleotide sequence <SEQ ID 37>:

```

1  ATGGCTTGTA CAGGTTTGAT GGTTTTCCG TTAATGGTTA TCGGAATATT

```

51 ACTTGCATCA AGCAAGCCTG CTCCTTTCCT TACTCTATTG TTAAATCCCG
 101 TCTTCCATGC ATCCAGTTGC GTATCGCGTT GGGCAATACG GAATAAAATC
 151 TGCTGTTCTG CTTTGGCTAA ATTTGCCAAA TTGTTTATTG TTTCTTTAGG
 201 AGCAGCTTGC TTAGCCGCCT TCGCTTTCGA CAACGCCCCC ACAGCGCCTT
 251 CCCAAGCGTT GCCTACCGTT ACCGCACCCG TGGCGATTCC CGCGCCCGCT
 301 TCGGCAGCCT GA

This corresponds to the amino acid sequence <SEQ ID 38; ORF56-1>:

1 MACTGLMVFP LMVIGILLAS SKPAPFLTLL LNPVFHASSC VSRWAIRNKI
 51 CCSALAKFAK LFIVSLGAAC LAFAFDNAP TGASQALPTV TAPVAIPAPA
 101 SAA*

Computer analysis of this amino acid sequence predicts a leader peptide (underlined) and suggests that ORF56 might be a membrane or periplasmic protein.

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

15 Example 9

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 39>

1 ATGTTCACTA TTTTAAATGT GTTCTTCAT TGTATTCTGG CTTGTGTAGT
 51 CTCTGGTGAG ACGCCTACTA TATTTGGTAT CCTTGCTCTT TTTTACTTAT
 101 TGTATCTTTC TTATCTTGCT GTTTTTAAGA TTTTCTTTC TTTTCTTCTA
 151 GACAGAGTTT CACTCCGGTC TCCCAGGCTG GAGTGCAAAT GGCATGACCC
 201 TTTGGCTCAC TGGCTCACGG CCACTTCTGC TATTCTGCCG CCTCAGCCTC
 251 CAGGG...

This corresponds to the amino acid sequence <SEQ ID 40; ORF63>:

1 MFSILNVFLH CILACVVSGE TPTIFGILAL FYLLYLSYLA VFKIFFSFFL
 251 DRVSLRSPRL ECKWHDPLAH WLTATSAILP PQPPG...

Computer analysis of this amino acid sequence predicts a transmembrane region.

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 10

30 The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 41>

1 ..GTGCGGACGT GGTGGGTTTT TTGGTTGCAG CGTTTGAAAT ACCCGTTGTT
 51 GCTTTGGATT GCGGATATGT TGCTGTACCG GTTGTGGGC GGCGCGGAAA
 101 TCGAATGCGG CCGTTGCCCT GTGCCGCCGA TGACGGATTG GCAGCATTTT
 151 TTGCCGGCGA TGGGAACGGT GTCGGCTTGG GTGGCGGTGA TTTGGGCATA
 35 201 CCTGATGATT GAAAGTGAAG AAAACGGAAG ATATTGA

This corresponds to the amino acid sequence <SEQ ID 42; ORF69>:

1 ..VRTWLVEWLQ RLKYPALLWI ADMLLYRLLG GAEIECGRCP VPPMTDWQHF
 51 LPAMGTVSAW VAVIWAYLMI ESEKNGRY*

Computer analysis of this amino acid sequence predicts a transmembrane region.

A corresponding ORF from strain A of *N.meningitidis* was also identified:

Homology with a predicted ORF from *N.meningitidis* (strain A)

ORF69 shows 96.2% identity over a 78aa overlap with an ORF (ORF69a) from strain A of *N.*

5 *meningitidis*:

```

      10      20      30      40      50      60
orf69.pep VRTWLVFQWLQRLKYPLLLWIADMLLYRLLGAEIECGRCVPVPMTDWQHFLPAMGTVSAW
          |||||
orf69a    VRTWLVFQWLQRLKYPLLLCIADMLLYRLLGAEIECGRCVPVPMTDWQHFLPTMGTVAAW
          10      20      30      40      50      60

      70      79
orf69.pep VAVIWAYLMIESEKNGRYX
          |||||
orf69a    VAVIWAYLMIESEKNGRYX
          70
  
```

The ORF69a nucleotide sequence <SEQ ID 43> is:

```

      1  GTGCGGACGT GGTGGTTTT TTGGTTGCAG CGTTTGAAAT ACCCGTTGTT
     51  GCTTTGTATT GCGGATATGC TGCTGTACCG GTTGTGGGC GCGCGGAAA
    101  TCGAATGCGG CCGTTGCCCT GTACCGCCGA TGACGATTG GCAGCATTTT
    151  TTGCCGACGA TGGGAACGGT GCGGCTTGG GTGGCGGTGA TTTGGGCATA
    201  CCTGATGATT GAAAGTGAAA AAAACGGAAG ATATTGA
  
```

This encodes a protein having amino acid sequence <SEQ ID 44>:

```

     1  VRTWLVFQWLQ RLKYPLLLCI ADMLLYRLLG GAEIECGRCP VPPMTDWQHF
    51  LPTMGTVAAW VAVIWAYLMI ESEKNGRY*
  
```

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 11

30 The following DNA sequence was identified in *N.meningitidis* <SEQ ID 45>

```

      1  ATGTTTCAAA ATTTTGATT GGGCGTGTT CTGCTTGCCG TCCTCCCCGT
     51  GCTGCCCTCC ATTACCGTCT CGCACGTGGC GCGCGGCTAT ACGGCGCGCT
    101  ACTGGGGAGA CAACACTGCC GAACAATACG GCAGGCTGAC ACTGAACCCC
    151  CTGCCCCATA TCGATTTGGT CGGCACAATC ATCgTACCGC TGCTTACTTT
    201  GATGTTACAG CCCTTCCTGT TCGGCTGGGC GCGTCCGATT CCTATCGATT
    251  CGCGCAACTT CCGCAACCCG cGCCTTGCTT GCGGTGCGT TGCCGCGTCC
    301  GGCCCGCTGT CGAATCTAGC GATGGCTGTW CTGTGGGGCG TGGTTTGGT
    351  GCTGACTCCG TATGTCGGCG GGGCGTATCA GATGCCGTTG GCTCAAATGG
    401  CAAACTACGG TATTCTGATC AATGCGATTG TGTTGCGGCT CAACATCATC
    451  CCCATCCTGC CTTGGGACGG CGGCATTTTC ATCGACACCT TCCTGTCGGC
    501  GAAATATTCG CAAGCGTTCC GCAAAATCGA ACCTTATGGG ACGTGGGATTA
    551  TCCTACTGCT GATGCTGACC sGGGTTTTGG GTGCGTTTAT WGCACCGATT
    601  sTGCGGmTGc GTGATTGCrT TTGTGCAGAT GTwCGTCTGA CTGGCTTCA
    651  GACGGCATAA
  
```

45 This corresponds to the amino acid sequence <SEQ ID 46; ORF77>:

Further work revealed the complete nucleotide sequence <SEQ ID 47>:

15

20 This corresponds to the amino acid sequence <SEQ ID 48; ORF77-1>:

25

Computer analysis of this amino acid sequence reveals a putative leader sequence and several transmembrane domains.

A corresponding ORF from strain A of *N.meningitidis* was also identified:

Homology with a predicted ORF from *N.meningitidis* (strain A)

30 ORF77 shows 96.5% identity over a 173aa overlap with an ORF (ORF77a) from strain A of *N. meningitidis*:

35

```

      10      20      30      40      50      60
orf77.pep  MFQNFDLGVFLLAVLPVLP SITVSHVARGYTARYWGDNTAEQYGRLTLNPLPHIDLVGTI
            |||||||
orf77a      RGYTARYWGDNTAEQYGRLTLNPLPHIDLVGTI
            10      20      30

```

40

orf77.pep IVPLLTLMFTPFLEFGWARPIPIDSRNFRNPRLAWRCVAASGPLSNLAMAVLWGVVLVLTP
 |||||
 orf77a IVPLLTLMFTPFLEFGWARPIPIDSRNFRNPRLAWRCVAASGPLSNLAMAVLWGVVLVLTP
 40 50 60 70 80 90

45

130 140 150 160 170 180
 orf77.pep YVGGAYQMPLAQMANYGILINAILFALNIIPILPWDGGIFIDTFLSAKYSQAFRKIEPYG
 ||||| ||||| ||||| ||||| ||||| |||||
 orf77a YVGGAYQMPLAQMANYXILINAILXALNIIPILPWDGGIFIDTFLSAKXSQAFRKIEPYG
 100 110 120 130 140 150

50

190
200
210
220
 orf77.pep TWIILLMLTXXVLGAFIAPIXRXRDCXCADVRLTGFQTAX
 ||||| ||||| |||||
 orf77a TWIIXLLMLTGVLGAXIAPIVQLVIAFVQMFX
 160 170 180

ORF77-1 and ORF77a show 96.8% identity in 185 aa overlap:

```

      10      20      30      40      50      60
orf77-1.pep MFQNFDLGVFLLAVLPVLLSITVREVARGYTARYWGDNTAEQYGRLLNPLPHIDLVGTI
5  orf77a      RGYTARYWGDNTAEQYGRLLNPLPHIDLVGTI
      10      20      30
      70      80      90      100     110     120
orf77-1.pep IVPLLLTLMFTFPLFGWARPIPIDSRNFRNPRLAWRCVAASGPLSNLAMAVLWGVVLVLT
10 orf77a      IVPLLLTLMFTFPLFGWARPIPIDSRNFRNPRLAWRCVAASGPLSNLAMAVLWGVVLVLT
      40      50      60      70      80      90
      130     140     150     160     170     180
orf77-1.pep YVGGAYQMPLAQMANYGILINAILFALNIIPILPWDGGIFIDTFLSAKYSQAFRKIEPYG
15 orf77a      YVGGAYQMPLAQMANYGILINAILFALNIIPILPWDGGIFIDTFLSAKXSQAFRKIEPYG
      100     110     120     130     140     150
      190     200     210
orf77-1.pep TWIILLMLTGVLGAFIPIVRLVIAFVQMFVX
20 orf77a      TWIIXLLMLTGVLGAXIPIVQLVIAFVQMFVX
      160     170     180

```

A partial ORF77a nucleotide sequence <SEQ ID 49> was identified:

```

      1  ..CGCGGCTATA CAGCGCGCTA CTGGGGTGAC AACACTGCCG AACAAATACGG
51  CAGGCTGACA CTGAACCCCTC TGCCCATAT CGATTGGTC GGCACAATCA
30 101 TCGTACCGCT GCTTACTTTG ATGTTTACGC CCTTCCTGTT CGGCTGGGCG
151 CGTCCGATTC CTATCGATTC GCGCACTTC CGCAACCCGC GCCTTGCCCTG
201 GCGTTGCGTT GCCGCGTCCG GCCCGCTGTC GAATCTGGCG ATGGCTGTTT
251 TGTGGGGCGT GGTTTTGGTG CTGACTCCGT ATGTCGGTGG GCGGTATCAG
301 ATGCCGTTGG CNCAATGGC AAACACNNN ATTCTGATCA ATGCGATTCT
351 GTNCGCGCTC AACATCATCC CCATCCTGCC TTGGGACGGC GGCATTTTCA
35 401 TCGACACCTT CCTGTCGGCN AAATANTCGC AAGCGTTCCG CAAAATCGAA
451 CCTTATGGGA CGTGGATTAT CCNGCTGCTT ATGCTGACCG GGGTTTTGGG
501 TGGCTNTATT GCACCGATTG TGCAGCTGGT GATTGCGTTT GTGCAGATGT
551 TCGTCTGA

```

This encodes a protein having amino acid sequence <SEQ ID 50>:

```

40      1  ..RGYTARYWGD NTAEQYGRLL LNPLPHIDLV GTIIVPLLLT MFTFPLFGWA
51  RPIPIDSRNF RNPRLAWRCV AASGPLSNLA MAVLWGVVLV LTPYVGGAYQ
101 MPLAQMANYS ILINAILXAL NIIPILPWDG GIFIDTFLSA KXSQAFRKIE
151 PYGTWIIIXLL MLTGVLGAXI APIVQLVIAF VQMFV*

```

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 12

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 51>

```

50      1  ATGAACCTGA TTTCACGTTA CATCATCCGT CAAATGGCGG TTATGGCGGT
51  TTACGCGCTC CTTGCCTTCC TCGCTTTGTA CAGCTTTTTT GAAATCCTGT
101 ACGAAACCGG CAACCTCGGC AAAGGCAGTT ACGGCATATG GGAAATGCTG
151 GGCTACACCG CCCTCAAAAT GCCCGCCCGC GCCTACGAAC TGATTCCCCT
201 CGCCGTCCTT ATCGGCGGAC TGGTCTCCCT CAGCCAGCTT GCCGCGGGCA
251 GCGAACTGAC CGTCATCAA AAGAGCGGCA TGAGCACCAA AAAGCTGCTG
301 TTGATTCTGT CGCAGTTCGG TTTTATTTTT GCTATTGCCA CCGTCGCGCT
55 351 CGGCGAATGG GTTGCGCCCA CACTGAGCCA AAAAGCCGAA AACATCAAAG

```

401 CCGCCGCCAT CAACGGCAAA ATCAGCACCG GCAATACCGG CCTTTGGCTG
451 AAAGAAAAAA ACAGCGTGAT CAATGTGCGC GAAATGTTGC CCGACCAT..

This corresponds to the amino acid sequence <SEQ ID 52; ORF112>:

5 1 MNLSRYIIR QMAVMVYAL LAFLALYSFF EILYETGNLG KGSYGIWEML
51 GYTALKMPAR AYELIPLAVL IGGLVLSLSQL AAGSELTVIK ASGMSTKKLL
101 LILSQFGFIF AIATVALGEW VAPTLSQKAE NIKAAAINGK ISTGNTGLWL
151 KEKNSVINVR EMLPDH...

Further work revealed further partial nucleotide sequence <SEQ ID 53>:

10 1 ATGAACCTGA TTTCACGTTA CATCATCCGT CAAATGGCGG TTATGGCGGT
51 TTACGCGCTC CTTGCCTTCC TCGCTTTGTA CAGCTTTTTT GAAATCCTGT
101 ACGAAACCGG CAACCTCGGC AAAGGCAGTT ACGGCATATG GGAAATGCTG
151 gGCTACACCG CCTCAAAAT GCCCGCCCGC GCCTACGAAC TGATTCCCT
201 CGCGTCTCTT ATCGGCGGAC TGGTCTCCCT CAGCCAGCTT GCCGCCGGCA
15 251 GCGAACTGAC CGTCATCAAA GCCAGCGGCA TGAGCACCAA AAAGCTGCTG
301 TTGATTCTGT CGCAGTTCGG TTTTATTTTT GCTATTGCCA CCGTCGCGCT
351 CGGCGAATGG GTTGCGCCCA CACTGAGCCA AAAAGCCGAA AACATCAAAG
401 CCGCCGCCAT CAACGGCAAA ATCAGCACCG GCAATACCGG CCTTTGGCTG
451 AAAGAAAAAA ACAGCTTKAT CAATGTGCGC GAAATGTTGC CCGACCATAC
501 GCTTTTGGGC ATCAAAATTT GGGCGCGCAA CGATAAAAC GAATTGGCAG
20 551 AGGCAGTGGA AGCCGATTCC GCCGTTTGA ACAGCGACGG CAGTTGGCAG
601 TTGAAAAACA TCCGCCGCG CACGCTTGGC GAAGACAAAG TCGAGGTCTC
651 TATTGCGGCT GAAGAAAAC TGGCCGATTTC CGTCAAACGC AACCTGATGG
701 ACGTATTGCT CGTCAAACCC GACCAAATGT CCGTCGGCGA ACTGACCACC
751 TACATCCGCC ACCTCCAAAA CAACAGCCAA AACACCCGAA TCTACGCCAT
25 801 CGCATGGTGG CGCAAATTGG TTTACCCCGC CGCAGCCTGG GTGATGGCGC
851 TCGTCGCCTT TGCCTTTACC CCGCAAACCA CCGCCACGG CAATATGGGC
901 TTAAACTCT TCGGCGGCAT CTGTsTCGGA TTGCTGTTCC ACCTTGCCGG
951 ACGGCTCTTT GGGTTTACCA GCCAACTCGG...

This corresponds to the amino acid sequence <SEQ ID 54; ORF112-1>:

30 1 MNLSRYIIR QMAVMVYAL LAFLALYSFF EILYETGNLG KGSYGIWEML
51 GYTALKMPAR AYELIPLAVL IGGLVLSLSQL AAGSELTVIK ASGMSTKKLL
101 LILSQFGFIF AIATVALGEW VAPTLSQKAE NIKAAAINGK ISTGNTGLWL
151 KEKNSXINVR EMLPDHTLLG IKIWARNDKN ELAEAVEADS AVLNSDGSWQ
35 201 LKNIRRTLK EDKVEVSIAA EENWPISVKR NLMVLLVKP DQMSVGLTT
251 YIRHLQNSQ NTRIIAIAWW RKLVPAAAW VMALVAFAFT PQTRHGNMG
301 LKLFGGICXG LLFHLAAGRLF GFTSQL...

Computer analysis of this amino acid sequence predicts two transmembrane domains.

A corresponding ORF from strain A of *N.meningitidis* was also identified:

Homology with a predicted ORF from *N.meningitidis* (strain A)

40 ORF112 shows 96.4% identity over a 166aa overlap with an ORF (ORF112a) from strain A of *N.meningitidis*:

		10	20	30	40	50	60
orfl12.pep	MNLSRYIIRQMAVMVYALLAFLALYSFFEILYETGNLGKGSYGIWEMLGYTALKMPAR						
45	orfl12a	MNLSRYIIRQMAVMVYALLAFLALYSFFEILYETGNLGKGSYGIWEMXGYTALKMXAR					
		10	20	30	40	50	60
		70	80	90	100	110	120
50	orfl12.pep	AYELIPLAVLIGGLVLSLSQLAAGSELTVIKASGMSTKKLLLILSQFGFIFAIATVALGEW					
	orfl12a	AYELMPLAVLIGGLVLSLSQLAAGSELXVIKASGMSTKKLLLILSQFGFIFAIATVALGEW					
		70	80	90	100	110	120

```

              130      140      150      160
orf112.pep  VAPTLSQKAENIKAAAINGKISTGNTGLWLKEKNSVINVREMLPDH
              |||||
orf112a     VAPTLSQKAENIKAAAINGKISTGNTGLWLKEKNSIINVREMLPDHTLLGIKIWARNDKN
              130      140      150      160      170      180
orf112a     ELAEAVEADSAVLNSDGSWQLKNIRRLSTLGEDKVEVSIAAEEXWPISVKRNLMDVLLVKP
              190      200      210      220      230      240

```

A partial ORF112a nucleotide sequence <SEQ ID 55> was identified:

```

10      1  ATGAACCTGA  TTTCACGTTA  CATCATCCGT  CAAATGGCGG  TTATGGCGGT
      51  TTACGCGCTC  CTTGCCTTCC  TCGCTTTGTA  CAGCTTTTTT  GAAATCCTGT
     101  ACGAAACCGG  CAACCTCGGC  AAAGGCAGTT  ACGGCATATG  GGAAATGNTG
     151  GGNTACACCG  CCTCAAAAT  GNCCGCCCGC  GCCTACGAAC  TGATGCCCTT
     201  CGCGTCTCTT  ATCGGCGGAC  TGGTCTCTNT  CAGCCAGCTT  GCCGCCGGCA
     15      251  GCGAACTGAN  CGTCATCAAA  GCCAGCGGCA  TGAGACCCAA  AAAGCTGCTG
     301  TTGATTCTGT  CGCAGTTCGG  TTTTATTTTT  GCTATTGCCA  CCGTCGCGCT
     351  CGGCGAATGG  GTTGCGCCCA  CACTGAGCCA  AAAAGCCGAA  AACATCAAAG
     401  CCGCGGCCAT  CAACGGCAAA  ATCAGTACCG  GCAATACCGG  CCTTTGGCTG
     451  AAAGAAAAAA  ACAGCATTAT  CAATGTGCGC  GAAATGTTGC  CCGACCATAC
     20      501  CCTGCTGGGC  ATTAAATCT  GGGCCCGCAA  CGATAAAAC  GAACTGGCAG
     551  AGCGAGTGA  AGCCGATTCC  GCCGTTTTGA  ACAGCGACGG  CAGTTGGCAG
     601  TTGAAAAACA  TCCGCCGCGC  CACGCTTGCG  GAAGACAAAG  TCGAGGTCTC
     651  TATTGCGGCT  GAAGAAAANT  GGCCGATTTC  CGTCAAACGC  AACCTGATGG
     701  ACGTATTGCT  CGTCAAACCC  GACCAAATGT  CCGTCGGCGA  ACTGACCACC
     25      751  TACATCCGCC  ACCTCCAAAN  NNACAGCCAA  AACACCCGAA  TCTACGCCAT
     801  CGCATGGTGG  CGCAAATTGG  TTTACCCCGC  CGCAGCCTGG  GTGATGGCGC
     851  TCGTCGCCTT  TGCCTTTACC  CCGCAAACCA  CCCGCCACGG  CAATATGGGC
     901  TTAATAANTCT  TCGGCGGCAT  CTGTCTCGGA  TTGCTGTTCC  ACCTTGCCGG
     951  NCGGCTCTTC  NGGTTTACCA  GCCAACTCTA  CGGCATCCCG  CCCTTCCTCG
     30      1001  NCGGCGCACT  ACCTACCATA  GCCTTCGCCT  TGCTCGCCGT  TTGGCTGATA
     1051  CGCAAACAGG  AAAAACGCTA  A

```

This encodes a protein having amino acid sequence <SEQ ID 56>:

```

      1  MNLISRYIIR QMAVMVYAL LAFLALYSFF EILYETGNLG KGSYGIWEMX
     51  GYTALKMXAR AYELMPLAVL IGGLVSXSQ L AAGSELXVIK ASGMSTKKLL
     35      101  LILSQFGFIF AIATVALGEW VAPTLSQKAE NIKAAAINGK ISTGNTGLWL
     151  KEKNSIINVR EMLPDHTLLG IKIWARNDKN ELAEAVEADS AVLNSDGSWQ
     201  LKNIRRLSTL EDKVEVSIAA EEXWPISVKR NLMDVLLVKP DQMSVGELTT
     251  YIRHLQXXSQ NTRIYAIAWW RKLVPAAAW VMALVAFAPT PQTTRHGNMG
     301  LKXFGGICLG LLFHLAGRLF XFTSOLYGIP PFLXGALPTI AFALLAVWLI
     40      351  RKQEKR*

```

ORF112a and ORF112-1 show 96.3% identity in 326 aa overlap:

```

orf112a.pep  MNLISRYIIRQMAVMVYALLAFLALYSFFEILYETGNLGKGSYGIWEMXGYTALKMXAR
              |||||
orf112-1     MNLISRYIIRQMAVMVYALLAFLALYSFFEILYETGNLGKGSYGIWEMGYTALKMPAR
45
orf112a.pep  AYELMPLAVLIGGLVSXSQLAAGSELXVIKASGMSTKKLLLILSQFGFIFAIATVALGEW
              ||||:|||||
orf112-1     AYELIPLAVLIGGLVLSQLAAGSELTVIKASGMSTKKLLLILSQFGFIFAIATVALGEW
50
orf112a.pep  VAPTLSQKAENIKAAAINGKISTGNTGLWLKEKNSIINVREMLPDHTLLGIKIWARNDKN
              |||||
orf112-1     VAPTLSQKAENIKAAAINGKISTGNTGLWLKEKNSXINVREMLPDHTLLGIKIWARNDKN
55
orf112a.pep  ELAEAVEADSAVLNSDGSWQLKNIRRLSTLGEDKVEVSIAAEEXWPISVKRNLMDVLLVKP
              |||||
orf112-1     ELAEAVEADSAVLNSDGSWQLKNIRRLSTLGEDKVEVSIAAEENWPISVKRNLMDVLLVKP
60
orf112a.pep  DQMSVGELTTYIRHLQXXSQNTRIYAIAWWRKLVYPAAAWVMALVAFAPT PQTTRHGNMG
              |||||
orf112-1     DQMSVGELTTYIRHLQNNQNTRIYAIAWWRKLVYPAAAWVMALVAFAPT PQTTRHGNMG
orf112a.pep  LKXFGGICLGLLFHLAGRLF XFTSOLYGIP PFLXGALPTI AFALLAVWLIRKQEKRX
              || ||||

```


orf112-1

LKLFGGICXGLLFHLAGRLFGFTSQL

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

5 Example 13

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 57>

```

1  ..GCAGTAGCCG AAACTGCCAA CAGCCAGGGC AAAGGTAAAC AGGCAGGCAG
51  TTCGGTTTCT GTTTCACCTGA AAACCTCAGG CGACCTTTGC GGCAAACTCA
101 AAACCAACCT TAAAACTTTG GTCTGCTCTT TGGTTTCCCT GAGTATGGTA
151 TTGCCTGCCC ATGCCCAAAT TACCACCGAC AAATCAGCAC CTAAAAACCA
201 GCAGGTGCGT ATCCTTAAAA CCAACACTGG TGCCCCCTTG GTGAATATCC
251 AAATCCGAA TGGACGCGGA TTGAGCCACA ACCGCTA.TA CGCATTTGAT
301 GTTGACAACA AAGGGGCAGT GTTAAACAAC GACCGTAACA ATAATCCGTT
351 TGTGGTCAAA GGCAGTGCGC AATTGATTTT GAACGAGGTA CGCGTACGG
15 401 CTAGCAAACCT CAACGGCATC GTTACCGTAG GCGGTCAAAA GGCCGACGTG
451 ATTATTGCCA ACCCCAACGG CATTACCGTT AATGGCGGCG GCTTTAAAAA
501 TGTGCGTCGG GGCATCTTAA CTACCGGTGC GCCCCAAATC GGCAAAGACG
551 GTGCACTGAC AGGATTTGAT GTGCGTCAAG GCACATTGga CCGTAGrAGC
601 AGCAGGTTGG AATGATAAAG GCGGAGCmrm yTACACCGGG GTACTTGCTC
20 651 GTGCAGTTGC TTTGCAGGGG AAATTWmmGG GTAAA.AACT GGCAGTTTCT
701 ACCGGTCCTC AGAAAGTAGA TTACGCCAGC GGCGAAATCA GTGCAGGTAC
751 GCGAGCGGTG ACGAAACCGA CTATTGCCCT TGATACTGCC GCACCTGGCG
801 GTATGTACGC CGACAGCATC ACACTGATTG CCAATGAAAA AGGCGTAGGC
851 GTCTAA

```

25 This corresponds to the amino acid sequence <SEQ ID 58; ORF114>:

```

1  ..AVAETANSQG KGKQAGSSVS VSLKTSGLDL GKLTTLKTL VCSLVLSMV
51  LPAHAQITTD KSAPKNQQVV ILKNTNGAPL VNIQTPNGRG LSHNRXYAFD
101 VDNKGAVLNN DRNNNPFVVK GSAQLILNEV RGTASKLNGI VTVGGQKADV
151 IIANPNGITV NGGGFKNVGR GILTTGAPQI GKDGALTGFD VVKAHWTVXA
30 201 AGWNDKGGAX YTGVLARAVA LQGKXXGKXL AVSTGPQKVD YASGEISAGT
251 AAGTKPTIAL DTAALGMYA DSITLIANEK GVG*

```

Further work revealed the complete nucleotide sequence <SEQ ID 59>:

```

1  ATGAATAAAG GTTTACATCG CATTATCTTT AGTAAAAAGC ACAGCACCAT
35 51  GGTTGCAGTA GCCGAACTG CCAACAGCCA GGGCAAAGGT AAACAGGCAG
101 GCAGTTCCGT TTCTGTTTCA CTGAAAACTT CAGGCGACCT TTGCGCAAAA
151 CTCAAAACCA CCTTAAAC TTTGGTCTGC TCTTTGGTTT CCCTGAGTAT
201 GGTATTGCCCT GCCCATGCCC AAATTACCAC CGACAAATCA GCACCTAAAA
251 ACCAGCAGGT CGTTATCCTT AAAACCAACA CTGGTGCCCC CTTGGTGAAT
301 ATCCAACTC CGAATGGACG CGGATTGAGC CACAACCGCT ATACGCAGTT
40 351 TGATGTTGAC AACAAAGGGG CAGTGTTAAA CAACGACCGT AACATAATC
401 CGTTTGTGGT CAAAGGCAGT GCGCAATTGA TTTTGAACGA GGTACGCGGT
451 ACGGCTAGCA AACTCAACGG CATCGTTACC GTAGGCGGTC AAAAGGCCGA
501 CGTGATTATT GCCAACCCCA ACGGCATTAC CGTTAATGGC GCGGCTTTA
551 AAAATGTCGG TCGGGGCATC TTAACCTACG GTGCGCCCCA AATCGGCAAA
45 601 GACGGTGCAC TGACAGGATT TGATGTGCGT CAAGGCACAT TGACCGTAGG
651 AGCAGCAGGT TGGAATGATA AAGGCGGAGC GCACTACACC GGGGTACTTG
701 CTCGTGCAGT TGCTTTGCAG GGGAAATTAC AGGGTAAAAA CCTGGCGGTT
751 TCTACCGGTC CTCAGAAAGT AGATTACGCC AGCGGCGAAA TCAGTGCAGG
801 TACGGCAGCG GGTACGAAAC CGACTATTGC CCTTGATACT GCCGCACTGG
50 851 GCGGTATGTA CGCCGACAGC ATCACACTGA TTGCCAATGA AAAAGGCGTA
901 GCGTCAAAA ATGCCGGCAC ACTCGAAGCG GCCAAGCAAT TGATTGTGAC
951 TTCGTCAGGC CGCATTGAAA ACAGCGGCCG CATCGCCACC ACTGCCGACG
1001 GCACCGAAGC TTCACCGACT TATCTCTCCA TCGAAACCAC CGAAAAAGGA
1051 GCGGCAGGCA CATTATCTC CAATGGTGGT CGGATCGAGA GCAAAGGCTT
55 1101 ATGGTTATT GAGACGGGAG AAGATATCAG CTTGCGTAAC GGAGCCGTGG
1151 TGCAGAATAA CGGCAGTCGC CCAGCTACCA CGGTATTAAA TGCTGGTCAT
1201 AATTTGGTGA TTGAGAGCAA AACTAATGTG AACAAATGCCA AAGGCCCGCG

```

5

10

15

20

25

30

35

40

45

50

55

60

65

70

1251	TACTCTGTCTG	GCCGACGGCC	GTACCGTCAT	CAAGGAGGCC	AGTATTTCAGA
1301	CTGGCACTAC	CGTATACAGT	TCCAGCAAAG	GCAACGCCGA	ATTAGGCAAT
1351	AACACACGCA	TTACCGGGG	AGATGTTACC	GTATTATCCA	ACGGCACCAT
1401	CAGCAGTTCC	GCCGTAATAG	ATGCCAAAGA	CACCGCACAC	ATCGAAGCAG
1451	GCAAACCGCT	TTCTTTGGAA	GCTTCAACAG	TTACCTCCGA	TATCCGCTTA
1501	AACGGAGGCA	GTATCAAGGG	CGGCAAGCAG	CTTGCTTTAC	TGGCAGACGA
1551	TAACATTACT	GCCAAAACCTA	CCAATCTGAA	TACTCCCAGC	AATCTGTATG
1601	TTCATACAGG	TAAAGATCTG	AATTTGAATG	TTGATAAAGA	TTTGTCTGCC
1651	GCCAGCATCC	ATTTGAAATC	GGATAACGCT	GCCCATATTA	CCGGCACCAG
1701	TAAAACCCCTC	ACTGCCTCAA	AAGACATGGG	TGTGGAGGCA	GGCTCGCTGA
1751	ATGTTACCAA	TACCAATCTG	CGTACCAACT	CGGGTAATCT	GCACATTTCAG
1801	GCAGCCAAAG	GCAATATTCA	GCTTCGCAAT	ACCAAGCTGA	ACGCAGCCAA
1851	GGCTCTCGAA	ACCACGCGAT	TGCAGGGCAA	TATCGTTTCA	GACGGCCTTC
1901	ATGCTGTTTC	TGCAGACGGT	CATGTATCCT	TATTTGGCCAA	CGGTAATGCC
1951	GACTTTTACC	GTCACAATAC	CCTGACAGCC	AAGGCCGATG	TCAATGCAGG
2001	ATCGGTTGGT	AAAGGCCGTC	TGAAAGCAGA	CAATACCAAT	ATCACTTCAT
2051	CTTCAGGAGA	TATTACGTTG	GTTGCCGGCA	ACGGTATTCA	GCTTGGTGAC
2101	GGAAAACAAC	GCAATTCAAT	CAACGGAAAA	CACATCAGCA	TCAAAAACAA
2151	CGGTGGTAAT	CGCGACTTAA	AAAACCTTAA	CGTCCATGCC	AAAAGCGGGG
2201	CATTGAACAT	TCATTCCGAC	CGGGCATTGA	GCATAGAAAA	TACCAAGCTG
2251	GAGTCTACCC	ATAATACGCA	TCTTAATGCA	CAACACGAGC	GGGTAACGCT
2301	CAACCAAGTA	GATGCCTACG	CACACCGTCA	TCTAAGCATT	ACCGGCAGCC
2351	AGATTGCGCA	AAACGACAAA	CTGCCTTCTG	CCAACAAGCT	GGTGGCTAAC
2401	GGTGTATTGG	CACTCAATGC	GCGCTATTCC	CAAAATTGCCG	ACAACACCAC
2451	GCTGAGAGCG	GGTGCAATCA	ACCTTACTGC	CGGTACCGCC	CTAGTCAAGC
2501	GCGGCAACAT	CAATTGGAGT	ACCGTTTCGA	CCAAAACCTT	GGAAAGATAAT
2551	GCCGAATTAA	AACCATTGGC	CGGACGGCTG	AATATTGAAG	CAGGTAGCGG
2601	GCATTAAC	ATCGAACCTG	CCAACCGCAT	CAGTGCGCAT	ACCGACCTGA
2651	GCATCAAAAC	AGGCGGAAAA	TTGCTGTTGT	CTGCAAAAGG	AGGAAATGCA
2701	GGTGCGCCTA	GTGCTCAAGT	TTCTCATTTG	GAAGCAAAAG	GCAATATCCG
2751	TCTGGTTACA	GGAGAAACAG	ATTTAAGAGG	TTCTAAAATT	ACAGCCGGTA
2801	AAAACCTTGGT	TGTCGCCACC	ACCAAAGGCA	AGTTGAATAT	CGAAGCCGTA
2851	AACAACCTCAT	TCAGCAATTA	TTTTCTTACA	CAAAAAGCGG	CTGAACCTCA
2901	CCAAAATCC	AAAGAATTGG	AACAGCAGAT	TGCGCAGTTG	AAAAAAAGCT
2951	CGCCTAAAAG	CAAGCTGATT	CCAACCTGCG	AAGAAGAAGC	CGACCGCTCT
3001	GCTTTCTATA	TTCAAGCCAT	CAACAAGGAA	GTTAAAGGTA	AAAAACCCAA
3051	AGGCAAAGAA	TACCTGCAAG	CCAAGCTTTC	TGCACAAAAT	ATTGACTTGA
3101	TTTCCGCACA	AGGCATCGAA	ATCAGCGGTT	CCGATATTAC	CGCTTCCAAA
3151	AAACTGAACC	TTCAACGCCG	AGGCGTATTG	CCAAAGGCAG	CAGATTTCAG
3201	GGCGGCTGCT	ATTCTGATTG	ACGGCATAAC	CGACCAATAT	GAAATTGGCA
3251	AGCCCACCTA	CAAGAGTCAC	TACGACAAAG	CTGCTCTGAA	CAAGCCTTCA
3301	CGTTTGACCG	GACGTACAGG	GGTAAGTATT	CATGCAGCTG	CGGCACTCGA
3351	TGATGCACGT	ATTATTATCG	GTGCATCCGA	AATCAAAGCT	CCCTCAGGCA
3401	GCATAGACAT	CAAAGCCCAT	AGTGATATTG	TACTGGAGGC	TGGACAAAAC
3451	GATGCCTATA	CCTTCTTAAA	AACCAAAGGT	AAAAGCGGCA	AAATCATCAG
3501	AAAAACCAAG	TTTACCAGCA	CCCGCGACCA	CCTGATTATG	CCAGCCCCCG
3551	TCGAGCTGAC	CGCCAACGGC	ATAACGCTTC	AGGCAGGCGG	CAACATCGAA
3601	GCTAATACCA	CCCGCTTCAA	TGCCCCTGCA	GGTAAAGTTA	CCCTGGTTGC
3651	GGGTGAAGAG	CTGCAACTGC	TGGCAGAAGA	AGGCATCCAC	AAGCACGAGT
3701	TGGATGTCCA	AAAAAGCCGC	CGCTTTATCG	GCATCAAGGT	AGGCAAGAGC
3751	AATTACAGTA	AAACGAACT	GAACGAAACC	AAATTGCCCTG	TCCGCGTCGT
3801	CGCCCAAACT	GCAGCCACCC	GTTCAAGGCTG	GGATACCGTG	CTCGAAGGTA
3851	CCGAATTCAA	AACCACGCTG	GCCGGTGCGG	ACATTACAGC	AGGTGTAGGC
3901	GAAAAAGCCC	GTGCCGATGC	GAAAATTATC	CTCAAAGGCA	TTGTGAACCG
3951	TATCCAGTCG	GAAGAAAAAT	TAGAAACCAA	CTCAACCGTA	TGGCAGAAAC
4001	AGGCCGGACG	CGGCAGCACT	ATCGAAACGC	TGAAACTGCC	CAGCTTCGAA
4051	AGCCCTACTC	CGCCCAAACT	GACCGCCCCC	GGTGGCTATA	TCGTTCGACAT
4101	TCCGAAAGGC	AATTTGAAAA	CCGAAATCGA	AAAGCTGGCC	AAACAGCCCCG
4151	AGTATGCCTA	TCTGAAACAG	CTCCAAGTAG	CGAAAAACGT	CAACTGGAAC
4201	CAGGTGCAAC	TGGCTTACGA	TAAATGGGAC	TATAAGCAGG	AAGGCTTAAAC
4251	CAGAGCCGGT	GCAGCGATTG	TTACCATAAT	CGTAACCGCA	CTGACTTATG
4301	GATACGGCGC	AACCGCAGCG	GGCGGTGTAG	CCGCTTCAGG	AAGTAGTACA
4351	GCCGCAGCTG	COGGAACAGC	CGCCACAACG	ACAGCAGCAG	CTACTACCGT
4401	TTCTACAGCG	ACTGCCATGC	AAACCGCTGC	TTTAGCCCTC	TTGTATAGCC
4451	AAGCAGCTGT	ATCCATCATC	AATAATAAAG	GTGATGTCGG	CAAAGCGTTG
4501	AAAGATCTCG	GCACCACTGA	TACGGTCAAG	CAGATTGTCA	CTTCTGCCCT
4551	GACGGCGGGT	GCATTAAATC	AGATGGGCGC	AGATATTGCC	CAATTGAACA
4601	GCAAGGTAAG	AACCGAACTG	TTCAGCAGTA	CGGGCAATCA	AACATTGGCC
4651	AACCTTGGAG	GCAGACTGCG	TACCAATCTC	AGTAATGCAG	GTATCTCAGC
4701	TGGTATCAAT	ACCGCCGCTG	ACGGCGGCAG	CCTGAAAGAC	AACTTAGGCA
4751	ATGCCGCATT	AGGAGCATTG	GTTAATAGCT	TCCAAGGAGA	AGCCGCGCAG
4801	AAAATCAAAA	CAACCTTCAG	CGACGATTAT	GTTGCCAAAC	AGTTGCCCAA

5

10

15

20

```

4851 CGCTTTGGCT GGGTGTGTTA GCGGATTGGT ACAAGGAAAA TGTAAGACG
4901 GGGCAATTGG CGCAGCAGTT GGGGAAATCG TAGCCGACTC CATGCTTGGC
4951 GGCAGAAACC CTGCTACACT CAGCGATGCG GAAAAGCATA AGGTATACAG
5001 TTACTIONGAG ATTATTGCCG GCAGCGTGGC GGCACCAAC GCGCGCGATG
5051 TGAATACTGC GCGCAATGCG GCTGAGGTGG CCGTAGTGAA TAATGCTTTG
5101 AATTTTGACA GTACCCCTAC CAATCGGAAA AAGCATCAAC CGCAGAAGCC
5151 CGACAAAACC GCACTGGAAA AAATTATCCA AGGTATTATG CCTGCACATG
5201 CAGCAGGTGC GATGACTAAT CCGCAGGATA AGGATGCTGC CATTGTGATA
5251 AGCAATATCC GTAATGGCAT CACAGGCCCG ATTGTGATTA CCAGCTATGG
5301 GGTTTATGCT GCAGGTGGGA CAGCTCCGCT GATCGGTACA GCGGGTAAAT
5351 TAGCTATCAG CACCTGCATG GCTAATCCTT CTGGTTGTAC TGTATGGTC
5401 ACTCAGGCTG CCGAAGCGGG CGCGGGAATC GCCACGGGTG CCGTACGGT
5451 AGGCAACGCT TGGGAAGCGC CTGTGGGGGC GTTGTGAAA GCGAAGGCGG
5501 CCAAGCAGGC TATACCAACC CAGACAGTTA AAGAAGTTGA TGGCTTACTA
5551 CAAGAATCAA AAAATATAGG TGCTGTAAAT ACACGAATTA ATATAGCGAA
5601 TAGTACTACT CGATATACAC CAATGAGACA AACGGGACAA CCGGTATCTG
5651 CTGGCTTTGA GCATGTTCTT GAGGGGCACT TCCATAGGCC TATTGCGAAT
5701 AACCCTTCAG TTTTACCAT CTCCCAAAT GAATTGAAGG TTATACTTCA
5751 AAGTAATAAA GTAGTTTCTT CTCCCGTATC GATGACTCCT GATGGCCAA
5801 ATATCGGAC TGTCGATGTA GGAAAAGTTA TTGGTACTAC TTCTATTAAA
5851 GAAGGTGGAC AACCCACAAC TACAATTAAA GTATTACAG ATAAGTCAGG
5901 AAATTTGATT ACTACATACC CAGTAAAAGG AAATAA

```

This corresponds to the amino acid sequence <SEQ ID 60; ORF114-1>:

25

30

35

40

45

50

55

60

```

1 MNKGLHRIIF SKKHSTMVAV AETANSQKG KQAGSSVSVS LKTSGLDLCGK
51 LKTTLLKTLVC SLVSLSMVLP AHAQITTDKS APKNQOVVIL KTNTGAPLVN
101 IQTPNGRGLS HNRYTQFDVD NKGAVLNDR NNNPFVVKGS AQLILNEVRG
151 TASKLNGIVT VGGQKADVII ANPNGITVNG GGFKNVGRGI LTTGAPQIGK
201 DGALTGFDVR QGTLTVGAAG WNDKGGADYT GVLARAVALQ GKLOGKNLAV
251 STGPQKVDYA SGEISAGTAA GTKPTIALDT AALGGMYADS ITLIANEKGV
301 GVKNAGTLEA AKQLIVTSSG RIENSGRIAT TADGTEASPT YLSIETTEKG
351 AAGTFISNGG RIESKGLLVI ETGEDISLRN GAVVQNNNGSR PATTVLNAGH
401 NLVIESKTNV NNAKGPATLS ADGRTVIKEA SIQTGTTVYS SSKGNAELGN
451 NTRITGADVT VLSNGTISSS AVIDAKDTAH IEAGKPLSLE ASTVTDIRL
501 NGGSIKGGKQ LALLADDNIT AKTTLNTPG NLYVHTGKDL NLNVDKDLSA
551 ASIHLKSDNA AHITGTSKTL TASKDMGVEA GSLNVTNTNL RTNSGNLHIQ
601 AAKNGIQLRN TKLNAAKALE TTALQGNIVS DGLHAVSADG HVSLLANGNA
651 DFTGHNTLTA KADVNAGSVG KGRLKADNTN ITSSSGDITL VAGNGIQLGD
701 GKQRNSINGK HISIKNNGGN ADLKNLNVHA KSGALNIHSD RALSIENTKL
751 ESTHNTLNA QHERVTLNQV DAYAHRHLSI TGSQIWQNDK LPSANKLVAN
801 GVLALNARYS QIADNTTLRA GAINLTAGTA LVKRGNNINWS TVSTKTLEDN
851 AELKPLAGRL NIEAGSGTTL IEPANRISAH TDLSTKTGGK LLSAKGGNA
901 GAPSAQVSSL EAKGNIRLVT GETDLRSGKI TAGKNLVVAT TKGKLNIEAV
951 NNSFSNYFPT QKAAELNQKS KELEQQIAQL KKSSPKSKLI PTLQEERDRL
1001 AFYIQAINKE VKGKKPKGKE YLQAKLSAQN IDLISAQGIE ISGSDITASK
1051 KLNLAHAGVL PKAADSEAAA ILIDGITDQY EIGKPTYKSH YDKAALNPKS
1101 RLTGRTGVSI HAAAALDDAR IIGASEIKA PSGSIDIKAH SDIVLEAGQN
1151 DAYTFLKTKG KSGKIIRKTK FTSTRDHLIM PAPVELTANG ITLQAGGNIE
1201 ANTTFRNAPA GKVTTLVAGEE LQLLAEEGIH KHELDVQKSR RFIGIKVGKS
1251 NYSKNELNET KLPVRVVAQT AATRSWDTV LEGTEFKTTL AGADIQAGVG
1301 EKARADAKII LKGIVNRIQS EEKLETNSTV WQKQAGRGST IETLKLPSFE
1351 SPTPPKLTA GGYIVDIPKG NLKTEIEKLA KQPEYAYLKQ LQVAKNVNWN
1401 QVQLAYDKWD YKQEGLTRAG AAIVTIIVTA LTYGYGATAA GGVAASGSST
1451 AAAAGTAATT TAAATTVSTA TAMQTAALAS LYSQAASVSI NNKGDVKGAL
1501 KDLGTSDTV KQIVTSALTAG ALNQMAGADIA QLNSKVRTEL FSSTGNQITIA
1551 NLGGRLATNL SNAGISAGIN TAVNGGSLKD NLGNAALGAL VNSFQGEAAS
1601 KIKTTFSDDY VAKQFAHALA GCVSGLVQGK CKDGAIGAAV GEIVADSMILG
1651 GRNPATLSDA EKHKVISYSK IIAGSVAALN GGDVNTAANA AEAUVNNAL
1701 NFDSTPTNAK KHQPQKPKDKT ALEKIIQGIM PAHAAGAMTN PQDKDAIWI
1751 SNIRNGITGP IVITSYGUYA AGWTAPLIGT AGKLAISTCM ANPSGCTVMV
1801 TQAAEAGAGI ATGAVTVGNA WEAPVGALSK AKAQAIPPT QTVKLEDGLL
1851 QESKNIGAVN TRINIANSTT RYTPMRQTGQ PVSAGFEHVL EGHFHRPIAN
1901 NRSVFTISPN ELKVILQSNK VVSSPVSMTP DGQYMRTVDV GKVIGTTSIK
1951 EGGQPTTTIK VFTDKSGNLI TTPVVKGN*

```

Computer analysis of this amino acid sequence predicts a transmembrane region and also gives the following results:

Homology with a predicted ORF from *N.meningitidis* (strain A)

ORF114 shows 91.9% identity over a 284aa overlap with an ORF (ORF114a) from strain A of *N. meningitidis*:

5	orf114.pep		10	20	30	40
			AVAETANSQKGKQAGSSVSVSLKTS	SGDLCGKLKTTL	KLTVLC	
	orf114a	MNKG	LHRIIFSKKHST	MVA	VAETANSQKGKQAGSSVSVSLKTS	SGDLCGKLKTTLKLTVLC
		10	20	30	40	50
10	orf114.pep	50	60	70	80	90
		SLVSLSMVLP	HAQA	ITTDKSAPKNQ	QVVILKTN	TGAPLVNIQT
	orf114a	SLVSLSMXXXXXX	QITTDKSAPKNQ	QVVILKTN	TGAPLVNIQT	PNGRGLSHNRYTQ
		70	80	90	100	110
15	orf114.pep	110	120	130	140	150
		NKGAVLNDRNN	PNFVVK	GSAQLILNE	VRGTASKL	NGI
	orf114a	NKGAVLNDRNN	PNFVVK	GSAQLILNE	VRGTASKL	NGI
		130	140	150	160	170
20	orf114.pep	170	180	190	200	210
		GGFKNVGRG	ILTTGAPQ	IGKD	GALTGF	DVVK
	orf114a	GGFKNVGRG	ILTTGAPQ	IGKD	GALTGF	DVVK
		190	200	210	220	230
25	orf114.pep	230	240	250	260	270
		GKXXGKXL	AVSTG	PQKVDY	ASGEIS	AGTAAG
	orf114a	GKXXGKXL	AVSTG	PQKVDY	ASGEIS	AGTAAG
		250	260	270	280	290
30	orf114.pep	310	320	330	340	350
		GVX				
	orf114a	GVKNAG	TLAAKQL	LIVTSS	GRIENSG	RIATTAD
		310	320	330	340	350

The complete length ORF114a nucleotide sequence <SEQ ID 61> is:

40	1	ATGAATAAAG	GTTTACATCG	CATTATCTTT	AGTAAAAAGC	ACAGCACCAT
	51	GGTTGCAGTA	GCCGAAACTG	CCAACAGCCA	GGGCAAAGGT	AAACAGGCAG
	101	GCAGTTCGGT	TTCTGTTTCA	CTGAAAACTT	CAGGCGACCT	TTGCGGCAAA
	151	CTCAAAACCA	CCCTTAAAC	CTTGGTCTGC	TCTTTGGTTT	CCCTGAGTAT
	201	GACATTNCNN	NNCNTNCCC	AAATTACCAC	CGACAAATCA	GCACCTAAAA
45	251	ACCANCAGGT	CGTTATCCTT	AAAACCAACA	CTGGTGCCCC	CTTGGTGAAT
	301	ATCCAAACTC	CGAATGGACG	CGGATTGAGC	CACAACCGCT	ATACGCAGTT
	351	TGATGTTGAC	AACAAAGGGG	CAGTGTAA	CAACGACCGT	AACAATAATC
	401	CGTTTCTGGT	CAAAGGCAGT	GCGCAATTGA	TTTGAACGA	GGTACGCGGT
	451	ACGGCTAGCA	AACTCAACGG	CATCGTTACC	GTAGGCGGTC	AAAAGGCCGA
50	501	CGTGATTATT	GCCAACCCCA	ACGGCATTAC	CGTTAATGGC	GGCGGCTTTA
	551	AAAATGTCGG	TCGGGGCATC	TTAACTATCG	GTGCGCCCCA	AATCGGCAAA
	601	GACGGTGCAC	TGACAGGATT	TGATGTGCGT	CAAGGCACAT	TGACCGTAGG
	651	AGCAGCAGGT	TGGAATGATA	AAGGCGGAGC	CGACTACACC	GGGGTACTTG
	701	CTCGTGCAGT	TGCTTTGCAG	GGGAAATTAC	AGGGTAAAAA	CCTGGCGGTT
55	751	TCTACCGGTC	CTCAGAAAGT	AGATTACGCC	AGCGGCGAAA	TCAGTGCAGG
	801	TACGGCAGCG	GGTACGAAAC	CGACTATTGC	CCTTGATACT	GCCGCACTGG
	851	GCGGTATGTA	CGCCGACAGC	ATCACA	CTGA	TTGCCANTGA
	901	GGCGTCAAAA	ATGCGGGCAC	ACTCGAAGCG	GCCAAGCAAT	TGATTGTGAC
	951	TTCTGCAGGC	CGCATTGAAA	ACAGCGGCCG	CATCGCCACC	ACTGCCGACG
60	1001	GCACCGAAGC	TTCAACGACT	TATCTNNCNA	TCGAAACCAC	CGAAAAAGGA
	1051	GCNNCAGGCA	CATTTATCTC	CAATGGTGGT	CGGATCGAGA	GCAAAGGCTT
	1101	ATTGGTTATT	GAGACGGGAG	AAGATATCAN	CTTGCGTAAC	GGAGCCGTGG
	1151	TGCAGAATAA	CGGCAGTCGC	CCAGCTACCA	CGGTATTAAA	TGCTGGTCAT
	1201	AATTTGGTGA	TTGAGAGTAA	AACAAATGTG	AACAATGCCA	AAGGCTCGNC

5

10

15

20

25

30

35

40

45

50

55

60

65

```

1251 TAATCTGTCG GCCGGCGGTC GTACTACGAT CAATGATGCT ACTATTCAAG
1301 CGGGCAGTTC CGTGACAGC TCCACCAAAG GCGATACTGA NTTGGGTGAA
1351 AATACCCGTA TTATTGCTGA AAACGTAACC GTATTATCTA ACGGTAGTAT
1401 TGGCAGTGCT GCTGTAATTG AGGCTAAAGA CACTGCACAC ATTGAATCGG
1451 GCAAACCGCT TTCTTTAGAA ACCTCGACCG TTGCCTCCAA CATCCGTTTG
1501 AACACCGTA ACATTAAAGG CGGAAAGCAG CTTGCTTTAC TGGCAGACGA
1551 TAACATTACT GCCAAACTA CCAATCTGAA TACTCCCGGC AATCTGTATG
1601 TTCATACAGG TAAAGATCTG AATTGGAATG TTGATAAAGA TTTGTCTGCC
1651 GCCAGCATCC ATTTGAAATC GGATAACGCT GCCCATATTA CCGGCACCAG
1701 TAAAACCCTC ACTGCCTCAA AAGACATGGG TGTGGAGGCA GGCTTGCTGA
1751 ATGTTACCAA TACCAATCTG CGTACCAACT CGGGTAATCT GCACATTAG
1801 GCAGCCAAAG GCAATATTCA GCTTCGCAAT ACCAAGCTGA ACGCAGCCAA
1851 GGCTCTCGAA ACCACCGCAT TGCAGGGCAA TATCGTTTCA GACGGCCTTC
1901 ATGCTGTTTC TGCAGACGGT CATGTATCCT TATTGGCCAA CGGTAATGCC
1951 GACTTTACCG GTCACAATAC CCTGACAGCC AAGGCCGATG TCNATGCAGG
2001 ATCGGTTGGT AAAGGCCGTC TGAAAGCAGA CAATACCAAT ATCACTTTCAT
2051 CTTACAGAGA TATTACGTTG GTTGCCGNNN NCGGTATTCA CGTTGGTGAC
2101 GGAAAACAAC GCAATTCAT CAACGGAAAA CACATCAGCA TCAAAAACAA
2151 CGGTGGTAAT GCCGACTTAA AAAACCTTAA CGTCCATGCC AAAAGCGGGG
2201 CATTGAACAT TCATTCCGAC CGGGCATTGA GCATAGAAAA TACNAAGCTG
2251 GAGTCTACCC ATAATACGCA TCTTAATGCA CAACACGAGC GGGTAACGCT
2301 CAACCAAGTA GATGCCTACG CACACCGTCA TCTAAGCATT ANCGGCAGCC
2351 AGATTTGGCA AAACGACAAA CTGCCTTCTG CCAACAAGCT GGTGGCTAAC
2401 GGTGTATTGG CANTCAATGC GCGCTATTCC CAAATTGCCG ACAACACCAC
2451 GCTGAGAGCG GGTGCAATCA ACCTTACTGC CGGTACCGCC CTAGTACAGC
2501 GCGGCAACAT CAATTGGAGT ACCGTTTCGA CCAAGACTTT GGAAGATAAT
2551 GCCGAATTAA AACCATTGGC CGGACGGCTG AATATTGAAG CAGGTAGCGG
2601 CACATTAACC ATCGAACCTG CCAACCGCAT CAGTGCGCAT ACCGACCTGA
2651 GCATCAAAAC AGGCGGAAAA TTGCTGTTGT CTGCAAAAGG AGGAAATGCA
2701 GGTGCGCNTA GTGCTCAAGT TTCCTCAT TG AAGCAAAAG GCAATATCCG
2751 TCTGGTTACA GGAGNAACAG ATTTAAGAGG TTCTAAATTT ACAGCCGGTA
2801 AAAAGTTGGT TGTGCGCACC ACCAAAGGCA AGTTGAATAT CGAAGCCGTA
2851 AACAACTCAT TCAGCAATTA TTTTCNTACA CAAAAGNGN NNGNNCTCAA
2901 CAAAAAATCC AAAGAAATGG AACAGCAGAT TGCGCAGTTG AAAAAAGCT
2951 CGCNTAAAG CAAGCTGATT CCAACCCCTG AAGAAGAACG CGACCGTCTC
3001 GCTTCTATA TTCAAGCCAT CAACAAGGAA GTTAAAGGTA AAAAAACCA
3051 AGGCAAGAA TACCTGCAAG CCAAGCTTTC TGCACAAAAT ATTGACTTGA
3101 TTTCCGCACA AGGCATCGAA ATCAGCGGTT CCGATATTAC CGCTTCCAAA
3151 AAAGTGAACC TTCACGCCGC AGGCGTATTG CCAAAGGCAG CAGATTGAGA
3201 GCGGCTGCT ATTCTGATTG ACGGCATAAC CGACCAATAT GAAATTGGCA
3251 AGCCACCTA CAAGAGTCAC TACGACAAAG CTGCTCTGAA CAAGCCTTCA
3301 CGTTTGACCG GACGTACGGG GGTAAGTATT CATGCAGCTG CGGCACTCGA
3351 TGATGCACGT ATTATTATCG GTGCATCCGA AATCAAAGCT CCCTCAGGCA
3401 CATAAGACAT CAAAGCCCAT AGTGATATTG TACTGGAGGC TGGACAAAC
3451 GATGCCTATA CCTTCTTANA AACCAAAGGT AAAAGCGGCA NAATNATCAG
3501 AAAAACNAAG TTTACAGCA CCNGCGANCA CCTGATTATG CCAGCCCNCG
3551 TCGAGCTGAC CGCCAACGGT ATCAGCTTTC AGGCAAGCGG CAACATCGAA
3601 TCAATACCA CCGCTTCAA TGCCCTGCA GGTAAGTTA CCCTGGTTGC
3651 GGGTGAANAG NTGCAACTGC TGGCAGAAGA AGGCATCCAC AAGCAGGAGT
3701 TGGATGTCCA AAAAGCCGC CGCTTTATCG GCATCAAGGT AGGTNAGAGC
3751 AATTACAGTA AAAACGAACT GAACGAAACC AAATTGCCTG TCGCGCTCGT
3801 CGCCCAANT GCAGCCACCC GTTCAGGCTG GGATACCGTG CTCGAGGTA
3851 CCGAATTCAA AACCACGCTG GCCGGTGCCG ACATTAGGC AGGTGTANGC
3901 GAAAAAGCCC GTGTCGATGC GAAAAATATC CTCAAAGGCA TTGTGAACCG
3951 TATCCAGTCG GAAGAAAAAT TAGAAACCAA CTCACCGTA TGGCAGAAAC
4001 AGGCCGACG CGGCAGCACT ATCGAAACGC TAAAGCTGCC CAGCTTCGAA
4051 AGCCCTACTC CGCCCAATTT GTCCGCACCC GGCGGNTATA TCGTCGACAT
4101 TCCGAAAGGC AATCTGAAAA CCGAAATCGA AAAGCTGTCC AAACAGCCCG
4151 AGTATGCCTA TCTGAAACAG CTCCAAGTAG CGAAAAACAT CAACTGGAAT
4201 CAGGTGCAGC TTGCTTACGA CAGATGGGAC TACAAACAGG AGGGCTTAAC
4251 CGAAGCAGGT GCGGCGATTA TCGCACTGGC CGTTACCGTG GTCACCTCAG
4301 GCGCAGGAAC CGGAGCCGTA TTGGGATTAA ACGGTGCGNC CGCCGCCGCA
4351 ACCGATGCAG CATTCGCCTC TTTGGCCAGC CAGGCTTCCG TATCGTTTAT
4401 CAACAACAAA GCGGATGTCG GCAAAACCCT GAAAGAGCTG GGCAGAGCA
4451 GCACGGTGAA AAATCTGGTG GTTGCCGCCG CTACCGCAGG CGTAGCCGAC
4501 AAAATCGGCG CTTCCGGCACT GANCAATGTC AGCGATAAGC AGTGGATCAA
4551 CAACCTGACC GTCAACCTAG CCAATGNCGG GCAGTGCCGC ACTGAttaa

```

This encodes a protein having amino acid sequence <SEQ ID 62>:

1 MNKGLHRIIF SKKHSTMVAV AETANSQGKG KQAGSSVSVS LKTSGLCGK

5 51 LKTTTLKTLVC SLVSLSMXXX XXXQITTDKS APKNXQVVIL KTNTGAPLVN
 101 IQTPNGRGLS HNRYTQFDVD NKGAVLNNDNR NNNPFLVKGS AQLILNEVRG
 151 TASKLNGIVT VGGQKADVII ANPNGITVNG GGFKNVGRGI LTIGAPQIGK
 201 DGALTGFVDVR QGTLTVGAAG WNDKGGADYT GVLARAVALQ GKLGKKNLAV
 251 STGPQKVDYA SGEISAGTAA GTKPTIALDT AALGGMYSADS ITLIAXEKGV
 301 GVKNAGTLEA AKQLIVTSSG RIENSGRIAT TADGTEASPT YLXIETTEKG
 351 AXGTFISNGG RIESKGLLVI ETGEDIXLRN GAVVQNNNGSR PATTVLNAGH
 401 NLVIESKTNV NNAKGSXNLS AGGRTTINDA TIQAGSSVYS STKGDTXLGE
 451 NTRIIAENVV VLSNGSIGSA AVIEAKDTAH IESGKPLSLE TSTVASNIRL
 10 501 NNGNIKGGKQ LALLADDNIT AKTTNLNTPG NLYVHTGKDL NLNVDKDLSA
 551 ASIHLSDNA AHITGTSKTL TASKDMGVEA GLLNVTNTNL RTNSGNLHIQ
 601 AAKGNIQLRN TKLNAAKALE TTALQGNIVS DGLHAVSADG HVSLLANGNA
 651 DFTGHNTLTA KADVXAGSVG KGRKADNTN ITSSSGDITL VAXXGQQLGD
 701 GKQRNSINGK HISIKNNGGN ADLKNLNVHA KSGALNIHSD RALSIENTKL
 15 751 ESTHNTLNA QHERVTLNQV DAYAHRHLSI XGSQIWQNDK LPSANKLVAN
 801 GVLAXNARYS QIADNTTLRA GAINLTAGTA LVKRGNNINWS TVSTKTLNEDN
 851 AELKPLAGRL NIEAGSGTIL IEPANRISAH TDLSIKTGGK LLSAKGGNA
 901 GAXSAQVSSL EAKGNIRLVT GXTDLRSGSKI TAGKNLVVAT TKGKLNIEAV
 951 NNSFSNYFXT QKXXXLNQKS KELEQQIAQL KKSSXKSKLI PTLQEERDRL
 20 1001 AFYIQAINKV VKGKKPKGKE YLQAKLSAQN IDLISAQIE ISGSDITASK
 1051 KLNLAAGVL PKAADSEAAA ILIDGITDQY EIGKPTYKSH YDKAALNKPS
 1101 RLTRGTGVSII HAAALDDAR IIIGASEIKA PSGSIDIKAH SDIVLEAGQN
 1151 DAYTFLXTKG KSGXXIRKTK FTSTXXHLIM PAPVELTANG ITLQAGGNIE
 25 1201 ANTTFRNAPA QKVTLVAGEX XQLLAEEGIH KHELDVQKSR RFIGIKVXGS
 1251 NYSKNELNET KLPVRVVAQX AATRSWDTV LEGTEFKTTL AGADIQAGVX
 1301 EKARVDAKII LKGIVNRIQS EEKLETNSTV WQKQAGRGST IETLKLPSFE
 1351 SPTPPKLSAP GGYIVDIPKG NLKTEIEKLS KQPEYAYLKO LQVAKNNINWN
 1401 QVQLAYDRWD YKQEGLTEAG AAIILAVTV VTSAGTGAV LGLNGAXAAA
 30 1451 TDAAFASLAS QASVSFINNK GGVGKTLKEL GRSSTVKNLV VAAATAGVAD
 1501 KIGASALXNV SDKQWINNLT VNLANXGQCR TD*

ORF114-1 and ORF114a show 89.8% identity in 1564 aa overlap

35 orf114a.pep MNKGLHRIIFSKKHSTMVAVAETANSQKGKQAGSSVSLSLKTSGDLCGKLTTLKTLVC
 orf114-1 MNKGLHRIIFSKKHSTMVAVAETANSQKGKQAGSSVSLSLKTSGDLCGKLTTLKTLVC
 40 orf114a.pep SLVSLSMXXXXXXQITTDKSAPKNXQVVILKTNTGAPLVNIQTPNGRGLSHNRYTQFDVD
 orf114-1 SLVSLSMVLPAAHAQITTDKSAPKNQVVILKTNTGAPLVNIQTPNGRGLSHNRYTQFDVD
 45 orf114a.pep NKGAVLNNDNRNNNPFLVKGSAQLILNEVRGTASKLNGIVTVGGQKADVIIANPNGITVNG
 orf114-1 NKGAVLNNDNRNNNPVVKGSAQLILNEVRGTASKLNGIVTVGGQKADVIIANPNGITVNG
 50 orf114a.pep GGFKNVGRGILTIGAPQIGKDGALTGFVDVRQGTLTVGAAGWNDKGGADYTGVLARAVALQ
 orf114-1 GGFKNVGRGILTTGAPQIGKDGALTGFVDVRQGTLTVGAAGWNDKGGADYTGVLARAVALQ
 55 orf114a.pep GKLGKKNLAVSTGPQKVDYASGEISAGTAAGTKPTIALDTAALGGMYSADITLIAXEKGV
 orf114-1 GKLGKKNLAVSTGPQKVDYASGEISAGTAAGTKPTIALDTAALGGMYSADITLIANEKGV
 60 orf114a.pep GVKNAGTLEAAKQLIVTSSGRIENSGRIATTADGTEASPTYLXIETTEKGAXGTFISNGG
 orf114-1 GVKNAGTLEAAKQLIVTSSGRIENSGRIATTADGTEASPTYLSIETTEKGAGTFISNGG
 65 orf114a.pep RIESKGLLVIETGEDIXLRNGAVVQNNNGSRPATTVLNAGHNLVIESKTNVNNAKGSXNLS
 orf114-1 RIESKGLLVIETGEDISLRNGAVVQNNNGSRPATTVLNAGHNLVIESKTNVNNAKGPATLS
 70 orf114a.pep AGGRTTINDATIQAGSSVYSSTKGDTXLGENTRIIAENVTVLSNGSIGSAAVIEAKDTAH
 orf114-1 ADGRTVIKEASIQGTGTVYSSSKGNAELGNNTTRITGADVTVLSNGTISSSAVIDAKDTAH
 75 orf114a.pep IESGKPLSLETSTVASNIRLNNNGNIKGGKQLALLADDNITAKTTNLNTPGNLYVHTGKDL
 orf114-1 IEAGKPLSLEASTVTSDIRLNGGSIKGGKQLALLADDNITAKTTNLNTPGNLYVHTGKDL
 80 orf114a.pep NLNVDKDLSAASIHLSDNAAHITGTSKTLTASKDMGVEAGLLNVTNTNLRTNSGNLHIQ
 orf114-1 NLNVDKDLSAASIHLSDNAAHITGTSKTLTASKDMGVEAGLLNVTNTNLRTNSGNLHIQ

	orf114-1	NLNVDKOLSAASIHLKSDNAAHITGTSKTLTASKDMGVEAGSLNVTNTNLRNNSGNLHIQ	
	orf114a.pep	AAKGNIQLRNTKLNAKALETTALQGNIVSDGLHAVSADGHVSLLANGNADFTGHNTLTA	
5	orf114-1	AAKGNIQLRNTKLNAKALETTALQGNIVSDGLHAVSADGHVSLLANGNADFTGHNTLTA	
	orf114a.pep	KADVXAGSVGKGRLLKADNTNITSSSGDITLVAXXGQIQLGDGKQRNSINGKHSIKNNGGN	
10	orf114-1	KADVXAGSVGKGRLLKADNTNITSSSGDITLVAGNGIQLGDGKQRNSINGKHSIKNNGGN	
	orf114a.pep	ADLKNLNVHAKSGALNIHSDRALSIENTKLESTHNTLNAQHERVTLNQVDAYAHRHLSI	
	orf114-1	ADLKNLNVHAKSGALNIHSDRALSIENTKLESTHNTLNAQHERVTLNQVDAYAHRHLSI	
15	orf114a.pep	XGSQIWQNDKLP SANKLVANGVLAXNARYS QIADNTTLRAGAINLTAGTALVKRGNINWS	
	orf114-1	TGSQIWQNDKLP SANKLVANGVLALNARYS QIADNTTLRAGAINLTAGTALVKRGNINWS	
20	orf114a.pep	TVSTKTLEDNAELKPLAGRLNIEAGSGTLTIEPANRISAHTDLSIKTGGKLLLSAKGGNA	
	orf114-1	TVSTKTLEDNAELKPLAGRLNIEAGSGTLTIEPANRISAHTDLSIKTGGKLLLSAKGGNA	
	orf114a.pep	GAXSAQVSSLEAKGNIRLVGTGXTDLRGSKITAGKNLVVATTGKGLNIEAVNNSFSNYFXT	
25	orf114-1	GAPSAQVSSLEAKGNIRLVGTGETDLRGSKITAGKNLVVATTGKGLNIEAVNNSFSNYFFT	
	orf114a.pep	QKXXLNQKSKELEQQIAQLKKSSXKSKLIPTLQEERDRLAFYIQAINEVKGKPKGKE	
30	orf114-1	QKAAELNQKSKELEQQIAQLKKSSPKSKLIPTLQEERDRLAFYIQAINEVKGKPKGKE	
	orf114a.pep	YLQAKLSAQNIDLISAQGIEISGSDITASKKLNLAAGVLPKAADSEAAAAILIDGITDQY	
	orf114-1	YLQAKLSAQNIDLISAQGIEISGSDITASKKLNLAAGVLPKAADSEAAAAILIDGITDQY	
35	orf114a.pep	EIGKPTYKSHYDKAALNKPSRLTGRTGVSIIHAAAALDDARIIGASEIKAPSGSIDIKAH	
	orf114-1	EIGKPTYKSHYDKAALNKPSRLTGRTGVSIIHAAAALDDARIIGASEIKAPSGSIDIKAH	
40	orf114a.pep	SDIVLEAGQNDAYTFLXTKGKSGXXIRKTKFTSTXXHLIMPAPVELTANGITLQAGGNIE	
	orf114-1	SDIVLEAGQNDAYTFLXTKGKSGKIIKTKFTSTRDHLIMPAPVELTANGITLQAGGNIE	
	orf114a.pep	ANTTRFNAPAGKVTLVAGEXXQLLAEEGIHKHELDVQKSRRFIGIKVGXSNYSKNELNET	
45	orf114-1	ANTTRFNAPAGKVTLVAGEELQLLAEEGIHKHELDVQKSRRFIGIKVGXSNYSKNELNET	
	orf114a.pep	KLPVRVVAQXAATRSWDTVLEGTEFKTTLGADIQAGVXEKARVDAKIIKLGIVNRIQS	
50	orf114-1	KLPVRVVAQTAATRSWDTVLEGTEFKTTLGADIQAGVGEKARADAKIIKLGIVNRIQS	
	orf114a.pep	EEKLETNSTVWQKQAGRGSTIETLKLPSFESPTPPKLSAPGGYIVDIPKGNLKTEIEKLS	
	orf114-1	EEKLETNSTVWQKQAGRGSTIETLKLPSFESPTPPKLTAPGGYIVDIPKGNLKTEIEKLA	
55	orf114a.pep	KQPEYAYLKQLQVAKNINWNQVQLAYDRWDYKQEGLTEAGAAIIALAVTVVTSAGTGAV	
	orf114-1	KQPEYAYLKQLQVAKNINWNQVQLAYDKWDYKQEGLTRAGAAIVTIIVTALTYGYGATAA	
60	orf114a.pep	LGLNGA-----XAAATD-----AAFASLASQASVSFINNKGDVGKTL 1477	
	orf114-1	GGVAASGSSTAAAAGTAATTTAAATTVSTATAMQTAALASLYSQAASVIINNKGDVGKAL 1500	
	orf114a.pep	KELGRSSTVKNLVVAATAGVADKIGA-----SALXNVSDKQWINNL-----TVNL 1523	
65	orf114-1	KDLGTSDTVKQIVTSALTAGALNQMGAIAQLNSKVRTELFSSSTGNQTIANLGGRLATNL 1560	
	orf114a.pep	ANXGQCRTDX	
70	orf114-1	SNAGISAGINTAVN...	

Homology with pspA putative secreted protein of *N.meningitidis* (accession number AF030941)

ORF114 and pspA protein show 36% aa identity in 302aa overlap:

```

5      Orf114: 1  AVAETANSQGKGKQAGSSVSLSL-----KTS GDXXXXXXXXXXXXXXXXXXXXXXPAHAQ 56
      pspA: 19  AVAENVHRDGKSMQDSEASVRVTGAASVSSARAAGFRMAAFSVMLALGVAAFPAPAS 78
      Orf114: 57  -ITTDKSAPKNQQVILKTNTGAPLVNIQTPNGRGLSHNRXYAFDVDNKGAVLNNDRNN- 114
      pspA: 79  GIIADKSAPKNQQAVILQTANGLPQVNIQTPSSQGVSVNRFKQFDVDEKGVILNNSRSNT 138
10     Orf114: 115 -----NPFVVGKSAQLILNEV-RGTASKLNGIVTVGGQKADVIIANPNGITVNGG 163
      pspA: 139 QTQLGGWIIQGNPHLARGEARVIVNQIDSSNPSSLNGYIEVGGKRAEVVVPNGSIRVNGG 198
15     Orf114: 164 GFKNVGRGILTTGAPQIGKDGALTGFDDVVKAHWTVXAAGWNDKGGAXYTGVLARAVALQG 223
      pspA: 199 GLINAASVTLTSGVPVL-NNGNLTGFDVSSGKVVIGGKGL-DTSDADYTRILSRAAEINA 256
20     Orf114: 224 KXXGKXLAVSTGPKQVDYASGEISAGTAAGTK----PTIALDTAALGGMYADSITLIANE 279
      pspA: 257 GVWGDVKVSVSGKNKLDGSLAKTASAPSSSDSVTPTVAIDTATLGGMYADKITLISTD 316
      Orf114: 280 KG 281
      pspA: 317 NG 318
25

```

ORF114a is also homologous to pspA:

```

30     gi|2623258 (AF030941) putative secreted protein [Neisseria meningitidis] Length
      = 2273
      Score = 261 bits (659), Expect = 3e-68
      Identities = 203/663 (30%), Positives = 314/663 (46%), Gaps = 76/663 (11%)
35     Query: 1  MNKGLHRIIFSKKHSTMVAVAETANSQGKGKQAGSSVSLSL-----TSGDXXXXXXXXXX 55
      Sbjct: 1  MNK +++IF+KK S M+AVAE + GK Q + SV + +S
      Query: 56  XXXXXXXXXXXXXXXXXXXXQITTDKSAPKNXQVILKTNTGAPLVNIQTPNGRGLSHNRXY 115
      Sbjct: 61  FSVMLALGVAAFPAPASGIIADKSAPKNQQAVILQTANGLPQVNIQTPSSQGVSVNRFK 120
40     Query: 116 QFDVDNKGAVLNNDRNN-----NPFVVGKSAQLILNEV-RGTASKLNGIVTVGG 163
      Sbjct: 121 QFDVDEKGVILNNSRSNTQTQLGGWIIQGNPHLARGEARVIVNQIDSSNPSSLNGYIEVGG 180
45     Query: 164 QKADVIIANPNGITVNGGGGFKNVGRGILTTGAPQIGKDGALTGFDDVQGTLTGVAAGWND 223
      Sbjct: 181 KRAEVVVPNGSIRVNGGGGLINAASVTLTSGVPVL-NNGNLTGFDVSSGKVVIGGKGL-D 238
50     Query: 224 KGGADYTGVLARAVALQGKLGKKNLAVSTGPKQVDYASGEISAGTAAGTK----PTIALD 279
      Sbjct: 239 TSDADYTRILSRAAEINAGVWGDVKVSVSGKNKLDGSLAKTASAPSSSDSVTPTVAID 298
55     Query: 280 TAALGGMYADSITLIAXEKGVGVKNAGTLEAAK-QLIVTSSGRIENSGRIATTADGTEAS 338
      Sbjct: 299 TATLGGMYADKITLISTDNGAVIRNKGRIFAATGGVTLSDGKLSNSGSI-----DAA 351
60     Query: 339 PTYLXIETTEKGAXGTFISNGGRIESKGLLVIIETGEDIXLRNGAVVQNNGSRPATTVLNA 398
      Sbjct: 352 EITISAQTVD-----NRQGFIRSGKGSVLKVS DGINNQAGLI-----GSAGLLDIRDT 399
65     Query: 399 GHNLVIESKTNVNNAGKS-----XNLSAGGRTTINDATIQAGSSSVYSTKGDXTLGENTRI 454
      Sbjct: 400 G-----KSSLHINNTDGTIIAGKDVSLQAKSLDNDGILTAARDV-SVSLHDDFAGKRDIE 453
      Query: 455 IAENVTVLSNGSIGSAAVIEAKDTAHIESGKPLSLETSTVASNIRLNNNGNIKGGKQLALL 514
      Sbjct: 455 +T + G + + +I+A DT + + + + + S R G L+

```


Sbjct: 454 AGRTLTFSTQGRLLKNTRI IQAGDTVSLTAAQIDNTVSGKIQSGNRTGLNGKNGITNRGLI 513
 Query: 515 ADDNIT-----AKTTNLTNTPGNLYVHTGKDLNLDKDLAASIHLSKSDNAAHITGTSKT 569
 + IT AK+ N T G +Y G + + D L+ AA
 Sbjct: 514 NSNGITLLQTEAKSDNAGT-GRIY---GSRVAVEADTLLNREETVNGETKAA-----V 562
 Query: 570 LTASKDMGVEAGXXXXXXXXXXXXSGNLHIQAA--KGNIQLRNTKL-NAKALETALQ 625
 + A + + + A SG+LHI +A +Q NT L N + A+E++
 Sbjct: 563 IAAERLDIGAREIENREAAALSSSGDLHIGSALNGSRQVQGANTSLHNRSAAIESS--- 619
 Query: 626 GNI 628
 GNI
 Sbjct: 620 GNI 622

 Score = 37.5 bits (85), Expect = 0.53
 Identities = 87/432 (20%), Positives = 159/432 (36%), Gaps = 62/432 (14%)
 Query: 239 LQGLQGKNLAVSTGPQKVDYASGEISAGTAAGTKPTIALDTAALGGMYADSITLIAEXE 298
 LQ LQGN+ + G + +G I A A K A + + S T +
 Sbjct: 1023 LQGD LQGNIFAAAGSDITN--TGSIGAENALLK-----ASNNIERSSETRSNQNE 1072
 Query: 299 GVGKVNAGTLEAAKQLIVTSSGRI--ENSGRIATTADGTEASPTYLXIETTEKGAXG-TF 355
 V+N G + A L +G + + I TA E T + G T
 Sbjct: 1073 QGSRVNI GRV-AGIYLTGRQNGSVLLDAGNNIVLTAS-----ELTNQSEDGQTV 1120
 Query: 356 ISNGGRIESKGLLVETGEDI XLNRGAVVQNNNGSRPATTVLNAGHNLVIESK-----T 408
 ++ GG I S + I + V++ + +T+ G NL + +K
 Sbjct: 1121 LNAGGDIRSDTTGISRNQNTIFDS DNYVIRKEQNEVGSTIRTRG-NLSLNAKGDIRIRAA 1179
 Query: 409 NVNNAKGSXNLSAGGRTTINDATI QAGSS-----VYSSTKGDTXLGENTRIIAENV 460
 V + +G L+AG D ++AG + Y+ G + TR +
 Sbjct: 1180 EVGSEQRLKLAAG-----RDIKVEAGKAHTETEDALKYTGRSGGGIKQKMTRHLKNQNG 1234
 Query: 461 VLSNGSIGSAAVIEAKDTAHIESGKPLSLETSTVASNIRLNNGNIKGGKQLALLADDNIT 520
 +G++ +I +G + + T+ S NN +K + + A+ N
 Sbjct: 1235 QAVSGTLDGKEIILVSGRDITVTGSNIIADNHTILS--AKNNIVLKAETR SRSAEMNKK 1292
 Query: 521 AKTTNLTNTPG-NLYVHTGKDLNLDKDLAASIHLSKSDN-----AAHITGTSKTLTA 572
 K+ + + G + KD N + +S + S N H T T T+++
 Sbjct: 1293 EKSGLMGSGGIGFTAGSKKDTQTNRSETVSHTESVVGSLNGNTLISAGKHYTQTGSTISS 1352
 Query: 573 SK-DMGVEAGXXXXXXXXXXXXSGNLHIQAAKG-----NIQLRNTKLNAKALETALQ 626
 + D+G+ +G + + KG ++ + NT + A A++ G
 Sbjct: 1353 PQGDVGISSGKISIDAAQNRYSQESKQVYEKGVTVAISVPVNTVMGAVDKAVQTVG 1412
 Query: 627 NIVSDGLHAVSA 638
 + ++A++A
 Sbjct: 1413 KSKNSRVNAMA 1424

Amino acids 1-1423 of ORF114-1 were cloned in the pGex vector and expressed in *E.coli*, as described above. GST-fusion expression was visible using SDS-PAGE, and Figure 5 shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF114-1.

Based on these results, including the homology with the putative secreted protein of *N.meningitidis* and on the presence of a transmembrane domain, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 14

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 63>

```

1  ..CGCTTCATTC ATGATGAAGC AGTCGGCAGC AACATCGGCG GCGGCAAAAT
51  GATTGTTGCA GCCGGGCAGG ATATCAATGT ACGCGGCAnA AGCCTTATTT
101 CTGATAAGGG CATTGTTTTA AAAGCAGGAC ACGACATCGA TATTCTACT
151 GCCCATAATC GCTATACCGG CAATGAATAC CACGAGAGCA wAAAwTCAGG
5  201 CGTCATGGGT ACTGGCGGAT TGGGCTTTAC TATCGGTAAC CGGAAACTA
251 CCGATGACAC TGATCGTACC AATATTGTsC ATACAGGCAG CATTATAGGC
301 AGCCTGAaTG GAGACACCGT TACAGTTGCA GGAAACCGCT ACCGACAAAC
351 CGGCAGTACC GTCTCCAGCC CCGAGGGGCG CAATACCGTC ACAGCCAAAw
10  401 GCATAGATGT AGAGTTCGCA AACAAACCGGT ATGCCACTGA CTACGCCAT
451 ACCCAgGGAA CAAAAAGGCC TTACCGTCGC CCTCAATGTC CCGGTTGTCC
501 AAGCTGCACA AAACCTTCATA CAAGCAGCCC AAAATGTGGG CAAAAGTAAA
551 AATAAACGCG TTAATGCCAT GGCTGCAGCC AATGCTGCAT GGCAGAGTTA
601 TCAAGCAACC CAACAATGC AACAAATTTGC TCCAAGCAGC AGTCCGGGAC
15  651 AAGGTCAAAA CTACAATCAA AGCCCCAGTA TCAGTGTGTC CATTAC.TAC
701 GGCGAACAGA AAAGTCGTAA CGAGCAAAAA AGACATTACA CCGAaGCGGC
751 AgCAAGTCAA ATTATCGGCA AAGGGCAAAC CACACTTGCG GCAACAGGAA
801 GTGGGGAGCA GTCCAATATC AATATTACAG GTTCCGATGT CATCGGCCAT
851 GCAGGTACTC C.CTCATTGC CGACAACCAT ATCAGACTCC AATCTGCCAA
901 ACAGGACGGC AGCGAGCAAA GCAAAAAACAA AAGCAGTGGT TGGAATGCAG
20  951 GCGTACGTnn CAAAATAGGC AACGGCATCA GGTTTGGAAT TACCGCCGGA
1001 GGAATATCG GTAAAGGTAA AGAGCAAGGG GGAAGTACTA CCCACGGCCA
1051 CACCCATGTC GGCAGCACAA CCGGCAAAC TACCATCCGA AGCGGCGGGG
1101 GATACCACCC TCAAAGGTGT GCAGCTCATC GGCAAGGCA TACAGGCAGA
1151 TACGCGCAAC CTGCATATAG AAAGTGTTCa AGATACTGAA ACCTATCAGA
25  1201 GCAAAACAGCA AAACGGCAAT GTCCAAGTTt ACTGTCGGTT ACGGATTcAG
1251 TGCAAGCGGC AGTTACCGCC AAAGCAAAGT CAAAGCAGAC CATGCCTCCG
1301 TAACCGGGCA AAgCGGTATT TATGCCGGAG AAGACGGCTA TCAAATyAAA
1351 GTyAGAGACA ACACAGACCT yAAGGGCGGT ATCATCACGT CTAGCCAAAG
1401 CGCAGAAGAT AAGGGCAAAA ACCTTTTTTCa GACGGCCACC CTACTGGCCA
30  1451 GCGACATTCA AAACCACAGC CGCTACGAAG GCAGAAGCTT CGGCATAGGC
1501 GGCAGTTTCG ACCTGAACGG CGGCTGGGAC GGCACGGTTA CCGACAAACA
1551 AGGCAGGCCT ACCGACAGGA TAAGCCCGGC AGCCGGCTAC GGCAGCGACG
1601 GAGACAGCAA AAACAGCACC ACCCGCAGCG GCGTCAACAC CCACAACATA
1651 CACATCACCG ACGAAGCGGG ACAACTTGCC CGAACAGGCA GGA CTGCAAA
35  1701 AGAAACCGAA GCGCGTATCT ACACCGGCAT CGACACCGAA ACTGCGGATC
1751 AACACTCAGG CCATCTGAAA AACAGCTTCG AC...

```

This corresponds to the amino acid sequence <SEQ ID 64; ORF116>:

```

1  ..RFIHDEAVGS NIGGGKMIVA AGQDINVRGX SLISDKGIVL KAGHDIDIST
40  51  AHNRYTGNEY HESXXSGVMG TGGLGFTIGN RKTDDTDRT NIVHTGSIIG
101  SLNGDTVTVa GNRYRQTGST VSSPEGRNTV TAKXIDVEFA NNRyATDYAH
151  TQEQKGLTVA LNPVvVQAAQ NFIQAAQNVG KSKNKRVNAM AAANAaWQSY
201  QATQOMQQFA PSSSAGQQQN YNQSPSISVS IXyGEQKSRN EQKRHYTEAA
251  ASQIIIGKQT TLAATGSSEQ SNINITGSDV IGHAGTXLIA DNHIRLQSAK
301  QDGSEQSKNK SSGWNAGVRX KIGNGIRFGI TAGGNIGKGK EQGGSTTHRH
45  351  THVGSTTGKT TIRSGDttl KGVQLIGKI QADTRNLHIE SVQDTETyQS
401  KQONGNVQVT VYGfSASGS YRQSKVKADH ASVTGQSGIY AGEDGYQIKV
451  RDNTDLKGGI ITSSQSAEDK GKnlFQTATL TASDIQNHSR YEGRSFGIGG
501  SFDLNGGWDG TVTDKQGRPT DRISPAAGYG SDGDSKNSTT RSGVNTHNIH
551  ITDEAGQLAR TGRtAKETEA RIYTGIDTET ADQHSgHLKN SFD...

```

Computer analysis of this amino acid sequence gave the following results:

Homology with pspA putative secreted protein of *N.meningitidis* (accession number AF030941)

ORF116 and pspA protein show 38% aa identity in 502aa overlap:

```

Orf116: 6  EAVGSNIGGGKMIVAAGQDINVRGXSLISDKGIVLKAGHDIDISTAHNRYTGNEYHESXX 65
55  +AV + G ++I+ +G+DI V G ++I+D +L A ++I + A R E ++
PspA: 1 235 QAVSGTLdGKEIILVSGRDITVTGSNIADNHtILSAKNNIVLKAaETRSRSaEMNKKEK 1294
Orf116: 66  XXXXXXXXXXXXXXXNRKXXXXXXXXRTNIVHTGSIIGSLNGDTVTVAGNRYRQTGSTVSSPE 125
++K + HT S++GSLNG+T+ AG Y QTGST+SSP+
PspA: 1295 SGLMGSGGIGFTAGSKKDTQTNrSETVSHTESVVGSLNGNTLISAGKHytQTGSTISSPQ 1354
60

```

Orf116: 126 GRNTVTAKXIDVEFANNRYATDYAHTQEOKGLTVALNVPXXXX---XXXXXXXXXXXXGKS 182
 G +++ I ++ A NRY+ + EQKG+TVA++VP GKS
 PspA: 1355 GDVGISSGKISIDAAQNRYSQESKQVYEQKGVTV AISVPVNTVMGAVDAVKAVQTVGKS 1414

5 Orf116: 183 KNKRXXXXXXXXXXWQSYQATQMQQFA--PSSSAGQGQNYNQSPSISVSIXYGEQKSRN 240
 KN RV + + + A P +AGQG ISVS+ YGEQK+ +
 PspA: 1415 KNSRVNMAAANALNKGVD SGVALYNAARNPKKAAGQG-----ISVSVTYGEQKNTS 1466

10 Orf116: 241 EQKRHYTEAAASQIIIGKQTTLAATGSGEQSNINITGSDVIGHAGTXLIADNHIRLQSAK 300
 E + T+ +I G G+ +L A+G+G+ S I ITGSDV G GT L A+N +++++A+
 PspA: 1467 ESRIKGTQVQEGKITGGGKVS LTASGAGKDSRITITGSDVYGGKGT RLKAENAVQIEAAR 1526

15 Orf116: 301 QDGSEQSKNKSSGWNAGVRXKIGNGIRFGITAXXXXXXXXXXXXXSTHRRHTHVGSTTGKT 360
 Q E+S+NKS+G+NAGV I GI FG TA T +R++H+GS +T
 PspA: 1527 QTHQERSENKSAGFNAGVAIAINKGISFGFTAGANYGKG YGNGDETAYRNSHIGSKDSQT 1586

20 Orf116: 361 TIRSGGDTTLKGVQLIGKGIQADTRNLHIESVQDTET YQSKQQNGNVQVTVG YGFSASGS 420
 I SGGDT +KG QL GKG+ +LHIES+QDT ++ KQ+N + QVTVG YGFS GS
 PspA: 1587 AIESGGDTVIKGGQLKGKGVGVTAE SLHIESLQDTAVFKGKQENVSAQVTVG YGFSVGGGS 1646

25 Orf116: 421 YRQSKVKADHASVTGQS GYAGEDGYQIKVRDNTDLKGGIITSSQSAEDKGNL FQTATL 480
 Y +SK +D+ASV QSGI+AG DGY+I+V T L G + S DK KNL +T+ +
 PspA: 1647 YNRKSSSDYASVNEQSGIFAGGDGYRIRVNGKTGLVGA AVVSD---ADKSKNLLKTSEI 1703

25 Orf116: 481 TASDIQNHSRYEGRSFGIGGSF 502
 DIQNH+ + G+ G F
 PspA: 1704 WHKDIQNHASAAASALGLSGGF 1725

Based on homology with *pspA*, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

30 Example 15

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 65>

35 1 ..ACGACCGGCA GCCTCGGCGG CATACTGGCC GGCGGCGGCA CTTCCCTTGC
 51 CGCACCGTAT TTGGACAAAG CGGCGGAAAA CCTCGGTCCG GCGGGCAAAG
 101 CGGCGGTCAA CGCACTGGGC GGTGCGGCCA TCGGCTATGC AACTGGTGGT
 151 AGTGGTGGTG CTGTGGTGGG TGCGAATGTA GATTGGAACA ATAGGCAGCT
 201 GCATCCGAAA GAAATGGCGT TGGCCGACAA ATATGCCGAA GCCCTCAAGC
 251 GCGAAGTTGA AAAACGCGAA GGCAGAAAAA TCAGCAGCCA AGAAGCGGCA
 301 ATGAGAATCC GCAGGCAGAT ATGCGTTGGG TGGACAAAGG TTCCCAAGAC
 351 GGCTATACCG ACCAAAGCGT CATATCCCTT ATCGGAATGA

40 This corresponds to the amino acid sequence <SEQ ID 66; ORF118>:

1 ..TTGSLGGILA GGGTSLAAPY LDKAAENLGP AGKAAVNALG GAAIGYATGG
 51 SGGAVVGANV DWNRRQLHPK EMALADKYAE ALKREVEKRE GRKISSQEAA
 101 MRIRRQICVG WTKVPKTAIP TKASYPLSE*

Computer analysis of this amino acid sequence reveals two putative transmembrane domains.

45 Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 16

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 67>

1 ..CAATGCCGTC TGAAAAGCTC ACAATTTTAC AGACGGCATT TGTATGCAA

51 GTACATATAC AGATTCCCTA TATACTGCCC AGrkGCGTGC GTgGCTGAAG
 101 ACACCCCCTA CGCTTGCTAT TTGrAACAGC TCCAAGTCAC CAAAGACGTC
 151 AACTGGAACC AGGTACwACT GGC GTACGAC AAATGGGACT ATAAACAGGA
 201 AGGCTTAACC GGAGCCGAG CAGCGATTAT TGCGCTGGCT GTTACCGTGG
 5 251 T TACTGCGGG CGCGGGAgCC GGAGCCGCAC TGGGcTTAAA CGGCGCGGcC
 301 GCAGCGGCAA CCGATGCCGC ATTCCGCTCG CTGGCCAGCC AGGcTTCCGT
 351 ATCGCTCATC AaCAACAAAG GCAATATCGG TAaCACCCCTG AAAGAGCTGG
 401 GCAGAAGCAG CACGGTGAAA AATCTGATGG TTGCCGTCGc tACCGCAgGC
 10 451 GTagCcgaCA AAATCGGTGC TTCGGCACTG AACAAATGTCA GCATAAGCA
 501 GTGGATCAAC AACCTGACCG TCAACCTGGC CAATGCGGGC AGTGCCGCAC
 551 TGATTAATAC CGCTGTCAAC GCGCGCAGCc tgAAAGACAA TCTGGAAGCG
 601 AATATCCTTG CGGCTTGGT GAATACTGCG CATGGAGAAG CAGCCAGTAA
 651 AATCAAAACAG TTGGATCAGC ACTACATTAC CCACAAGATT GCCCA TGCCA
 15 701 TAGCGGGCTG TCGGgTGCG GCGGCGAATA AGGGCAAGTG TCAGGATGGT
 751 GCGATAgGTG CCGCTGTGGG CGAGATAGTC GGGGAgGCTT TGACAAACGG
 801 CAAAAATCCT GACACTTTGA CAGCTAAAgA ACGCGaACAG ATTTTGGCAT
 851 ACAGCAAAC TGGTTGCCGT ACGGTAAGCG GTGTGGTCCG CGGCGATGTA
 901 AATGCGGCGG CGAATGCGGC TGAGGTAGCG GTGAAAAATA ATCAGCTTAG
 951 CGACAAAtGA

20 This corresponds to the amino acid sequence <SEQ ID 68; ORF41>:

1 ..QCRLKSSQFY RRHLLCKYIY RFPIYCPKAC VAEDTPYACY LXQLQVTKDV
 51 NWNQVXLAYD KWDYKQEGLT GAGAAIALA VTVVTAGAGA GAALGLNGAA
 101 AAATDAAFAS LASQASVSLI NNKGNIGNTL KELGRSSTVK NLMVAVATAG
 151 VADKIGASAL NNVSDKQWIN NLTVNLANAG SAALINTAVN GGLSKDNLEA
 201 NILAALVNTA HGEAAASKIKQ LDQHYITHKI AHAIAGCAAA AANKGKCQDG
 251 AIGAAVGEIV GEALTNGKNP DTLTAKEREQ_ILAYSKLVAG TVSGVVGVDV
 301 NAAANAAEVA VKNNQLSDK*

Further work revealed the complete nucleotide sequence <SEQ ID 69>:

30 1 ATGCAAGTAA ATATTAGAT TCCCTATATA CTGCCCAGAT GCGTGCGTGC
 51 TGAAGACACC CCCTACGCTT GCTATTTGAA ACAGCTCCAA GTCACCAAAG
 101 ACGTCAACTG GAACCAGGTA CAACTGGCGT ACGACAAATG GGAATAAAA
 151 CAGGAAGGCT TAACCGGAGC CGGAGCAGCG ATTATTGCGC TGGCTGTTAC
 201 CGTGGTTACT GCGGGCGCGG GAGCCGGAGC CGCACTGGGC TTAACGGCG
 251 CGGCCGAGC GGCAACCGAT GCCGCATTG CCTCGCTGGC CAGCCAGGCT
 35 301 TCCGTATCGC TCATCAACAA CAAAGGCAAT ATCGGTAACA CCCTGAAAGA
 351 GCTGGGCAGA AGCAGCACGG TGA AAAATCT GATGGTTGCC GTCGCTACCG
 401 CAGGCGTAGC CGACAAAATC GGTGCTTCGG CACTGAACAA TGTGAGCAT
 451 AAGCAGTGA TCAACAACT GACCGTCAAC CTGGCCAATG CGGCGAGTGC
 501 CGCACTGATT AATACCGCTG TCAACGGCGG CAGCCTGAAA GACAATCTGG
 40 551 AAGCGAATAT CCTTGCGGCT TTGGTGAATA CTGCGCATGG AGAAGCAGCC
 601 AGTAAATCA AACAGTTGGA TCAGCACTAC ATTACCCACA AGATTGCCCCA
 651 TGCCATAGCG GGCTGTGCGG CTGCGGCGCG GAATAAGGGC AAGTGTCAAG
 701 ATGGTGCGAT AGGTGCGGCT GTGGGCGAGA TAGTCGGGGA GGCTTTGACA
 751 AACGGCAAAA ATCCTGACAC TTTGACAGCT AAAGAACCGC AACAGATTTT
 45 801 GGCATACAGC AAACCTGGTTG CCGGTACGGT AAGCGGTGTG GTCGGCGGGC
 851 ATGTAAATGC GCGGCGAAT GCGGCTGAGG TAGCGGTGAA AAATAATCAG
 901 CTTAGCGACA AAGAGGGTAG AGAATTTGAT AACGAAATGA CTGCATGCGC
 951 CAAACAGAAT AATCCTCAAC TGTGCAGAAA AAATACTGTA AAAAAGTATC
 1001 AAAATGTTGC TGATAAAGA CTTGCTGCTT CGATTGCAAT ATGTACGGAT
 50 1051 ATATCCCGTA GACTGAATG TAGAACATC AGAAAACAAC ATTTGATCGA
 1101 TAGTAGAAGC CTTATTTCAT CTTGGGAAGC AGGTCTAATT GGTAAAGATG
 1151 ATGAATGGTA TAAATTATTC AGCAATCTT ACACCCAAGC AGATTTGGCT
 1201 TTACAGTCTT ATCATTTGAA TACTGCTGCT AAATCTTGGC TTCAATCGGG
 1251 CAATACAAAG CTTTTATCCG AATGGATGTC CGACCAAGGT TATACACTTA
 55 1301 TTTCAAGAGT TAATCCTAGA TTCATTCCAA TACCAAGAGG GTTTGTAAAA
 1351 CAAAATACAC CTATTACTAA TGTCAAATAC CCGGAAGGCA TCAGTTTCGA
 1401 TACAAACCTA AAAAGACATC TGGCAAATGC TGATGGTTTT AGTCAAAAAC
 1451 AGGCGATTAA AGGAGCCCAT AACCGCACCA ATTTTATGGC AGAATCAAT
 1501 TCACGAGGAG GACGCGTAAA ATCTGAAACC CAACTGATA TTGAAGGCAT
 60 1551 TACCGAATT AAATATGAGA TTCCTACACT AGACAGGACA GGTAAACCTG
 1601 ATGGTGGATT TAAGGAAAT TCAAGTATAA AAATGTTTA TAATCCTAAA
 1651 AAATTTCTG ATGATAAAAT ACTTCAAATG GCTCAAAATG CTGCTTACCA
 1701 AGGATATTCA AAAGCCTCTA AAATTGCTCA AAATGAAAGA ACTAAATCAA
 1751 TATCGGAAAG AAAAAATGTC ATTCAATTCT CAGAAACCTT TGACGGAATC
 65 1801 AAATTTAGAT CATATTTTGA TGTAAATACA GGAAGAATTA CAAACATTCA
 1851 CCCAGAATAA

This corresponds to the amino acid sequence <SEQ ID 70; ORF41-1>:

```

1  MQVNIQIPYI LPRCVR AEDT PYACYLKQLQ VTKDVNWNQV QLAYDKWDYK
51  QEGLTGAGAA IIALAVTVVT AGAGAGAALG LNGAAAAATD AAFASLASQA
101 SVSLINNKG N IGNTLKE LGR SSTVKNLMVA VATAGVADKI GASALNNVSD
151 KQWINNLT V N LANAGSAAL I NTAVNGGSLK DNLEANILAA LVNTAHGEAA
201 SKIKQLDQHY ITHKIAHAIA GCAAAAANKG KCQDGAIGAA VGEIVGEALT
251 NGKNPDTLT A KEREQILAYS KLVAGTVSGV VGGDVNAAAN AAEVAVKNNQ
301 LSDKEGREFD NEMTACAKQN NPQLCRKNTV KKYQNVADKR LAASIAICTD
351 ISRSTECRTI RKQHLIDSR S LHSSWEAGLI GKDDWEYKLF SKSYTQADLA
401 LQSYHLNTAA KSWLQSGNTK PLSEWMSDQG YTLISGVNPR FIPIPRGFVK
451 QNTFITNVKY PEGISFDTNL KRHLANADGF SQKQGIKGAH NRTNFM AELN
501 SRGGRVKSET QTDIEGITRI KYEIPDLRT GKPDGGFKEI SSIKTVYNPK
551 KFSDDKILQM AQNAASQGY S KASKIAQNER TKSISERKNV IQFSETFDGI
601 KFSYFDVNT GRITNIHPE*

```

- 15 Computer analysis of this amino acid sequence predicts a transmembrane domain, and homology with an ORF from *N.meningitidis* (strain A) was also found.

ORF41 shows 92.8% identity over a 279aa overlap with an ORF (ORF41a) from strain A of *N. meningitidis*:

```

20  orf41.pep      10      20      30      40      50      60      69
    orf41a              YRRHLLCKYIYRFPIYCPXACVAEDTPYACYLXQLQVTKDVNWNQVXLAYDKWDYKQEGL.
                                || |||:|:| ||| |||:|:| |||
                                YLKQLQVAKNINWNQVQLAYDRWDYKQEGL
                                10      20      30
25  orf41.pep      70      80      90      100     110     120     129
    orf41a      TGAGAAIIALAVTVVTAGAGAGAALGLNGAAAAATDAAFASLASQASVSLINNKG NIGNT
    orf41a      TEAGAAIIALAVTVVTSAGTGAVLGLNGAXAAATDAAFASLASQASVSFINNKG DVGKT
                                40      50      60      70      80      90
30  orf41.pep      130     140     150     160     170     180     189
    orf41a      LKELGRSSTVKNLMVAVATAGVADKIGASALNNVSDKQWINNLT VNLNAGSAALINTAV
    orf41a      LKELGRSSTVKNLVVAAATAGVADKIGASALXNVSDKQWINNLT VNLNAGSAALINTAV
35  orf41a              100     110     120     130     140     150
40  orf41.pep      190     200     210     220     230     240     249
    orf41a      NGGSLKDNLEANILAALVNTAHGEAASKIKQLDQHYITHKIAHAIAGCAAAAANKGKCQD
    orf41a      NGGSLKDXLEANILAALVNTAHGEAASKIKQLDQHYIVHKIAHAIAGCAAAAANKGKCQD
                                160     170     180     190     200     210
45  orf41.pep      250     260     270     280     290     300     309
    orf41a      GAIGAAVGEIVGEALTNGKNPDTLTAKEREQILAYSKLVAGTVSGVVGGDVNAAANA AEV
    orf41a      GAIGAAVGEIVGEALTNGKNPDTLTAKEREQILAYSKLVAGTVSGVVGGDVNAAANA AEV
                                220     230     240     250     260     270
50  orf41.pep      310     320
    orf41a      AVKNNQLSDKX
    orf41a      AVKNNQLSDXEGREFDNEMTACAKQNXNPQLCRKNTVKKYQNVADKRLAASIAICTDISRS
                                280     290     300     310     320     330

```

A partial ORF41a nucleotide sequence <SEQ ID 71> is:

```

55  1  ..TATCTGAAAC AGCTCCAAGT AGCGAAAAAC ATCAACTGGA ATCAGGTGCA
51  51 GCTTGCTTAC GACAGATGGG ACTACAAACA GGAGGGCTTA ACCGAAGCAG
101 101 GTGCGGCGAT TATCGCACTG GCCGTTACCG TGGTCACCTC AGGCGCAGGA
151 151 ACCGGAGCCG TATTGGGATT AAACGGTGCG NCCGCCGCCG CAACCGATGC

```

201 AGCATTTCGCC TCTTTGGCCA GCCAGGCTTC CGTATCGTTC ATCAACAACA
 251 AAGGCGATGT CGGCAAAACC CTGAAAGAGC TGGGCAGAAG CAGCACGGTG
 301 AAAAATCTGG TGGTTGCCGC CGCTACCGCA GGCGTAGCCG ACAAATCGCG
 351 CGCTTCGGCA CTGANCAATG TCAGCGATAA GCAGTGGATC AACAACTGA
 401 CCGTCAACCT AGCCAATGCG GGCAAGTCCG CACTGATTAA TACCGCTGTC
 451 AACGGCGGCA GCCTGAAAGA CANTCTGGAA GCGAATATCC TTGCGGCTTT
 501 GGTCAATACC GCGCATGGAG AAGCAGCCAG TAAATCAAA CAGTTGGATC
 551 AGCACTACAT AGTCCACAAG ATTGCCCATG CCATAGCGGG CTGTGCGGCA
 601 GCGGCGGCGA ATAAGGCAA GTGTCAGGAT GGTGCGATAG GTGCGCTGT
 651 GGGCGAGATA GTCGGGGAGG CTTTGACAAA CGGCAAAAT CCTGACACTT
 701 TGACAGCTAA AGAACGCGAA CAGATTTTGG CATACAGCAA ACTGGTTGCC
 751 GGTACGGTAA GCGGTGTGGT CCGCGGCGAT GTAAATGCGG CGGCGAATGC
 801 GGCTGAGGTA GCGGTGAAAA ATAATCAGCT TAGCGACNAA GAGGTCAGAG
 851 AATTTGATAA CGAAATGACT GCATGCGCCA AACAGAATAN TCCTCAACTG
 901 TGCAGAAAAA ATACTGTAAA AAAGTATCAA AATGTTGCTG ATAAAAGACT
 951 TGCTGCTTCG ATTGCAATAT GTACGGATAT ATCCCGTAGT ACTGAATGTA
 1001 GAACAATCAG AAAACAACAT TTGATCGATA GTAGAAGCCT TCATCTGTA
 1051 TGGGAAGCAG GTCTAATTGG TAAAGATGAT GAATGGTATA AATTATTGAG
 1101 CAAATCTTAC ACCCAAGCAG ATTTGGCTTT ACAGTCTTAT CATTTGAATA
 1151 CTGCTGCTAA ATCTTGGCTT CAATCGGGCA ATACAAAGCC TTTATCCGAA
 1201 TGGATGTCCG ACCAAGGTTA TACACTTATT TCAGGAGTTA ATCCTGATT
 1251 CATTCCAATA CCAAGAGGGT TTGTAAAACA AAATACACCT ATTACTAATG
 1301 TCAAATACCC GGAAGGCATC AGTTTCGATA CAAACCTANA AAGACATCTG
 1351 GCAAATGCTG ATGGTTTTAG TCAAGAACAG GGCATTAAAG GAGCCCATAA
 1401 CCGCACCAAT NTTATGGCAG AACTAAATTC ACGAGGAGGA NGNGTAAAT
 1451 CTGAAACCCA NACTGATATT GAAGGCATTA CCCGAATTAA ATATGAGATT
 1501 CCTACACTAG ACAGGACAGG TAAACCTGAT GGTGGATTTA AGGAAATTTT
 1551 AAGTATAAAA ACTGTTTATA ATCCTAAAAA NTTTTNNGAT GATAAAATAC
 1601 TTCAAATGGC TCAANATGCT GNTTCACAAG GATATTCAA AGCCTCTAAA
 1651 ATTGCTCAA ATGAAAGAAC TAAATCAATA TCGGAAAGAA AAAATGTCAT
 1701 TCAATTCTCA GAAACCTTTG ACGGAATCAA ATTTAGANNN TATNTNGATG
 1751 TAAATACAGG AAGAATTACA AACATTACCC CAGAATAA

This encodes a protein having the partial amino acid sequence <SEQ ID 72>:

1 YLKQLQVAKN INWNQVQLAY DRWDYKQEGL TEAGAAIAL AVTVVTSGAG
 51 TGAVLGLNGA XAAATDAFA SLASQASVSF INNKGDVGKT LKELGRSSTV
 101 KNLVVAATA GVADKIGASA LXNVSDKQWI NNLTVNLANA GSAALINTAV
 151 NGGSLKDXLE ANILAALVNT AHGEAASKIK QLDQHYIVHK IAHAAGCAA
 201 AAANKGKCQD GAIGAAGEI VGEALTNGKN PDTLTAKERE QILAYSKLVA
 251 GTVSGVVGVD VNAANAAEV AVKNNQLSDX EGREFDNEMT ACAKQXPQL
 301 CRKNTVKKYQ NVADKRLAAS IAICTDISRS TECRTIRKQH LIDSRSLHSS
 351 WEAGLIGKDD EWYKLFKSY TQADLALQSY HLNTAAKSWL QSGNTKPLSE
 401 WMSDQGYTLI SGVNPFIPI PRGFVKQNTF ITNVKYPEGI SFDTNLXRHL
 451 ANADGFSQEQ GIKGAHNRTN XMAELNSRGG XVKSETXTDI EGITRIKYEI
 501 PTLDRGTGPD GGFKEISSIK TVYNPKXFXD DKILQMAQXA XSQGYSKASK
 551 IAQNERTKSI SERKNVIQFS ETFDGIKFRX YXDVTNGRIT NIHPE*

ORF41a and ORF41-1 show 94.8% identity in 595 aa overlap:

orf41a.pep 10 20 30
 YLKQLQVAKNINWNQVQLAYDRWDYKQEGLTEAGAA
 50 orf41-1 MQVNIQIPYILPRCVRAEDTPYACYLKQLQVTKDVNWNQVQLAYDKWDYKQEGLTGAGAA
 10 20 30 40 50 60
 orf41a.pep 40 50 60 70 80 90
 IIALAVTVVTSGAGTGAVLGLNGAXAAATDAAFASLASQASVSFINNKGDVGKTLKELGR
 55 orf41-1 IIALAVTVVTAGAGAGAALGLNGAAAAATDAAFASLASQASVSLINNKGNIQNTLTKELGR
 70 80 90 100 110 120
 orf41a.pep 100 110 120 130 140 150
 SSTVKNLVVAATAGVADKIGASALXNVSDKQWNNLTVNLANAGSAAALINTAVNGGSLK
 orf41-1 SSTVKNLVAVATAGVADKIGASALNNVSDKQWNNLTVNLANAGSAAALINTAVNGGSLK
 130 140 150 160 170 180
 65 orf41a.pep 160 170 180 190 200 210
 DXLEANILAALVNTAHGEAASKIKQLDQHYIVHKIAHAAGCAAAAANKGKCQDGAIGAA

-100-

	orf41-1	 DNLEANILAA LVNTAHGEAASKIKQLDQHYITHKIAHAIAGCAAAAANKGKCQDGAIGAA 190 200 210 220 230 240
5	orf41a.pep	220 230 240 250 260 270 VGEIVGEALTNGKNPDTLTAKEREQILAYSKLVAGTVSGVVGGDVNAAANAAEVAVKNNQ
10	orf41-1	250 260 270 280 290 300 VGEIVGEALTNGKNPDTLTAKEREQILAYSKLVAGTVSGVVGGDVNAAANAAEVAVKNNQ
	orf41a.pep	280 290 300 310 320 330 LSDXEGREFDNEMTACAKQNXPLCRKNTVKKYQNVADKRLAASIAICTDISRSTECRTI
15	orf41-1	310 320 330 340 350 360 LSDKEGREFDNEMTACAKQNNPQLCRKNTVKKYQNVADKRLAASIAICTDISRSTECRTI
	orf41a.pep	340 350 360 370 380 390 RKQHLIDSRSLHSSWEAGLIGKDDWEYKLFSSYQADLALQSYHLNTAAKSWLQSGNTK
20	orf41-1	370 380 390 400 410 420 RKQHLIDSRSLHSSWEAGLIGKDDWEYKLFSSYQADLALQSYHLNTAAKSWLQSGNTK
	orf41a.pep	400 410 420 430 440 450 PLSEWMSDQGYTLISGVNPRFIPIPRGFVKQNTPTITNVKYPEGISFDTNLXRHLANADGF
25	orf41-1	430 440 450 460 470 480 PLSEWMSDQGYTLISGVNPRFIPIPRGFVKQNTPTITNVKYPEGISFDTNLKRHLANADGF
	orf41a.pep	460 470 480 490 500 510 SQEQGIKGAHNRTNXMAELNSRGGXVKSETXTDIEGITRIKYEIPTLDRGTGKPDGGFKEI :
30	orf41-1	490 500 510 520 530 540 SQKQGIKGAHNRTNFMALNSRGGRVKSETQTDIEGITRIKYEIPTLDRGTGKPDGGFKEI
	orf41a.pep	520 530 540 550 560 570 SSIKTVYNPKKFXDDKILQMAQXASQGYSKASKIAQNERTKSISERKNVIQFSETFDGI
35	orf41-1	550 560 570 580 590 600 SSIKTVYNPKKFSDDKILQMAQNAASQGYSKASKIAQNERTKSISERKNVIQFSETFDGI
40	orf41a.pep	580 590 KFRXYXDVNTGRITNIHPEX
45	orf41-1	610 620 KFRSYFDVNTGRITNIHPEX

Amino acids 25-619 of ORF41-1 were amplified as described above. Figure 6 shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF41-1.

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

50. Example 17

The following DNA sequence was identified in *N.meningitidis* <SEQ ID 73>

	1	ATGGCAATCA	TTACATTGTA	TTATTCTGTC	AATGGTATTT	TAAATGTATG
	51	TGCAAAAGCA	AAAAATATTC	AAGTAGTTGC	CAATAATAAG	AATATGGTTC
	101	TTTTTGGGTT	TTTGGsmrGC	ATCATCGGCG	GTTCAACCAA	TGCCATGTCT
55	151	CCCATATTGT	TAATATTTT	GCTTAGCGAA	ACAGAAAATA	AAAATcgTAT
	201	CGTAAAATCA	AGCAATCTAT	GCTATCTTTT	GGCGAAAATT	GTTCAAATAT
	251	ATATGCTAAG	AGACCAGTAT	TGGTTATTAA	ATAAGAGTGA	ATACGdTTTA
	301	ATATTTTAC	TGTCCGTATT	GTCTGTTATT	GGATTGTATG	TTGGAATTCG
	351	GTTAAGGACT	AAGATTAGCC	CAaATTTTTT	TAAAATGTTA	ATTTTTATTG

401 tTTTATTGGT ATTGGCtCTG AAAATCGGGC AttCGGGTTT AAtCAAACCTT
451 TAA

This corresponds to the amino acid sequence <SEQ ID 74; ORF51>:

5 1 MAIITLYYSV NGILNVCAKA KNIQVVANNK NMVLFGLXX IIGGSTNAMS
 51 PILLIFLLSE TENKNRIVKS SNLCYLLAKI VQIYMLRDQY WLLNKSEYXL
 101 IFLLSVLSVI GLYVGIRLRT KISPNNFFKML IFIVLLVLAL KIGHSGLIK
 151 *

Further work revealed the complete nucleotide sequence <SEQ ID 75>:

10 1 ATGCAAGAAA TAATGCAATC TATCGTTTTT GTTGCTGCCG CAATACTGCA
 51 CGGAATTACA GGCATGGGAT TTCCGATGCT CGGTACAACC GCATTGGCTT
 101 TTATCATGCC ATTGTCTAAG GTTGTTGCCT TGGTGGCATT ACCAAGCCTG
 151 TTAATGAGCT TGTTGGTTCT ATGCAGCAAT AACAAAAAGG GTTTTTGGCA
 201 AGAGATTGTT TATTATTTAA AAACCTATAA ATTGCTTGCT ATCGGCAGCG
 251 TCGTTGGCAG CATTGTTGGG GTGAAGTTGC TTTTGATACT TCCAGTGTCT
15 301 TGGCTGCTTT TACTGATGGC AATCATTACA TTGTATTATT CTGTCAATGG
 351 TATTTTAAAT GTATGTGCAA AAGCAAAAAA TATTCAGTA GTTGCCAATA
 401 ATAAGAATAT GGTCTCTTTT GGGTTTTTGG CAGGCATCAT CGGCGGTTCA
 451 ACCAATGCCA TGTCTCCCAT ATTGTTAATA TTTTGCTTA GCGAAACAGA
 501 AAATAAAAAT CGTATCGTAA AATCAAGCAA TCTATGCTAT CTTTGGCGA
20 551 AAATTGTTCA AATATATATG CTAAGAGACC AGTATTGGTT ATTAAATAAG
 601 AGTGAATACG GTTTAATATT TTTACTGTCC GTATTGTCTG TTATTGGATT
 651 GTATGTTGGA ATTCGGTTAA GGACTAAGAT TAGCCCAAT TTTTAAAAA
 701 TGTTAATTTT TATTGTTTTA TTGGTATTGG CTCTGAAAAT CGGGCATTCG
 751 GGTTAATCA AACTTTAA

25 This corresponds to the amino acid sequence <SEQ ID 76; ORF51-1>:

30 1 MQEIMQSIVF VAAAILHGIT GMGFPM LGTT ALAFIMPLSK VVALVALPSL
 51 LMSLLVLC SN NKKGFWEIV YYLKTYKLLA IGSVGSILG VKLLILPVS
 101 WLLLLMAIIT LYYSVNGILN VCAKAKNIQV VANNKNMVL FFLAGIIGGS
 151 TNAMSPILLI FLLSETENKN RIVKSSNL CY LLAKIVQIYM LRDQYWLLNK
 201 SEYGLIFLLS VLSVIGLYVG IRLRTKISP N FFKMLIFIVL LVLALKHIGHS
 251 GLIKL*

Computer analysis of this amino acid sequence reveals three putative transmembrane domains. A corresponding ORF from strain A of *N.meningitidis* was also identified:

Homology with a predicted ORF from *N.meningitidis* (strain A)

35 ORF51 shows 96.7% identity over a 150aa overlap with an ORF (ORF51a) from strain A of *N.meningitidis*:

					10	20	30
orf51.pep					MAIITLYYSV	NGILNVCAKA	KNIQVVANNK
40 orf51a	YKLLAIGSVVGSILGVKLL	LILPVS	WLLLLMAIITLYYSV	NGILNVCAKA	KNIQVVANNK		
	80	90	100	110	120	130	
		40	50	60	70	80	90
orf51.pep		NMVLFGFLXXI	IIGGSTNAMSPILLI	FLLSETENKNR	IVKSSNL	CYLLAKIVQIY	MLRDQY
45 orf51a		NMVLFGFLAGI	IIGGSTNAMSPILLI	FLLSETENKNRI	AKSSNL	CYLLAKIVQIY	MLRDQY
		140	150	160	170	180	190
	100	110	120	130	140	150	
50 orf51.pep	WLLNKSEYXL	IFLLSVLSVIGLYVG	IRLRTKISP	NFFKMLIFIVL	LVLALKIGHSG	GLIKL	
orf51a	WLLNKSEYGL	IFLLSVLSVIGLYVG	IRLRTKISP	NFFKMLIFIVL	LVLALKIGYSGLIKL		
	200	210	220	230	240	250	

ORF51-1 and ORF51a show 99.2% identity in 255 aa overlap:

```

5  orf51a.pep  MQEIMQSIVFVAAAILHGITGMGFMLGTTALAFIMPLSKVVALVALPSLLMSLLVLCSN
   orf51-1    MQEIMQSIVFVAAAILHGITGMGFMLGTTALAFIMPLSKVVALVALPSLLMSLLVLCSN

10 orf51a.pep  NKKGFWQEIVYYLKYTKLLAIGSVVGSILGVKLLLLILPVSWLLLLLMAITLYYSVNGILN
   orf51-1    NKKGFWQEIVYYLKYTKLLAIGSVVGSILGVKLLLLILPVSWLLLLLMAITLYYSVNGILN

15 orf51a.pep  VCAKAKNIQVVANNKNMVLFGFLAGIIGGSTNAMSPILLIFLLSETENKNRIAKSSNLCY
   orf51-1    VCAKAKNIQVVANNKNMVLFGFLAGIIGGSTNAMSPILLIFLLSETENKNRIVKSSNLCY

20 orf51a.pep  LLAIVQIYMLRDQYWLLNKSEYGLIFLLSVLSVIGLYVGIRLRTKISPNEFKMLIFIVL
   orf51-1    LLAIVQIYMLRDQYWLLNKSEYGLIFLLSVLSVIGLYVGIRLRTKISPNEFKMLIFIVL

   orf51a.pep  LVLALKIGYSGLIKLX
   orf51-1    LVLALKIGHSGLIKLY

```

The complete length ORF51a nucleotide sequence <SEQ ID 77> is:

```

1  ATGCAAGAAA TAATGCAATC TATCGTTTTT GTTGCTGCCG CAATACTGCA
51 CGGAATTACA GGCATGGGAT TTCCGATGCT CGGTACAACC GCATTGGCCT
25 101 TTATCATGCC ATTGTCTAAG GTTGTGCCT TGGTGGCATT ACCAAGCCTG
   151 TTAATGAGCT TGTTGGTTCT ATGCAGCAAT AACAAAAAGG GTTTTTGGCA
   201 AGAGATTGTT TATTATTTAA AAACCTATAA ATTGCTTGCT ATCGGCAGCG
   251 TCGTTGGCAG CATTTTGGGG GTGAAGTTGC TTTTGATACT TCCAGTGTCT
   301 TGGCTGCTTT TACTGATGGC AATCATTACA TTGTATTATT CTGTCAATGG
   351 TATTTTAAAT GTATGTGCAA AAGCAAAAAA TATTCAAGTA GTTGCCAATA
30 401 ATAAGAATAT GGTCTTTTTT GGGTTTTTGG CAGGCATCAT CGGCGGTTCA
   451 ACCAATGCCA TGTCTCCCAT ATTGTTAATA TTTTGCTTCA GCGAAACAGA
   501 GAATAAAAAA CGTATCGCAA AATCAAGCAA TCTATGCTAT CTTTGGCAA
   551 AAATTGTTCA AATATATATG CTAAGAGACC AGTATTGGTT ATTAAATAAG
35 601 AGTGAATACG GTTAAATATT TTTACTGTCC GTATTGTCTG TTATTGGATT
   651 GTATGTTGGA ATTCGGTTAA GGACTAAGAT TAGCCCAAAT TTTTAAAAA
   701 TGTTAATTTT TATTGTTTTA TTGGTATTGG CTCTGAAAAT CGGGTATTCA
   751 GGTTTAATCA AACTTTAA

```

This encodes a protein having amino acid sequence <SEQ ID 78>:

```

40 1  MQEIMQSIVF VAAAILHGIT GMGFMLGTT ALAFIMPLSK VVALVALPSL
   51  LMSLLVLCSN NKKGFWQEIV YYLKYTKLLA IGSVVGSILG VKLLLLILPV
   101 WLLLLLMAIT LYYSVNGILN VCAKAKNIQV VANNKNMVL FFLAGIIGGS
   151 TNAMSPILLI FLLSETENKN RIAKSSNLCY LLAIVQIYM LRDQYWLLNK
   201 SEYGLIFLLS VLSVIGLYVG IRLRTKISPN FFKMLIFIVL LVLALKIGYS
   251 GLIKL*

```

45 Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 18

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 79>

```

50 1  ATGAGACATA TGAAAATACA AAATTATTTA CTAGTATTTA TAGTTTACAA
   51  TATAGCCTTG ATAGTAATTA ATATAGTGTT TGGTTATTTT GTTTTCTAT
   101 TTGATTTTTT TGCGTTTTTG TTTTGTGCAA ACGTCTTCTT TGCTGTAAAT
   151 TTATTATTTT TAGAAAAAAC CATAAAAAAC AAATTATTGT TTTTATTGCC
   201 GATTTCTATT ATTATATGGA TGGTAATTCA TATTAGTATG ATAAATATAA
55 251 AATTTTATAA ATTTGAGCAT CAAATAAAGG AACAAATAT ATCCTCGATT
   301 ACTGGGGTGA TAAACCACA TGATAGTTAT AATTATGTTT ATGACTCAAA

```

```

351 TGGATATGCT AAATTTAAAG ATAATCATAG ATATGGTAGG GTAATTAGAG
401 AAACACCTTA TATTGATGTA GTTGCATCTG ATGTTAAAAA TAAATCCATA
451 AGATTAAGCT TGGTTTGTGG TATTCATTCA TATGCTCCAT GTGCCAATTT
501 TATAAAATTT GTCAGG..

```

5 This corresponds to the amino acid sequence <SEQ ID 80; ORF82>:

```

1 MRHMKIQNYL LVFIVLHIAL IVINIVFGYF VFLDFFAFL FFANVFLAVN
51 LFLFLEKNIKN KLLFLLPISI IIWMVIHISM INIKFYKFEH QIKEQNISSI
101 TGVIKPHDSY NYVYDSNGYA KLKDNHRYGR VIRETPYIDV VASDVKNKSI
151 RLSLVCGIHS YAPCANFIKF VR..

```

10 Further work revealed the complete nucleotide sequence <SEQ ID 81>:

```

1 ATGAGACATA TGAAAAATAA AAATTATTTA CTAGTATTTA TAGTTTTACA
51 TATAGCCTTG ATAGTAATTA ATATAGTGTT TGGTTATTTT GTTTTTCTAT
101 TTGATTTTTT TCGGTTTTTG TTTTGTGCAA ACGTCTTTCT TGCTGTAAAT
151 TTATTATTTT TAGAAAAAAA CATAAAAAAC AAATTATTGT TTTTATTGCC
15 201 GATTTCTATT ATTATATGGA TGGTAATTCA TATTAGTATG ATAAATATAA
251 AATTTTATAA ATTTGAGCAT CAAATAAAGG AACAAAATAT ATCCTCGATT
301 ACTGGGGTGA TAAAACCACA TGATAGTTAT AATTATGTTT ATGACTCAAA
351 TGGATATGCT AAATTTAAAG ATAATCATAG ATATGGTAGG GTAATTAGAG
401 AAACACCTTA TATTGATGTA GTTGCATCTG ATGTTAAAAA TAAATCCATA
20 451 AGATTAAGCT TGGTTTGTGG TATTCATTCA TATGCTCCAT GTGCCAATTT
501 TATAAAATTT GCAAAAAAAC CTGTTAAAAT TTATTTTAT AATCAACCTC
551 AAGGAGATTT TATAGATAAT GTAATATTG AAATTAATGA TGGAAACAAA
601 AGTTTGTACT TGTTAGATAA GTATAAAACA TTTTCTTCTT TTGAAAACAG
651 TGTTTGTATC GTATTAATTA TTTTATATTT AAAATTTAAT TTGCTTTTAT
25 701 ATAGGACTTA CTTCAATGAG TTGGAATAG

```

This corresponds to the amino acid sequence <SEQ ID 82; ORF82-1>:

```

1 MRHMKKNKNYL LVFIVLHIAL IVINIVFGYF VFLDFFAFL FFANVFLAVN
51 LFLFLEKNIKN KLLFLLPISI IIWMVIHISM INIKFYKFEH QIKEQNISSI
101 TGVIKPHDSY NYVYDSNGYA KLKDNHRYGR VIRETPYIDV VASDVKNKSI
30 151 RLSLVCGIHS YAPCANFIKF AKKPVKIYFY NQPGDFIDN VIFEINDGNK
201 SLYLLDKYKT FFLIENSVC I VLIILYLKFN LLYRITYFNE LE*

```

Computer analysis of this amino acid sequence reveals a predicted leader peptide.

A corresponding ORF from strain A of *N.meningitidis* was also identified:

Homology with a predicted ORF from *N.meningitidis* (strain A)

35 ORF82 shows 97.1% identity over a 172aa overlap with an ORF (ORF82a) from strain A of *N.meningitidis*:

		10	20	30	40	50	60
orf82.pep		MRHMKIQNYLLVFIVLHIALIVINIVFGYFVFLDFFAFLFFANVFLAVNLLFLEKNIKN					
40 orf82a		MRHMKKNKNYLLVFIVLHITLLVINIVFGYFVFLDFFAFLFFANVFLAVNLLFLEKNIKN					
		10	20	30	40	50	60
		70	80	90	100	110	120
orf82.pep		KLLFLLPISIIIIWMVIHISM INIKFYKFEHQIKEQNISITGVIKPHDSYNYVYDSNGYA					
45 orf82a		KLLFLLPISIIIIWMVIHISM INIKFYKFEHQIKEQNISITGVIKPHDSYNYVYDSNGYA					
		70	80	90	100	110	120
		130	140	150	160	170	
50 orf82.pep		KLKDNHRYGRVIRETPYIDVVASDVKNKSIRLSLVCGIHSYAPCANFIKFVR					
orf82a		KLKDNHRYGRVIRETPYIDVVASDVKNKSIRLSLVCGIHSYAPCANFIKFAKKPVKIYFY					
		130	140	150	160	170	180

ORF82a and ORF82-1 show 99.2% identity in 242 aa overlap:

```

5  orf82a.pep  MRHMKNKNYLLVFIVLHITLIVINIVFGYFVFLDFFAFLFFANVFLAVNLLFLEKNIKN
   orf82-1    MRHMKNKNYLLVFIVLHIALIVINIVFGYFVFLDFFAFLFFANVFLAVNLLFLEKNIKN

10 orf82a.pep  KLLFLLPISIIWMVIHISMINIKFYKFEHQIKEQNISSITGVIKPHDSYNYVYDSNGYA
   orf82-1    KLLFLLPISIIWMVIHISMINIKFYKFEHQIKEQNISSITGVIKPHDSYNYVYDSNGYA

15 orf82a.pep  KLKDNHRYGRVIRETPYIDVVASDVKNKSIRLSLVCGIHSYAPCANFIKFAKKPVKIYFY
   orf82-1    KLKDNHRYGRVIRETPYIDVVASDVKNKSIRLSLVCGIHSYAPCANFIKFAKKPVKIYFY

   orf82a.pep  NQPQGD FIDNVIFEINDGKKS LYL LDKYKTFFLIENSVCIVLIILYLKFN LLLYRTYFNE
   orf82-1    NQPQGD FIDNVIFEINDGKKS LYL LDKYKTFFLIENSVCIVLIILYLKFN LLLYRTYFNE

20 orf82a.pep  LEX
   orf82-1    LEX

```

The complete length ORF82a nucleotide sequence <SEQ ID 83> is:

```

25 1  ATGAGACATA TGAAAAATAA AAATTATTTA CTAGTATTTA TAGTTTAC
   51 TATAACCTTG ATAGTAATTA ATATAGTGTT TGGTTATTTT GTTTTCTAT
  101 TTGATTTTTT TCGGTTTTTG TTTTTCGCAA ACGTCTTTCT TGCTGTAAAT
  151 TTATTATTTT TAGAAAAAAA CATAAAAAAC AAATTATTGT TTTTATTGCC
  201 GATTTCTATT ATTATATGGA TGGTAATTCA TATTAGTATG ATAAATATAA
  251 AATTTTATAA ATTTGAGCAT CAAATAAAGG AACAAAATAT ATCCTCGATT
  301 ACTGGGGTGA TAAAACCACA TGATAGTTAT AATTATGTTT ATGACTCAAA
  351 TGGATATGCT AAATTAAAAG ATAATCATAG ATATGGTAGG GTAATTAGAG
  401 AAACACCTTA TATTGATGTA GTTGCATCTG ATGTTAAAAA TAAATCCATA
  451 AGATTAAGCT TGGTTTGTGG TATTCATTCA TATGCTCCAT GTGCCAATTT
  501 TATAAAATTT GCAAAAAAAC CTGTTAAAAT TTATTTTAT AATCAACCTC
  551 AAGGAGATT TATAGATAAT GTAATATTG AAATTAATGA TGGAAAAAAA
  601 AGTTTGTACT TGTTAGATAA GTATAAAACA TTTTTCCTTA TTGAAAAACAG
  651 TGTTTGTATC GTATTAATTA TTTTATATTT AAAATTTAAT TTGCTTTTAT
  701 ATAGGACTTA CTTCAATGAG TTGGAATAG

```

This encodes a protein having amino acid sequence <SEQ ID 84>:

```

40 1  MRHMKNKNYL LVFIVLHITL IVINIVFGYF VFLDFFAFL FFANVFLAVN
   51 LLFLEKNIKN KLLFLLPISI IIWMVIHISM INIKFYKFEH QIKEQNISSI
  101 TGVIKPHDSY NYVYDSNGYA KLKDNHRYGR VIRETPYIDV VASDVKNKSI
  151 RLSLVCGIHS YAPCANFIK AKKPVKIYFY NQPQGD FIDN VIFEINDGKK
  201 SLYLLDKYKT FFLIENSVC I VLIILYLKFN LLLYRTYFNE LE*

```

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

Example 19

The following partial DNA sequence was identified in *N.meningitidis* <SEQ ID 85>

```

50 1  ..ACCCCAACA GCGTGACCGT CTTGCCGTCT TTCGGCGGAT TCGGGCGTAC
   51 CGGCGCGACC ATCAATGCAG CAGGCGGGGT CGGCATGACT GCCTTTTCGA
  101 CAACCTTAAT TTCCGTAGCC GAGGGCGCGG TTGTAGAGCT GCAGGCCGTG
  151 AGAGCCAAAG CCGTCAATGC AACCGCCGCT TGCATTTTTA CGGTCTTGAG
  201 TAAGGACATT TTCGATTTC TTTTATTTT CCGTTTTTCA ACGGTGACTT
  251 TCCGCCTGTA TTTTCGCCAA AGCCATGCCG ACAGCGTGCG CCTTGACTTC
  301 ATATTAAAAA GCTTCCGCGC GTGCCAGTTC CAGTTCGCGC GCATAGTTTT
  351 GAGCCGACAA CAGCAGGCT TGCGCCTTGT CGCGCTCCAT CTTGTCGATG

```

```

401 ACCGCCTGCA GCTTCGCAAA TGCCGACTTG TAGCCTTGAT GGTGCGACAC
451 AGCCAAGCCC GTGCCGACAA GCGCGATAAT GGCAATCGGT TGCCAGTAAT
501 TCGCCAGCAG TTCACGAGA TTCATTCTCG ACCTCCTGAC GCTTCACGCT
551 GA

```

5 This corresponds to the amino acid sequence <SEQ ID 86; ORF124>:

```

1 ..TPNSVTVLPS FGGFGRTGAT INAAGGVGMT AFSTTLISVA EGAVVELQAV
51 RAKAVNATAA CIFTVLSKDI FDFLFIFRFQ TADFRLYFRQ SHADSVRLDF
101 IFKSFRACQF QFARIVLSRQ QQGLRLVALH LVDDRLLQLRK CRLVALMVRH
151 SQARADKRDN GNRLPVIRQQ FHEIHSRPPD ASR*

```

10 Computer analysis of this amino acid sequence predicts a transmembrane domain.

Further work revealed the complete nucleotide sequence <SEQ ID 87>:

```

1 ATGACTGCCT TTTTCGACAAC CTTAATTTCC GTAGCCGAGG GCGCGGTTGT
51 AGAGCTGCAG GCCGTGAGAG CCAAAGCCGT CAATGCAACC GCCGCTTGCA
101 TTTTACGGT CTTGAGTAAG GACATTTTCG ATTTCTTTT TATTTTCCGT
151 TTTCAGACGG CTGACTTCCG CCTGTTTTTT CGCCAAAGCC ATGCCGACAG
201 CGTGCGCCTT GACTTCATAT TTTTGTAGCTT CCGCGCGTGC CAGTTCAGT
251 TCGCGCGCAT AGTTTTGAGC CGACAACAGC AGGGCTTGCG CCTGTGCGCG
301 CTCCATCTTG TCGATGACCG CCTGCTGCTT CGCAAATGCC GACTTGTAGC
351 CTTGATGGTG CGACACAGCC AAGCCCGTGC CGACAAGCGC GATAATGGCA
401 ATCGGTTGCC AGTTATTCGC CAGCAGTTTC ACGAGATTCA TTCTCGACCT
451 CCTGACGCTT CACGCTGA

```

This corresponds to the amino acid sequence <SEQ ID 88; ORF124-1>:

```

1 MTAFTTLIS VAEGAVVELQ AVRAKAVNAT AACIFTVLSK DIFDFLFIER
51 FQTADFRLEF RQSHADSVRL DFIFFSFRAC QFQFARIVLS RQQQGLRLVA
101 LHLVDDRLLL RKRLVALMV RHSQARADKR DNGNRLPVIR QFHEIHSRP
151 PDASR*

```

A corresponding ORF from strain A of *N.meningitidis* was also identified:

Homology with a predicted ORF from *N.meningitidis* (strain A)

ORF124 shows 87.5% identity over a 152aa overlap with an ORF (ORF124a) from strain A of *N.*

30 *meningitidis*:

```

10 20 30 40 50 60
orf124.pep TPNSVTVLPSFGGFGRTGATINAAGGVGMTAFSTTLISVAEGAVVELQAVRAKAVNATAA
orf124a MTAFTTLISVAEGALVELQAVMAKAVNTTAA
10 20 30
70 80 90 100 110 120
orf124.pep CIFTVLSKDIFDFLFIFRFQTADFRLYFRQSHADSVRLDFIFKSFRACQFQFARIVLSRQ
orf124a CIFTVLSKDIFDFLFIFRFQTADFRLLFFRQSHADGVRLDFIFFSFRTRLFQFAGVVLRSRQ
40 40 50 60 70 80 90
130 140 150 160 170 180
orf124.pep QQGLRLVALHLLVDDRLLQLRKRLVALMVRHRSQARADKRDNGNRLPVIRQQFHEIHSRPPD
orf124a QQGLRLVALHFLNDRLLLRKSRLVALMVRHRRQTRADKRDDGNRLPVIRQQFHEIHSRPPD
100 110 120 130 140 150
orf124.pep ASRX
orf124a :
50 orf124a VX

```

ORF124a and ORF124-1 show 89.5% identity in 152 aa overlap:

```

    orf124-1.pep  MTAfstTLISVAEGAVVELQAVRAKAVNATAACIFTVLskDIFdFLFIfrFQTADfRLFF
    orf124a      MTAfstTLISVAEGALVELQAVMAKAVNTTAACIFTVLskDIFdFLFIfrFQTADfRLFF
5   orf124-1.pep  RQSHADSVRLDFIffSFRACQFQfARIVLSRQQQGLRLVALHLVDDRLLLRKCRlVALMV
    orf124a      RQSHADGVRLDFIffSFRTRLfQfAGVVLSRQQQGLRLVALHFLNDRLLLRKSRLVALMV
10  orf124-1.pep  RhsQARADKRDNGNRLPVIRQQfHEIHSRPPDASRX
    orf124a      RHRQTRADKRDDGNRLPVIRQQfHEIHSRPPDVX

```

The complete length ORF124a nucleotide sequence <SEQ ID 89> is:

```

1   ATGACCGCCT TTTGACAAC CTTAATTTCC GTAGCCGAGG GCGCGCTTGT
15  51  AGAGCTGCAA GCCGTGATGG CCAAAGCCGT CAATACAACC GCCGCCTGCA
    101  TTTTACGGT CTTGAGTAAG GACATTTTCG ATTCCTTTT TATTTCCGT
    151  TTTcAGACGG CTGACTTCCG CCTGTTTTT CGCCAAAGCC ATGCCGACGG
    201  CGTGCGCCTT GACTTCATAT TTTTtagCTT CCGCACGCGC CTGTTCCAGT
    251  TCGCGGGCGT AGTTTTGAGC CGACAACAGC AGGGCTTGCG CCTTGTGCGG
20  301  CTTcATTTTC TCAATGACCG CCTGCTGCTT CGCAAAGCC GACTTGTAGC
    351  CTTGATGGTG CGACACCGCC AAACCCGTGC CGACAAGCGC GATGATGGCA
    401  ATCGGTTGCC AGTTATTGCG CAGCAGTTTC ACGAGATTCA TTCTCGACCT
    451  CCTGACGTTT GA

```

This encodes a protein having amino acid sequence <SEQ ID 90>:

```

25  1   MTAfstTLIS VAEGALVELQ AVMAKAVNTT AACIFTVLsk DIFdFLFIfr
    51  FQTADfRLFF RQSHADGVRL DFIffSFRTR LFQfAGVVLS RQQQGLRLVA
    101  LHFLNDRLLL RKSRLVALMV RHRQTRADKR DDGNRLPVIR QQfHEIHSRP
    151  PDV*

```

ORF124-1 was amplified as described above. Figure 7 shows plots of hydrophilicity, antigenic index, and AMPHI regions for ORF124-1.

Based on this analysis, it is predicted that this protein from *N.meningitidis*, and its epitopes, could be useful antigens for vaccines or diagnostics.

It will be appreciated that the invention has been described by means of example only, and that modifications may be made whilst remaining within the spirit and scope of the invention.

TABLE I – PCR primers

ORF	Primer	Sequence	Restriction sites
ORF 38	Forward	CGCGGATCCCATATG-TCGCCGAAAATTCCGA	BamHI-NdeI XhoI
	Reverse	CCCGCTCGAG-TTTTGCCGCGTTAAAAGC	
ORF 40	Forward	CGCGGATCCCATATG-ACCGTGAAGACCGCC	BamHI-NdeI XhoI
	Reverse	CCCGCTCGAG-CCACTGATAACCGACAGA	
ORF 41	Forward	CGCGGATCCCATATG-TATTTGAAACAGCTCCAAG	BamHI-NdeI XhoI
	Reverse	CCCGCTCGAG-TTCTGGGTGAATGTTA	
ORF 44	Forward	GCGGATCCCATATG-GGCACGGACAACCCC	BamHI-NdeI XhoI
	Reverse	CCCGCTCGAG-ACGTGGGGAACAGTCT	
ORF 51	Forward	GCGGATCCCATATG-AAAAATATTCAAGTAGTTGC	BamHI-NdeI XhoI
	Reverse	CCCGCTCGAG-AAGTTTGATTAAACCCG	
ORF 52	Forward	CGCGGATCCCATATG-TGCCAACCGCAATCCG	BamHI-NdeI XhoI
	Reverse	CCCGCTCGAG-TTTTTCAGCTCCGGCA	
ORF 56	Forward	GCGGATCCCATATG-GTTATCGGAATATTACTCG	BamHI-NdeI XhoI
	Reverse	CCCGCTCGAG-GGCTGCAGAAGCTGG	
ORF 69	Forward	CGCGGATCCCATATG-CGGACGTGGTTGGTTTT	BamHI-NdeI XhoI
	Reverse	CCCGCTCGAG-ATATCTTCCGTTTTTTTCAC	
ORF 82	Forward	CGCGGATCCGCTAGC-GTAAATTTATTATTTTAGAA	BamHI-NheI XhoI
	Reverse	CCCGCTCGAG-TTCCAATCATTGAAGTA	
ORF 114	Forward	CGCGGATCCCATATG-AATAAAGGTTTACATCGCAT	BamHI-NheI XhoI
	Reverse	CCCGCTCGAG-AATCGCTGCACCGGCT	
ORF 124	Forward	CGCGGATCCCATATG-ACTGCCTTTTCGACA	BamHI-NheI XhoI
	Reverse	CCCGCTCGAG-GCGTGAAGCGTCAGGA	

TABLE II – Cloning, expression and purification




ORF	PCR/cloning	His-fusion expression	GST-fusion expression	Purification
orf 38	+	+	+	His-fusion
orf 40	+	+	+	His-fusion
orf 41	+	n.d.	n.d.	
orf 44	+	+	+	His-fusion
orf 51	+	n.d.	n.d.	
orf 52	+	n.d.	+	GST-fusion
orf 56	+	n.d.	n.d.	
orf 69	+	n.d.	n.d.	
orf 82	+	n.d.	n.d.	
orf 114	+	n.d.	+	GST-fusion
orf 124	+	n.d.	n.d.	

CLAIMS



1. A protein comprising an amino acid sequence selected from the group consisting of SEQ IDs 2, 4, and 6.
2. A nucleic acid molecule which encodes a protein according to claim 1.
- 5 3. A nucleic acid molecule according to claim 2, comprising a nucleotide sequence selected from the group consisting of SEQ IDs 1, 3, and 5.
4. A protein comprising an amino acid sequence selected from the group consisting of SEQ IDs 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, and 90.
- 10 5. A protein having 50% or greater sequence identity to a protein according to claim 4.
6. A protein comprising a fragment of an amino acid sequence selected from the group consisting of SEQ IDs 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, and 90.
7. An antibody which binds to a protein according to any one of claims 4 to 6.
- 15 8. A nucleic acid molecule which encodes a protein according to any one of claims 4 to 6.
9. A nucleic acid molecule according to claim 8, comprising a nucleotide sequence selected from the group consisting of SEQ IDs 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, and 89.
- 20 10. A nucleic acid molecule comprising a fragment of a nucleotide sequence selected from the group consisting of SEQ IDs 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, 63, 65, 67, 69, 71, 73, 75, 77, 79, 81, 83, 85, 87, and 89.
11. A nucleic acid molecule comprising a nucleotide sequence complementary to a nucleic acid molecule according to any one of claims 8 to 10.

12. A nucleic acid molecule comprising a nucleotide sequences having 50% or greater sequence identity to a nucleic acid molecule according to any one of claims 8 to 11.
13. A nucleic acid molecule which can hybridise to a nucleic acid molecule according to any one of claims 8 to 12 under high stringency conditions.
- 5 14. A composition comprising a protein, a nucleic acid molecule, or an antibody according to any preceding claim.
15. A composition according to claim 14 being a vaccine composition or a diagnostic composition.
16. A composition according to claim 14 or claim 15 for use as a pharmaceutical.
- 10 17. The use of a composition according to claim 14 in the manufacture of a medicament for the treatment or prevention of infection due to Neisserial bacteria, particularly *Neisseria meningitidis*.


1/11

FIG. 1A**FIG. 1B****FIG. 4A**





2/11

**FIG. 1C****FIG. 1D**

3/11



4/11

FIG. 2A**FIG. 2B****FIG. 2C****FIG. 2D**

5/11







FIG. 2E

6/11

FIG. 3A**FIG. 3B****FIG. 3C**

7/11





FIG. 3D


8/11

FIG. 4B




9/11

FIG. 5



10/11

FIG. 6



11/11

FIG. 7

